

# Range Plant Community Types and Carrying Capacity for the Subalpine and Alpine Subregions



**Alberta**

**Sustainable Resource Development**



**RANGE PLANT COMMUNITY TYPES AND CARRYING CAPACITY  
FOR THE SUBALPINE AND ALPINE SUBREGIONS**

**Second approximation**

**(Please note this edition is a revision of the 1<sup>st</sup> approximation of the Range Plant  
Community types and carrying capacity for the Subalpine and Alpine subregions. Pub. No.  
T/438)**

**2003**

**Prepared by**

**Michael G. Willoughby**

**and**

**Michael J. Alexander**

Edmonton  
2003

Sustainable Resource Development  
Public Lands Division

Pub. No.: T/034  
ISBN: 0-7785-2342-X

For copies of this report contact:

Michael Willoughby  
Rm 200 J.G. O' Donoghue Bldg.  
7000 113 th Str.  
Edmonton, Alta.  
T6H 5T6  
(780) 422-4598  
[mike.willoughby@gov.ab.ca](mailto:mike.willoughby@gov.ab.ca)

## Table of contents

	Page
Introduction	1
Climate of Subalpine and Alpine subregions	1
Methods	4
How to use guide	4
Results	26
Subalpine (Grassland and Shrubland ecology)	27
Central and Northern Foothills areas	28
Ecology of ecodistricts	29
Key to grasslands in the Foothills	33
Key to shrublands in the Foothills	34
Community types	
Grasslands	
SACFA1. Water sedge-Beaked sedge meadows	35
SACFA2. Tufted bulrush	36
SACFA3. Sedge-Cottongrass	37
SACFA4. Tufted hairgrass-Sedge	38
SACFA5. Sedge-Tufted hairgrass	39
SACFA6. Sedge-Rocky Mtn. fescue-Alpine timothy	40
SACFA7. Sedge-Slender wheatgrass-Fringed brome/Forb	41
SACFA8. California oatgrass-Sedge	42
SACFA9. Rough fescue-Hairy wildrye-Sedge	43
SACFA10. Sedge-Hairy wildrye	44
SACFA11. Blunt sedge-Junegrass/Bearberry	45
SACFA12. Fringed sage/Sedge-Junegrass	46
SACFA13. Sedge-Bog sedge-Tufted hairgrass	47
SACFA14. White Mtn. avens/Bog sedge	48
SACFA15. Creeping red fescue-Sedge	49
SACFA16. Kentucky bluegrass-Sedge/Dandelion	50
SACFA17. Fireweed-Meadow rue/Sedge-Hairy wildrye	51
Shrublands	
SACFB1. Willow-Bog birch/Water sedge	52
SACFB2. Willow/Horsetail	53

SACFB3. Willow/Graceful sedge	54
SACFB4. Willow-Bog birch/Tufted hairgrass	55
SACFB5. Willow-Bog birch/Clover-Dandelion	56
SACFB6. Willow-Bog birch/California oatgrass	57
SACFB7. Willow-Bog birch/Hairy wildrye	58
SACFB8. Bog birch/Bog sedge-Sedge	59
SACFB9. Bog birch-Willow/Rough fescue	60
SACFB10. Bog birch/Rough fescue-Bog sedge	61
SACFB11. Willow/Fringed brome-Sedge	62
Central and Northern Rocky Mountain areas	63
Ecology of ecodistricts	64
Key to the grasslands within the Mountain ecodistricts	68
Key to the shrublands within the Mountain ecodistricts	69
Community types	
Grasslands	
SACMA1. Bog sedge-California oatgrass	70
SACMA2. Forb meadows	71
SACMA3. Shrubby cinquefoil/Hairy wildrye	72
SACMA4. Bearberry-Juniper	73
SACMA5. Junegrass-Hairy wildrye-Brome	74
SACMA6. Hairy wildrye/Bearberry-Juniper	75
SACMA7. Northern wheatgrass	76
SACMA8. Alpine bluegrass	77
SACMA9. Yellow mountain avens	78
Shrublands	
SACMB1. Willow/Water sedge	79
SACMB2. Willow-Bog birch/Sedge	80
SACMB3. Willow-Bog birch/Hairy wildrye	81
SACMB4. Willow-Bog birch/Bog sedge	82
SACMB5. Bog birch/Juniper	83
SACMB6. Willow/Forb	84
SACMB7. Grouseberry-Juniper	85
SACMB8. Subalpine fir	86
Southern Rocky Mountains	87
Native grass and shrubland ecology	88
Key to grasslands and shrublands	116
Community types	
Grasslands	
North of Blairmore	

SASMA1. Water sedge	118
SASMA1a. Beaked sedge-Alpine foxtail-Tufted hairgrass	119
SASMA2. Rough fescue-Sedge	120
SASMA3. Rough fescue-Hairy wildrye	121
SASMA3a Hairy wildrye-Rough fescue-Sedge	122
SASMA4. Sedge-Hairy wildrye-Slender wheatgrass	123
SASMA5. Kentucky bluegrass/Dandelion	124
SASMA6. Yellow mountain avens	125
SASMA7. Tufted hairgrass-Sedge	126
SASMA7a Marsh reedgrass/Cow parsnip	127
South of Blairmore	
SASMA8. Rough fescue-Idaho fescue-Parry oatgrass	128
SASMA9. Rough fescue-Sedge/Bearberry	129
SASMA10. Parry oatgrass-Rough fescue-Sedge	130
SASMA11. Sedge/Bearberry	131
SASMA12. Silverberry-Rose	132
SASMA13. Fescue-Junegrass/Early yellow locoweed	133
SASMA14. White Mountain avens	134
SASMA15. Pinegrass-Hairy wildrye/Strawberry	135
SASMA16. Forb meadow	136
Shrublands	
SASMB1. Willow/Sedge	137
SASMB2. Willow/Richardson needlegrass	138
SASMB3. Whitebark pine	139
SASMB4. Willow-Bog birch/Rough fescue-Kentucky bluegrass	140
SASMB5. Willow/Marsh reedgrass	141
Grazing modified	
SASMC1. Parry oatgrass-Rough fescue-Kentucky bluegrass	142
SASMC2. Parry oatgrass-Kentucky bluegrass-Sedge	143
SASMC3. Meadow foxtail-Kentucky bluegrass	144
SASMC4. Fringed sage/Kentucky bluegrass-Parry oatgrass	145
SASMC5. Rough fescue-Kentucky bluegrass	146
SASMC6. Kentucky bluegrass-Rough fescue	147
SASMC7. Timothy-Slender wheatgrass/Fireweed	148
SASMC8. Marsh reedgrass-Timothy/Cow parsnip	149
SASMC9. Idaho fescue-Rough fescue/Bearberry	150
Subalpine Forest Ecology	152
Deciduous	
SASMD1 Pb/Silverberry	154

SASMD2. Aw/Rose/Pinegrass	155
SASMD3. Aw/Fireweed/Meadow foxtail	156
SASMD4. Aw/Rose/Canada bluegrass	157
SASMD5. Aw-Pb/Cow parsnip	158
SASMD6. Aw-Pb/Cow parsnip/Timothy	159
<b>Conifer</b>	
SASME1. Pl/Juniper	160
SASME2. Pl/Pinegrass	161
SASME3. Pl-Se/Moss	162
SASME4. Se-Aw/Alder/Hairy wildrye	163
<b>Alpine subregion</b>	
Ecology of Alpine communities	165
Community types	
ALPA1. Bog sedge	167
ALPA2. White mountain avens	168
ALPA3. Mountain heather	169
ALPA4. Blackening sedge	170
ALPA5. Simple bog sedge	171
ALPA6. Arctic willow (low willow types)	172
Literature cited	173

### **List of Figures**

Figure 1. Overview of native shrub and grassland complex in the Central and Northern Foothills	28
Figure 2. Grassland and shrubland community types in the landscape of the Central and Northern Rocky Mountain and Foothills areas of the Subalpine subregion	32
Figure 3. Overview of native shrub and grassland complex in the Central and Northern Rocky Mountains	63
Figure 4. Sequence of plant communities in the upper Subalpine and Alpine subregions of Alberta	67
Figure 5. Overview of native shrub and grassland complex in the Southern Rocky Mountains	87
Figure 6. Grassland and shrubland community types in the landscape of the Montane and Subalpine subregions of southern Alberta	115
Figure 7. Aspen/Rose/Pinegrass dominated community type in the southern zone of the Subalpine subregion	151
Figure 8. Overview of native shrub and grassland complex in the Alpine	



### List of Tables

Table 1. Ecosites, ecosite phases and plant community types for the Subalpine subregion of West-Central Alberta(adapted from Beckingham et al. 1996).	6
Table 2. Native grass and shrublands of the Central and Northern Foothills	30
Table 3. Native grass and shrublands of the Central and Northern Rocky Mountains	65
Table 4. Ecosites, ecosite phases and plant community types for the Subalpine subregion of Southern Alberta(adapted from Archibald and Klappstein 1996).	89
Table 5. Native grass and shrublands of the Southern Rocky Mountains	113
Table 6. Deciduous and conifer communities for the Southern Rocky Mtn. ecodistrict of the Subalpine subregion	153
Table 7. Native grasslands and shrublands of the Alpine subregion	166

### List of Maps

Map 1. Location of Alpine and Subalpine subregions in Alberta	3
---	---

## Abstract

The Subalpine subregion is a Rocky Mountain altitudinal vegetation zone extending from an elevation of 1575 m to 2175 m at its southernmost occurrence and 1365 m to 2000 m near Grande Cache (Strong 1992). The valley bottoms of the subalpine are extensively utilized for recreational horseback riding and commercial trail riding operations. As a result many of the grass and shrublands around the back country campsites are extensively utilized by horses. This utilization has an impact on the vegetation which can be detrimental to the wildlife populations in the area. Despite the importance of these vegetation types for wildlife and backcountry horse use there is little information on their ecology. The lack of information makes it difficult to development management prescriptions for multiple use. The Alpine subregion which occurs above the upper climatic forest-line in the Rocky Mountains (Strong 1992) overlaps with many of the subalpine community types and therefore can be important locally for wildlife. As a result this guide was developed for the Subalpine and Alpine subregions in order to provide a framework that will easily group the vegetative community types. It is hoped this classification system can be used by field staff to assess the ecology of the sites and develop management prescriptions on lands within each region. This guide represents the analysis of 651 plots described in the Subalpine subregion and 134 grass and shrubland plots described in the Alpine subregion. In the Subalpine subregion it was found there was distinct differences between the grassland and shrubland community types between the Central and Northern Foothills (West of Rocky Mountain house and Hinton), the Central and Northern Rocky Mountains (Banff and Jasper National Parks) and the Southern Rocky Mountains (southwest of Calgary). As a result the Subalpine was prestratified into 3 subdivisions. These types are split into:

### Subalpine

#### Central and Northern Foothills

- A. Native grasslands 17 community types
- B. Native shrublands 11 community types

#### Central and Northern Rocky Mountains

- A Native grasslands 9 community types
- B. Native shrublands 8 community types

#### Southern Rocky Mountains

- A. Native grasslands 19 community types
- B. Native shrublands 5 community types
- C. Grazed grasslands 9 community types
- D. Deciduous 6 community types
- E. Conifer 4 community types

### Alpine

- A. Native grasslands and shrublands 6 community types

The dominant plant species, canopy cover, environmental conditions, response to grazing, forage production and carrying capacity are outlined for each type.

## Acknowledgements

The creation of this report would not be possible without the data collected in other projects. We would like to acknowledge Parks Canada for allowing us to use data from the Ecological Land Classification of Banff and Jasper National Parks (Holland and Coen 1982). Much of the grass and shrubland vegetation data collected by Ian Corns and Peter Achuff for this project were incorporated into this guide. We would also like to acknowledge the Alberta Conservation Association and Corporate Management Service. They provided funding for the study of bighorn sheep winter range in Southern Alberta and backcountry horse use and elk carrying capacity in the Panther Corners. The vegetation data from these studies were also included in this guide. Finally appreciation and thanks go out to all members and former members of Land and Forest Service who were involved in data collection.



## **Introduction**

The province of Alberta is covered by a broad spectrum of vegetation regions from prairie in the South, to alpine vegetation in the mountains and dense forests in the Central and Northern part of the province. These broad vegetation regions have been classified into 6 regions and 20 subregions (Dept. of Environmental Protection 1994). Each of the 20 subregions consists of groups of plant communities which are influenced by environmental conditions and human impacts. Intensive management of these regions requires the ability to recognize the vegetative communities that have similar productivities and respond to disturbance in the same way.

The purpose of this guide was to develop a framework that would easily group the grassland and shrubland community types in the Subalpine and Alpine subregions of the province. Initially the guide was developed to provide a classification system that could be used by the field staff to assess carrying capacity and evaluate range condition on back country areas within the foothills of the Subalpine subregion, but it was realized that these community types were only transitional to subalpine and alpine communities described by Ogilvie (1969) and Corns and Achuff (1982). As a result, their original data was reclassified and added to the more recent data to form a more detailed classification.

It is hoped this classification system can be used by field staff to assess the ecology of the sites and develop management prescriptions on lands within the region. This guide supplements the work done by Beckingham et al. (1996), Beckingham and Archibald (1996) and Archibald et al. (1996) on the forested community types in the Subalpine subregion. Their guides are a good description of the forested community types found within the subregion, but it does not include forage production values and carrying capacities. It also does not provide a description of the native shrubland and grassland communities which are utilized by livestock at the lower elevations in this subregion.

The sections of the guide that describes the Central and Northern Rocky Mountain areas of the Subalpine and the Alpine subregions supplements the work done by Corns and Achuff (1982) in Banff and Jasper National Parks.

## **Climate**

### **Subalpine subregion**

The subalpine subregion is a Rocky Mountain altitudinal vegetation zone with its upper boundary formed by the Alpine subregion, whereas the lower boundary abuts the Montane, Foothills parkland and the Upper Foothills subregions. In Alberta, the subalpine extends from 1525 m to 2175 m at its southernmost occurrence and from 1360 m to 2000 m in the vicinity of Grande Cache (Strong 1992). The subalpine has a cordilleran climate characterized by snowy, cold winters and showery cool summers. Annual precipitation ranges from 329 mm to 916 mm, with maximum precipitation falling during July. The subalpine receives more precipitation during the winter months than any subregion (Strong 1992). The mean summer temperatures averages 9.4 °C and winter temperatures typically average -8.9 °C with December and January being the coldest months. The cold winter temperatures help to maintain the snowpack which

makes this an important watershed area.

The majority of the vegetation is dominated by seral lodgepole pine forests at lower elevations with Engelmann spruce and subalpine fir forests being more common at higher elevations. At timberline dwarf spruce, subalpine and whitebark pine are typical of the transition to the Alpine subregion. Imperfectly drained bottomlands are dominated by willow, bog birch, sedge, tufted hairgrass and californian oatgrass species and the steep south facing slopes are often dominated by fescue, hairy wildrye, wheatgrass and junegrass species.

### **Alpine subregion**

The Alpine subregion occurs above timberline in the Rocky Mountains of Alberta. Elevationally, alpine occupies areas greater than 2150 m in southern Alberta and declines to 2000 m in more northern portions. It is felt that the total annual precipitation is at least equal to the Subalpine subregion, which potentially makes the Alpine the wettest subregion in Alberta (Strong 1992). Summer temperatures are the coldest in Alberta with July mean temperatures averaging 10 °C. Freezing temperatures occur in all months of the year. Winter temperatures are probably colder than the subalpine subregion with temperatures probably never going above freezing for the whole winter. The cold temperatures help to maintain the snowpack for much of the year. Wind is also extremely important climatic factor in the Alpine. Although, precipitation is abundant the strong winds likely result in very large moisture deficits (Strong 1992).

The Alpine is characterized by low growing vegetation, which helps to protect the vegetation from the desiccating winds, and allows the plants to gain heat from the ground. Glaciers occur at the higher elevations where snow accumulation exceeds melt.



**Map 1. Location of Subalpine and Alpine subregions in Alberta**

## Methods

A community type approach (Mueggler 1988) to classification was chosen in preference to the habitat type approach (Daubenmire 1952) or ecosystem association approach (Corns and Annas 1986) because of the lack of understanding of the successional sequences of the communities. Community types are aggregates of similar plant communities based upon existing floristics regardless of successional status (Mueggler 1988). Community types are what is actually seen in the field. After defining the community types, they then can be linked to the ecosystem associations developed by Corns and Annas (1986) and Beckingham et al. (1996). In the mean time community types can be used as the basis for mapping and range management planning.

Individual plots were initially classified within a forest region using cluster analysis (SAS) and ordination (DECORANA, Gauch 1982). These types were described in individual carrying capacity guides for each forest. This led to differences in classification of the same types between forests, particularly for deciduous forest types. In an effort to standardize the community name and gain some understanding of each community types ecology, all plots sampled in each forest were reclassified. As the study progressed it became quite evident that there were differences in the productivity of the communities between subregions. As a result, it was decided to develop the classification within the subregion framework. A subregion is a geographical area that has broad vegetation zones combined with climatic data. As a result, the vegetation within each subregion is strongly influenced by the climatic conditions.

Sampling for this guide occurred within the Subalpine and Alpine subregions . This guide outlines the preliminary classification of 651 plots described in the Subalpine and 134 plots described in the Alpine subregion.

The procedure for inventory in the Southern Rocky Mtn. and Central and Northern Foothills areas followed the Range Survey Manual (1992) and uses the MF5 form. A plot consisted of a 10x10 m macroplot and ten randomly selected 1x1 m microplots to record the canopy cover of shrubs and ten nested 20x50 cm microplots to record the canopy cover of forbs and grass. At each macroplot a 50x100 cm was clipped and separated into trees, shrubs, forbs and graminoids, oven dried and weighed. The recommended stocking rate is based on 25 percent of the total production for forested types and 50% total production for grass and shrubland types and the fact that one animal requires 455 kg of dry weight material for one month of grazing.

For a description of the methodology for the remaining plots done in the Central and Northern Rocky Mountains of the Subalpine and Alpine subregions see Holland and Coen (1982).

### How to use the guide

In the Subalpine subregion it was found there was distinct differences between the grassland and shrubland community types between the Central and Northern Foothills (the foothills west of Sundre, Rocky Mtn. House, Hinton and Grande Cache), the Central and Northern Rocky



Mountains (mountains of Banff and Jasper National Park) and the Southern Rocky Mountains (foothills west of Calgary, Turner valley and areas south of Blairmore) (Strong and Thompson 1995). As a result the Subalpine was prestratified into 3 subdivisions. For the Subalpine first decide which area you are in then turn to the appropriate subdivision in the guide. The Alpine is generally all the vegetation types above timberline.

In order to understand how the community types in this guide are related to the ecosites and ecosite phases outlined in "Ecosites of West-Central Alberta and Southwestern Alberta" (Beckingham et al. 1996, Archibald et al. 1996), the community types in this guide are arranged by ecosite and ecosite phase for West-Central Alberta (Table 1) and Southwestern Alberta (Table 4). **Ecosites** are defined as ecological units that develop under similar environmental influences (climate, moisture and nutrient regime). An **ecosite phase** is a subdivision of the ecosite based on the dominant species in the canopy. Table 1 is a reproduction of Figure 20 in the Ecosites of West-Central Alberta guide and table 4 is a reproduction of Figure 20 in the Ecosites of Southwestern Alberta guide with the community types in this guide highlighted. For the most part the ecosites and ecosite phases are the same, particularly for the forested community types, but a number of new ecosites and ecosite phases had to be created for the grass and shrubland community types (Table 1). The ecosites included (bb)(subxeric/poor) yellow mountain avens, and (dd)(subhygric/medium) bog sedge meadow. The ecosite phases include (b2) bearberry grassland, (bb1) yellow mtn. avens, (c4) hairy wildrye grassland, (d4) California oatgrass, (d5) willow, (d6) grouseberry, (dd1) sedge-bog sedge, (dd2) shrubland, (e3) grass meadow, (g2) willow and (h3) grass bog for West-Central Alberta (Table 1). In Southwestern Alberta the new ecosites include (cc) (submesic/rich) rough fescue, and (i) fen. The ecosite phases included (a2) grassland, (b2) grassland, (c2) yellow mtn. avens, (cc1) rough fescue, (cc2) shrubland, (cc3) forb meadow, (f3) thimbleberry Aw, (f4) shrubby seepages, (i1)shrub fen and (i2) graminoid fen (Table 4). The "Grazing succession" category outlines the successional sequence the community type will undergo with increased grazing pressure. For a detailed description of the forested community types in the Subalpine please refer to the work done by Beckingham et al. (1996), Beckingham and Archibald (1996) and Archibald and Klappstein (1996). The dominant plant species, canopy cover, environmental conditions, response to grazing, forage production and carrying capacity of the grasslands and shrublands for the Subalpine and Alpine subregions are outlined in this guide.

Table 1. Ecosite, ecosite phases and community types for the Subalpine subregion of West-Central Alberta (adapted from Beekingham et al. 1996)(Highlighted communities are described in this guide, non-highlighted communities are outlined in guide to Ecosites of West-Central Alberta)

<b>Ecosite</b>	<b>Ecosite Phase</b>	<b>Forested Plant Community Type</b>	<b>Range plant community type</b>	<b>Grazing Succession</b>
a grassland (subxeric/medium)	a1 a1 shrubby grassland	a1.1 bearberry grassland	SACFA11 Blunt sedge-Junegrass/Bearberry	
		a1.2 willow-dwarf birch grassland	SACMB5 Bog birch/Juniper	
b bearberry/lichen (subxeric/poor)	a2 graminoid grassland	a2.1 Bellard's kobresia-hairy wild rye grassland	SACFA14. White Mtn. Avens/Bog sedge SACMA7 Northern wheatgrass SACFA10 Sedge-Hairy wildrye SACFA12 Fringed sage/Sedge-Junegrass	
	b1 bearberry/lichen P1	b1.1 P1/bearberry/lichen		
		b1.2 P1/bog cranberry/lichen		
bb yellow mtn. avens (submesic/poor)	b2 bearberry grassland	b1.3 P1/crowberry/lichen		
			SACMA4 Bearberry -Juniper	
c hairy wild rye (submesic/medium)	bb1 yellow mtn. avens		SACMA9 Yellow Mountain avens	
	c1 hairy wild rye P1	c1.1 P1/Canada buffalo-berry/hairy wild rye		
		c1.2 P1/juniper-bearberry/hairy wild rye		

d rhododendron-mesic (mesic/medium)	c2 hairy wild rye P1-Aw	c1.3	P1/green alder/hairy wild rye		
		c1.4	P1/hairy wild rye/feather moss		
		c2.1	P1-Aw/hairy wild rye		
		c3.1	Se/Canada buffalo-berry/hairy wild rye		
	c3 hairy wild rye Se	c3.2	Se/juniper-bearberry/hairy wild rye		
		c3.3	Se/willow/hairy wild rye		
		c3.4	Se/hairy wild rye/feather moss		
		c4	hairy wildrye grassland	SACMA3 Shrubby cinquefoil/Hairy wildrye SACMA5 Junegrass-Hairy wildrye-Brome SACMA6 Hairy wildrye/Bearberry-Juniper SACFA9 Rough fescue-H. wildrye-Sedge	
	d1 rhododendron-mesic P1	d1.1	P1/rhododendron/feather moss		
		d1.2	P1/false azalea/feather moss		
d1.3		P1/tall bilberry/feather moss			
d1.4		P1/Labrador tea/feather moss			
d1.5		P1/green alder/feather moss			
d1.6		P1/feather moss			
d2 rhododendron-mesic Se	d2.1	Se/rhododendron/feather moss			
	d2.2	Se/false azalea/feather moss			

		d2.3	Se/tall bilberry/feather moss		
		d2.4	Se/Labrador tea/feather moss		
		d2.5	Se/green alder/feather moss		
		d2.6	Se/subalpine fir/feather moss		
		d2.7	Se/feather moss		
	d3	d3.1	Fa/rhododendron/feather moss		
		d3.2	Fa/false azalea/feather moss		
		d3.3	Fa/tall bilberry/feather moss		
		d3.4	Fa/subalpine fir/feather moss		
		d3.5	Fa/feather moss	SACMB8	Subalpine fir
	d4		California oatgrass	SACFA8 California oatgrass-Sedge SACMA1 Bog sedge-California oatgrass SACMA8 Alpine bluegrass	
	d5		Willow	SACFB6 Willow-Bog birch/California oatgrass	
	d6		grouseberry	SACMB7 Grouseberry-Juniper	
dd bog sedge meadow (subhygric/medium)		dd1	sedge-bog sedge	SACFA13 Sedge-Bog sedge-Tufted hairgrass	


	dd2 shrubland		<p>SACFB8 Willow-Bog birch/Bog sedge-Sedge  SACFB9 Bog birch-Willow/Rough fescue  SACFB10 Bog birch/Rough fescue-Bog sedge  SACMB4 Willow-Bog birch/Bog sedge</p>	
e meadow (subhygric/very rich)	e1 shrubby meadow	e1.1 willow-dwarf birch meadow	<p>SACFB3 Willow/Graceful sedge  SACFB4 Willow-Bog birch/Tufted hairgrass  SACFB7 Willow-Bog birch/Hairy wildrye  SACFB11 Willow/Fringed brome-Sedge  SACMB2 Willow/Sedge  SACMB3 Willow-Bog birch/Hairy wildrye  SACMB6 Willow/Forb</p>	SACFB5 Willow-Bog birch/Clover-Dandelion
	e2 forb meadow	e2.1 meadow rue meadow	SACMA2 Forb meadows	
	e3 grass meadow		<p>SACFA4 Tufted hairgrass-Sedge  SACFA17 Fireweed-Meadow rue/Sedge-H. wildrye</p>	<p>SACFA5 Sedge-Tufted hairgrass  SACFA6 Sedge-Rocky Mtn. fescue-Alpine timothy  SACFA7 Sedge-Slender wheatgrass-Fringed brome/Forb  SACFA16 Kentucky bluegrass-Sedge/Dandelion  SACFA15 Creeping red fescue-Sedge</p>

f rhododendron- subhygric (subhygric/mediu m)	f1 rhododendron-subhygric P1	f1.1	P1/rhododendron/feather moss		
		f1.2	P1 false azalea/feather moss		
		f1.3	P1/Labrador tea/feather moss		
g horsetail (hygric/rich)	f2 rhododendron-subhygric Se-Fa	f2.1	Se-Fa/rhododendron/feather moss		
		f2.2	Se-Fa/false azalea/feather moss		
		f2.3	Se-Fa/Labrador tea/feather moss		
h bog (subhydric/poor)	g1 horsetail Se	g1.1	Se/willow/horsetail		
		g1.2	Se/feather moss		
i fen (subhydric/rich)	g2 willow	h1.1	Sb/cloudberry/feather moss-peat moss	SACFB2 Willow/Horsetail	
		h2.1	Labrador tea/cloudberry/peat moss		
		h3	grass bog	SACFA2. Tufted bulrush SACFA3. Sedge-Cottongrass	
i fen (subhydric/rich)	i1 treed fen	i1.1	Sb-Se/willow/sedge/peat moss		
		i1.2	Sb-Se/willow-dwarf birch/sedge/golden moss		
	i2 shrubby fen	i2.1	willow/sedge/tufted moss-peat moss	SACFB1 Willow-Bog birch/Water sedge SACMB1 Willow/Water sedge	
		i2.2	willow-dwarf birch/sedge/peat moss-golden moss		

	i3 graminoid fen	i2.3 dwarf birch/sedge/peat moss-golden moss  i3.1 sedge fen	SACFA1. Water sedge-Beaked sedge	
--	---------------------	--	----------------------------------	--

## b2 bearberry grassland (n=9)

---



## SITE CHARACTERISTICS

---

**Moisture regime:** subxeric, mesic  
**Nutrient regime:** poor, medium  
**Topographic position:** upper slope  
**Slope:** (16-30)(46-70)  
**Aspect:** southerly

## SOIL CHARACTERISTICS

---

**Organic thickness:** (0-5)  
**Humus form:** mor  
**Surface texture:** SL,S,L  
**Effective texture:** SL,LS  
**Depth to Mottles/Gley:** none  
**Drainage:** rapid, well  
**Parent material:** M,GF  
**Soil subgroup:** E.DYB, O.HFP, BR.GL, O.EB

## COMMUNITY TYPES

---

SACMA4. Bearberry-Juniper (n=9)

## CHARACTERISTIC SPECIES

---

### Trees

[ 2 ] Aspen

### Shrubs

[ 4 ] Willow

[ 2 ] White mtn. avens

[ 4 ] Shrubby cinquefoil

[ 11 ] Ground juniper

### Forbs

[ 16 ] Bearberry

[ 2 ] White camas

[ 1 ] Strawberry

[ 1 ] Yarrow

### Graminoids

[ 2 ] Sedge

[ 1 ] Spiked trisetum


[ 3 ] Hairy wildrye



**bb yellow mtn. avens (n=4)**

**GENERAL DESCRIPTION**

This ecosite is located on recent fluvial and glacialfluvial landforms with gentle slopes. The soils are poorly developed gravels and are rapidly drained. Yellow mountain avens, bearberry, juniper and junegrass are typical of these early successional river flats. The poor soil conditions limits the forage productivity and amount of regrowth after grazing. This ecosite should be rated as non-use range.




**SUCCESSIONAL RELATIONSHIPS**

Yellow mtn. avens generally dominates this community in the early successional stages. Succession in the absence of disturbance will be to balsam poplar, Engelmann spruce and subalpine fir.

**INDICATOR SPECIES**

- yellow mtn. avens
- willow
- silverberry
- juniper
- showy locoweed
- bearberry
- alpine fireweed
- sedge

**subxeric/medium**



**SITE CHARACTERISTICS**

- Moisture regime:** subxeric, submesic
- Nutrient regime:** poor, medium
- Topographic position:** floodplain
- Slope:** (0-5)
- Aspect:** variable


**SOIL CHARACTERISTICS**

- Organic thickness:** (0-2)
- Humus form:** mor
- Surface texture:** SL, SiL
- Effective texture:** SL,
- Depth to Mottles/Gley:** none
- Drainage:** rapid, well
- Parent material:** F, GF
- Soil subgroup:** O, R

**ECOSITE PHASES**

- bb1 yellow mtn. avens (n=4)

**bb1 yellow mtn. avens (n=4)**



**SITE CHARACTERISTICS**

**Moisture regime:** subxeric, submesic  
**Nutrient regime:** poor, medium  
**Topographic position:** floodplain  
**Slope:** (0-5)  
**Aspect:** variable

**SOIL CHARACTERISTICS**

**Organic thickness:** (0-2)  
**Humus form:** mor  
**Surface texture:** SL,SiL  
**Effective texture:** SL,  
**Depth to Mottles/Gley:** none  
**Drainage:** rapid, well  
**Parent material:** F,GF  
**Soil subgroup:** O.R

**COMMUNITY TYPES**

SACMA9. Yellow mountain avens (n=4)

**CHARACTERISTIC SPECIES**

**Trees**

- [ 2 ] Engelmann spruce
- [ 1 ] Subalpine fir
- [ 1 ] White spruce

**Shrubs**

- [ 5 ] Willow
- [ 1 ] Buffaloberry
- [ 50 ] Yellow mtn. avens
- [ 1 ] Shrubby cinquefoil


**Forbs**

- [ 1 ] Alpine fireweed
- [ 1 ] Showy locoweed
- [ 1 ] Alpine goldenrod

**Graminoids**

- [ 1 ] Sedge
- [ 1 ] Spiked trisetum

## c4 hairy wildrye grassland (n=81)



### CHARACTERISTIC SPECIES

#### Shrubs

- [ 2 ] Juniper
- [ 10 ] Shrubby cinquefoil

#### Forbs

- [ 9 ] Bearberry
- [ 2 ] Showy locoweed
- [ 2 ] Strawberry
- [ 1 ] Old man's whiskers
- [ 1 ] White mtn. avens
- [ 1 ] Yellow hedysarum

#### Graminoids

- [ 5 ] Rough fescue
- [ 15 ] Hairy wildrye
- [ 5 ] Sedge
- [ 5 ] Junegrass

### SITE CHARACTERISTICS

**Moisture regime:** subxeric, submesic,  
**Nutrient regime:** medium  
**Topographic position:** upper slope  
**Slope:** (16-30)(47-70)  
**Aspect:** southerly


### SOIL CHARACTERISTICS

**Organic thickness:** (0-2)(2-5)  
**Humus form:** no data  
**Surface texture:** SiL  
**Effective texture:** SL,  
**Depth to Mottles/Gley:** none  
**Drainage:** rapid, well  
**Parent material:** M  
**Soil subgroup:** O.R, O.EB

### COMMUNITY TYPES

SACMA3 Shrubby cinquefoil/Hairy wildrye (n=5)  
 SACMA5 Junegrass-Hairy wildrye-Brome (n=19)  
 SACMA6 Hairy wildrye/Bearberry-Juniper (n=44)  
 SACFA9 Rough fescue-Hairy wildrye-Sedge (n=13)

**d4 california oatgrass grassland (n=7)**



**SITE CHARACTERISTICS**

**Moisture regime:** mesic, subhygric, submesic,  
**Nutrient regime:** medium, rich, poor  
**Topographic position:** level, midslope  
**Slope:** (0-5)  
**Aspect:** variable

**SOIL CHARACTERISTICS**

**Organic thickness:** (0-2)(2-5)  
**Humus form:** no data  
**Surface texture:** SiL, L  
**Effective texture:** SL, SiL, L  
**Depth to Mottles/Gley:** none  
**Drainage:** mod. well, well  
**Parent material:** M  
**Soil subgroup:** O.R, O.EB, BR.GL

**COMMUNITY TYPES**

SACFA8 California oatgrass-Sedge (n=5)  
 SACMA1 Bog sedge-California oatgrass (n=1)  
 SACMA8 Alpine bluegrass (n=1)

**CHARACTERISTIC SPECIES**

**Shrubs**

- [ 11 ] Willow
- [ 13 ] Shrubby cinquefoil

**Forbs**


- [ 5 ] Bearberry
- [ 3 ] Veiny meadow rue
- [ 8 ] Strawberry
- [ 2 ] Yarrow
- [ 2 ] Slender blue beardtongue
- [ 1 ] Fireweed

**Graminoids**

- [ 30 ] California oatgrass
- [ 3 ] Hairy wildrye
- [ 8 ] Sedge
- [ 9 ] Bog sedge
- [ 15 ] Alpine bluegrass

d5

willow (n=19)



### CHARACTERISTIC SPECIES

#### Shrubs

- [ 30 ] Willow
- [ 7 ] Bog birch

#### Forbs

- [ 1 ] Graceful cinquefoil
- [ 1 ] Veiny meadow rue
- [ 9 ] Strawberry
- [ 2 ] Yarrow
- [ 2 ] Globeflower
- [ 1 ] Wandering daisy

#### Graminoids

- [ 24 ] California oatgrass
- [ 4 ] Mountain timothy
- [ 10 ] Sedge
- [ 1 ] Slender wheatgrass
- [ 1 ] Spiked trisetum

### SITE CHARACTERISTICS

**Moisture regime:** mesic, subhygric, submesic,  
**Nutrient regime:** medium, rich, poor  
**Topographic position:** level, midslope  
**Slope:** (0-5)  
**Aspect:** variable

### SOIL CHARACTERISTICS


**Organic thickness:** (0-2)(2-5)  
**Humus form:** no data  
**Surface texture:** SiL, L  
**Effective texture:** SL, SiL, L  
**Depth to Mottles/Gley:** none  
**Drainage:** mod. well, well  
**Parent material:** M  
**Soil subgroup:** O.R, O.EB, BR.GL

### COMMUNITY TYPES

SACFB6 Willow-Bog birch/California oatgrass(n=19)

## d5 grouseberry (n=4)

---



## SITE CHARACTERISTICS

---

**Moisture regime:** mesic, submesic,  
**Nutrient regime:** medium  
**Topographic position:** level, midslope  
**Slope:** (0-5)  
**Aspect:** variable

## SOIL CHARACTERISTICS

---

**Organic thickness:** (0-2)(2-5)  
**Humus form:** no data  
**Surface texture:** SiL, L  
**Effective texture:** SL, SiL, L  
**Depth to Mottles/Gley:** none  
**Drainage:** mod. well, well  
**Parent material:** M  
**Soil subgroup:** O.R, O.EB, BR.GL

## COMMUNITY TYPES

---

SACMB7 Grouseberry-Juniper (n=4)

## CHARACTERISTIC SPECIES

---

### Trees

[ 1 ] Subalpine fir

### Shrubs

[ 1 ] Willow  
[ 14 ] Grouseberry  
[ 5 ] Ground juniper  
[ 1 ] Crowberry

### Forbs

[ 12 ] Fireweed  
[ 3 ] Small lv'd everlasting  
[ 3 ] Strawberry  
[ 2 ] Yarrow

### Graminoids


[ 6 ] California oatgrass  
[ 1 ] Sedge  
[ 7 ] Spiked trisetum

**dd bog sedge meadow (n=30)**

veiny meadow rue  
sedge  
**subhygric/medium**

**GENERAL DESCRIPTION**

This ecosite is located on moist well drained lowland sites adjacent to rivers and streams at higher elevations in the Central and Northern Foothills of the Subalpine. The presence of bog sedge appears to indicate the transition to the higher Alpine subregion. Indeed, Oglivie (1969) described bog sedge dominated community types on windswept ridges at higher elevations in the Alpine subregion. The forage production of this community is only moderate. Perhaps, the higher elevation and colder climate which favours the growth of bog sedge limits total




productivity of this site.

**SUCCESSIONAL RELATIONSHIPS**

Due to the nature of the site grasslands often remain the climax vegetation on these sites. In the absence of disturbance willow and bog birch often invade to form willow and bog birch dominated shrublands.

**INDICATOR SPECIES**

- california oatgrass
- bog sedge
- rough fescue
- tufted hairgrass
- sedge
- willow
- bog birch



**SITE CHARACTERISTICS**

**Moisture regime:** mesic, subhygric  
**Nutrient regime:** medium, rich  
**Topographic position:** floodplain  
**Slope:** (0-5)  
**Aspect:** variable

**SOIL CHARACTERISTICS**


**Organic thickness:** (0-15)  
**Humus form:** no data  
**Surface texture:** L, SiL  
**Effective texture:** SiL,  
**Depth to Mottles/Gley:** none  
**Drainage:** well, mod. well  
**Parent material:** F, GF  
**Soil subgroup:**, O.R, O.HR

**ECOSITE PHASES**

- dd1 sedge-bog sedge(n=6)
- dd2 shrubland(n=24)

## dd1 sedge-bog sedge (n=6)

---



## SITE CHARACTERISTICS

---

**Moisture regime:** mesic, subhygric  
**Nutrient regime:** medium  
**Topographic position:** level, midslope  
**Slope:** (0-5)  
**Aspect:** variable

## SOIL CHARACTERISTICS

---

**Organic thickness:** (0-15)  
**Humus form:** no data  
**Surface texture:** SiL, L  
**Effective texture:** SL, SiL, L  
**Depth to Mottles/Gley:** none  
**Drainage:** mod. well, well  
**Parent material:** F  
**Soil subgroup:** O.R, O.HR

## COMMUNITY TYPES

---

SACFA13 Sedge-Bog sedge-Tufted hairgrass (n=6)

## CHARACTERISTIC SPECIES

---

### Shrubs

[ 1 ] Shrubby cinquefoil

### Forbs


[ 7 ] Alpine goldenrod  
[ 3 ] Graceful cinquefoil  
[ 1 ] Strawberry  
[ 5 ] Yarrow  
[ 6 ] Alpine bistort

### Graminoids

[ 20 ] Bog sedge  
[ 40 ] Sedge  
[ 13 ] Tufted hairgrass  
[ 7 ] Hairy wildrye  
[ 3 ] Rocky mtn. fescue



## dd2 shrubland (n=24)



## SITE CHARACTERISTICS

**Moisture regime:** mesic, subhygric  
**Nutrient regime:** medium  
**Topographic position:** level, midslope  
**Slope:** (0-5)  
**Aspect:** variable

## SOIL CHARACTERISTICS

**Organic thickness:** (0-15)  
**Humus form:** no data  
**Surface texture:** SiL, L  
**Effective texture:** SL, SiL, L  
**Depth to Mottles/Gley:** none  
**Drainage:** mod. well, well  
**Parent material:** F  
**Soil subgroup:** O.R, O.HR

## CHARACTERISTIC SPECIES

### Shrubs

- [ 7 ] Willow
- [ 30 ] Bog birch

### Forbs

- [ 1 ] Alpine goldenrod
- [ 2 ] Graceful cinquefoil
- [ 3 ] Old man's whiskers
- [ 1 ] Yarrow
- [ 1 ] Alpine bistort
- [ 2 ] False dandelion
- [ 2 ] Tall larkspur
- [ 1 ] Larkspur


### Graminoids

- [ 17 ] Bog sedge
- [ 10 ] Sedge
- [ 7 ] Rough fescue
- [ 1 ] Hairy wildrye
- [ 1 ] California oatgrass

## COMMUNITY TYPES

- SACFB8 Bog birch/Bog sedge-Sedge (n=5)
- SACFB9 Bog birch-Willow/Rough fescue(n=4)
- SACFB10 Bog birch/Rough fescue-Bog sedge(n=1)
- SACMB4 Willow-Bog birch/Bog sedge (n=14)

**e3 grass meadow (n=19)**



**CHARACTERISTIC SPECIES**

**Shrubs**

- [ 2 ] Willow
- [ 1 ] Bog birch
- [ 1 ] Shrubby cinquefoil

**Forbs**

- [ 3 ] Veiny meadow rue
- [ 2 ] Graceful cinquefoil
- [ 1 ] Old man's whiskers
- [ 2 ] Yarrow
- [ 1 ] Alpine bistort
- [ 2 ] False dandelion
- [ 2 ] Tall larkspur

**Graminoids**

- [ 10 ] Tufted hairgrass
- [ 10 ] Sedge
- [ 2 ] Slender wheatgrass
- [ 1 ] Hairy wildrye
- [ 1 ] California oatgrass

**SITE CHARACTERISTICS**

**Moisture regime:** mesic, subhygric  
**Nutrient regime:** rich, medium  
**Topographic position:** level, midslope  
**Slope:** (0-5)  
**Aspect:** variable


**SOIL CHARACTERISTICS**

**Organic thickness:** (0-15)  
**Humus form:** mor, moder  
**Surface texture:** SiL, L  
**Effective texture:** SiL, L, CL, C, SiCL, SCL, LS  
**Depth to Mottles/Gley:** (0-25)  
**Drainage:** mod. well, well, poor  
**Parent material:** F  
**Soil subgroup:** R.G, O.HG, O.HR, O.MB, CU.R

**COMMUNITY TYPES**

SACFA4 Tufted hairgrass-Sedge (n=18)  
 SACFA17 Fireweed-Meadow rue/Sedge-Hairy wildrye (n=1)

**e3 grazed grass meadow (n=33)**



**CHARACTERISTIC SPECIES**

**Shrubs**

- [ 2 ] Willow
- [ 1 ] Bog birch
- [ 1 ] Shrubby cinquefoil

**Forbs**

- [ 3 ] Veiny meadow rue
- [ 2 ] Graceful cinquefoil
- [ 1 ] Old man's whiskers
- [ 2 ] Yarrow
- [ 1 ] Alpine bistort
- [ 2 ] False dandelion
- [ 2 ] Tall larkspur
- [ 3 ] Dandelion

**Graminoids**

- [ 7 ] Tufted hairgrass
- [ 10 ] Sedge
- [ 2 ] Slender wheatgrass
- [ 1 ] Hairy wildrye
- [ 1 ] California oatgrass
- [ 10 ] Creeping red fescue
- [ 12 ] Kentucky bluegrass
- [ 2 ] Rocky mtn. fescue

**SITE CHARACTERISTICS**

**Moisture regime:** mesic, subhygric  
**Nutrient regime:** rich, medium  
**Topographic position:** level, midslope  
**Slope:** (0-5)  
**Aspect:** variable


**SOIL CHARACTERISTICS**

**Organic thickness:** (0-15)  
**Humus form:** mor, moder  
**Surface texture:** SiL, L  
**Effective texture:** SiL, L, CL, C, SiCL, SCL, LS  
**Depth to Mottles/Gley:** (0-25)  
**Drainage:** mod. well, well, poor  
**Parent material:** F  
**Soil subgroup:** R.G, O.HG, O.HR, O.MB, CU.R

**COMMUNITY TYPES**

- SACFA5 Sedge-Tufted hairgrass(n=7)
- SACFA6 Sedge-Rocky Mtn. fescue-Alpine timothy(n=6)
- SACFA7 Sedge-Slender wheatgrass-Fringed brome/Forb(n=18)
- SACFA15 Creeping red fescue-Sedge(n=1)
- SACFA16 Kentucky bluegrass-Sedge/Dandelion(n=1)

**g2 willow (n=4)**



**SITE CHARACTERISTICS**

Moisture regime: hygric, subhygric  
 Nutrient regime: rich, medium  
 Topographic position: level, midslope, toe  
 Slope: (0-5)  
 Aspect: variable

**SOIL CHARACTERISTICS**

Organic thickness: (6-15)  
 Humus form: mor  
 Surface texture: humic, SiL, L, SiCL  
 Effective texture: humic, SiL, L, SCL, SC  
 Depth to Mottles/Gley: (0-25)(26-50)  
 Drainage: mod. well, poor, imperfect  
 Parent material: F, M, FL  
 Soil subgroup: R, HG, O, R, T, H

**COMMUNITY TYPES**

SACFB2 Willow/Horsetail (n=4)

**CHARACTERISTIC SPECIES**

**Shrubs**

- [ 30 ] Willow
- [ 6 ] Bog birch


**Forbs**

- [ 12 ] Variegated horsetail
- [ 9 ] Common horsetail
- [ 3 ] Showy everlasting

**Graminoids**

- [ 6 ] Tufted hairgrass
- [ 16 ] Sedge

### h3 grass bog (n=19)



#### SITE CHARACTERISTICS

**Moisture regime:** subhydryc, hygic

**Nutrient regime:** medium, poor

**Topographic position:** depression

**Slope:** (0-5)

**Aspect:** variable

#### SOIL CHARACTERISTICS

**Organic thickness:** (>80)

**Humus form:** peatymor

**Surface texture:** fibric

**Effective texture:** mesic

**Depth to Mottles/Gley:** (0-25)(26-50)

**Drainage:** poor, imperfect

**Parent material:** O

**Soil subgroup:** T.M, FI.OC

#### COMMUNITY TYPES

SACFA2 Tufted bulrush (n=12)

SACFA3 Sedge-Cottongrass(n=7)

#### CHARACTERISTIC SPECIES

##### Shrubs

[ 2 ] Willow

[ 2 ] Bog birch

##### Forbs

[ 1 ] Elephant's head

[ 2 ] Common horsetail

[ 1 ] Woolly everlasting

[ 1 ] Lanced leaved paintbrush

##### Graminoids

[ 8 ] Tufted bulrush

[ 15 ] Sedge

[ 3 ] Cottongrass

[\(Back to table of contents\)](#)

## Results

The analysis of the 785 plots distinguished 94 community types. These types were split into 4 categories:

### Subalpine

#### Central and Northern Foothills

A. Native grasslands 17 community types

B. Native shrublands 11 community types

#### Central and Northern Rocky Mountains

A. Native grasslands 9 community types

B. Native shrublands 8 community types

#### Southern Rocky Mountains

A. Native grasslands 19 community types

B. Native shrublands 5 community types

C. Grazed grasslands 9 community types

D. Deciduous 6 community types

E. Conifer 4 community types

### Alpine

A. Native grasslands and shrublands 6 community types

The dominant plant species, canopy cover, environmental conditions, forage production and carrying capacity (when available) are outlined for each community type.

## SUBALPINE SUBREGION

### SUBALPINE SHRUB AND GRASSLAND ECOLOGY

The Subalpine subregion has highly variable ecological conditions. Much of the variation is the result of complex topography, with a strong ecotonal effect from the surrounding subregions. For instance the grasslands and shrublands of the foothills west of Sundre, Rocky Mtn. House, Hinton and Grande Cache are very similar to the tufted hairgrass, California oatgrass, sedge and rough fescue dominated grass and shrublands of the Upper Foothills subregion. This area is represented by the Central and Northern Foothills, an area dominated by morainal and residuum deposits, gentler slopes (16-45%) and Brunisolic and Luvisolic soils. The Central and Northern Foothills are transitional from the lower Upper Foothills subregion to the higher and steeper Central and Northern Rocky Mountains of the Subalpine subregion. The Central and Northern Rocky Mountains are typical of the morainal and talus deposits on steeper slopes (10-100%), with Brunisolic and Regosolic soils in Banff and Jasper National Parks. These areas are transitional from the lower Subalpine to the higher Alpine subregion. Many species such as bog sedge, heather spp., white mountain avens which are characteristic of alpine communities start to become predominant in these areas. Grasslands on steep, south-facing slopes are dominated by hairy wildrye, junegrass and shrubby cinquefoil.

In southern Alberta (west of Turner valley, south of Blairmore) the subalpine grasslands and shrublands are strongly influenced by the lower Montane subregion. Many of the grass species associated with the Montane (rough fescue, Parry oatgrass, Idaho fescue) dominate the south facing slopes of the Subalpine. These grasslands are very different from the hairy wildrye dominated community types found in the Central and Northern Rocky Mountains. The Southern Rocky Mountains are dominated by residuum, morainal and talus deposits on gentle to steep slopes (16-100%).

**SUBALPINE SUBREGION**

**CENTRAL AND NORTHERN FOOTHILLS**

**NATIVE GRASSLANDS AND SHRUBLANDS**



**Figure 1.** This figure is typical of the Tufted hairgrass-Sedge community with succession to a Willow-Bog birch dominated community type in the Central and Northern Foothills of the Subalpine subregion.



## Native grass and shrubland ecology of the Foothills

The native grass and shrubland community types in the Central and Northern Foothills of the Subalpine subregion (Table 2) are found in the valley bottoms adjacent to streams and rivers. The community types in this area are very similar to the grass and shrublands found in the Upper Foothills subregion and represent a transition from the lower Upper Foothills subregion to the Central and Northern Rocky Mountains of Banff and Jasper National Parks. The sequence of these community types along a moisture gradient from wet (Sedge meadows) to dry south facing slopes (Blunt sedge-Junegrass/Bearberry, Fringed sage/Junegrass-Sedge) is outlined in Figure 2. The change in species composition from the wet sedge meadows to tufted hairgrass, California oatgrass or rough fescue meadows may occur over a 3 foot elevational gradient. The presence of bog sedge (*Kobresia myosuroides*) in the White Mtn. avens-Bog sedge and Sedge-Bog sedge-Tufted hairgrass community types appears to indicate the transition from the lower Central and Northern Foothills to the higher Central and Northern Rocky Mountains. Ogilvie (1969) and Corns and Achuff (1982), described bog sedge dominated community types in the higher elevations of the subalpine and alpine of the Rocky Mountains of Banff and Jasper National Parks.

The maintenance of these grassland community types is extremely fire dependent. The lack of fire quickly allows bog birch and willow to expand shading the modal grassland community types. Prolonged shading causes the understory composition to shift from a tufted hairgrass-California oatgrass dominated understory to a slender wheatgrass-hairy wildrye dominated understory (Figure 2). Under a heavy shrub cover there is little forb or grass cover. Increased shrub cover also causes a decline in forage productivity and reduces the accessibility for livestock.

Many of these subalpine grass and shrublands are very fragile because of exposure and cold climate. The forage productivity is generally only half of what is found in the lower Upper Foothills subregion and recovery from overgrazing will likely take some time because of the poor growing conditions. As a result grazing by domestic livestock should be done with caution.

**Table 2.** Native grass and shrublands of the Central and Northern foothills ecodeistricts of the Subalpine subregion

Community number	Community type	Grass	Forb	Shrub	Total	Moisture	Drainage	Carrying capacity (Ha/AUM)	
<b>A. GRASSLANDS</b>									
SACFA1.	Water sedge-Beaked sedge	1215	774	0	1721	Subhydryc	Poorly	Non-use	
SACFA2.	Tufted bulrush	-	-	-	N/A	Subhydryc	Poorly	Non-use	
SACFA3.	Sedge-Cottongrass	-	-	-	N/A	Subhydryc	Poorly	Non-use	
SACFA4.	Tufted hairgrass-Sedge	997	288	0	1284	Subhydryc	Mod. Well	0.7	
SACFA5.	Sedge-Tufted hairgrass	785	146	0	931	Subhydryc	Mod. Well	1.0	
SACFA6.	Sedge-Rocky Mtn. fescue-Alpine Timothy	661	326	7	994	Mesic	Mod. Well	0.9	
SACFA7.	Sedge-Slender wheatgrass-Fringed Brome/Forb	539	156	7	702	Subhydryc	Mod. Well	1.3	
SACFA8.	California oatgrass-Sedge	921	352	0	1273	Submesic	Mod. Well	0.7	
SACFA9.	Rough fescue-Hairy wildrye-Sedge	1487	689	167	2343	Submesic	Well	0.4	
SACFA10.	Sedge-Hairy wildrye	771	390	284	1444	Subxeric	Rapidly	Non-use	
SACFA11.	Blunt sedge-Junegrass/Bearberry	1235	264	13	1512	Subxeric	Rapidly	Non-use	
SACFA12.	Fringed sage/Junegrass-Sedge	1133	545	250	2126	Subxeric	Rapidly	Non-use	
SACFA13.	Sedge-Bog sedge-Tufted hairgrass	582	158	0	740	Subhydryc	Well	Non-use	
SACFA14.	White Mtn. avens/Bog sedge	517	245	0	762	Subxeric	Rapidly	Non-use	
SACFA15.	Creeping red fescue-Sedge	1705	15	0	1720	Mesic	Well	0.6	
SACFA16.	Kentucky bluegrass-Sedge/Dandelion	380	224	0	604	Mesic	Well	1.5	
SACFA17.	Fireweed-Meadow rue/Sedge-Wildrye	1006	2134	0	3140	Subhydryc	Mod. well	0.3	
<b>B. SHRUBLANDS</b>									
SACFB1.	Willow-Bog birch/Water sedge	1234	129	1150	2514	Hygic	Poorly	Non-use	

Table 2. cont'd

Community number	Community type	Production(kg/ha)			Moisture	Drainage	Carrying capacity (ha/AUM)	
		Grass	Forb	Shrub				Total
SACFB2.	Willow/Horsetail	-	-	-	N/A	Subhygric	Poorly	Non-use
SACFB3.	Willow/Graceful sedge	806	109	3	919	Subhygric	Mod. well	1.0
SACFB4.	Willow-Bog birch/Tufted hairgrass	950	493	265	1803	Subhygric	Mod. well	0.5
SACFB5.	Willow-Bog birch/Clover-Dandelion	1728	199	0	1927	Subhygric	Mod. well	0.5
SACFB6.	Willow-Bog birch/California oatgrass	598	418	300	1316	Subhygric	Mod. well	0.7
SACFB7.	Willow-Bog birch/Hairy wildrye	868	713	135	1716	Subhygric	Well	0.5
SACFB8.	Bog birch/Bog sedge-Sedge	1333	390	202	1925	Subhygric	Mod. well	0.5
SACFB9.	Bog birch-Willow/Rough fescue	1807	705	559	3071	Mesic	Mod. well	0.4
SACFB10.	Bog birch/Rough fescue-Bog sedge	1201	147	0	1348	Mesic	Well	0.7
SACFB11.	Willow/Fringed brome-Sedge	472	288	0	760	Mesic	Imperfectly	1.1

\*Estimate

## Central and Northern Foothills Ecodistricts

## Central and Northern Rocky Mountain ecodistricts




Figure 2. Grassland and shrubland community types in the landscape of the Central and Northern Rocky Mountain and Foothills areas of the Subalpine subregion.

## Native grass and shrublands in Central and Northern Mountain and Foothills areas

1. Community above timberline (Alpine subregion).....	<b>Alpine section of guide</b>	
Community not above timberline, meadows, shrublands or south facing slopes dominated by grass.....		2
2. Moist sites, community dominated by shrubs >20% cover (willow, bog birch) or timberline grouseberry .....	<b>see shrub key pg 34</b>	
Drier to moist sites shrub cover <10% site dominated by grasses and forbs.....		3
3. Community very wet free standing water, dominated by sedge, cottongrass or tufted bulrush .....		4
Community drier, dominated by tufted hairgrass, rough fescue, california oatgrass, hairy wildrye, juniper or bearberry.....		6
4. Very wet nutrient rich sites dominated by water or beaked sedge.....	<b>SACFA1</b>	
Very wet nutrient poor, acidic sites dominated by tufted bulrush and cottongrass.....		5
5. Site dominated by tufted bulrush.....	<b>SACFA2</b>	
Site dominated by cottongrass and sedge.....	<b>SACFA3</b>	
6. Grasslands of meadows and lowland areas.....		7
Grasslands of south facing slopes, or windswept ridges, hairy wildrye, bearberry, white mtn. avens dominated .....		20
7. Higher elevation sites near or at timberline, moist sites dominated by wandering daisy, globeflower, mountain marigold.....	<b>SACMA2</b>	
Lower elevation sites, valley bottoms adjacent to streams or rivers or if higher elevation dry sites dominated by bog sedge.....		8
8. Disturbed or grazed community types dominated by Kentucky bluegrass, creeping red fescue, dandelion, alpine bluegrass, alpine timothy or fireweed.....		9
Undisturbed community types dominated by rough fescue, California oatgrass, bog sedge, tufted hairgrass, and sedge species.....		14
9. Moderately grazed site native increasers dominant (slender wheatgrass, sedge, fringed brome, rocky mtn. fescue, alpine timothy), Alpine or Kentucky bluegrass increasing in cover.....		10
Heavily grazed sites dominated by Kentucky bluegrass or abandoned airstrips dominated by creeping red fescue or native forb species (fireweed, veiny meadow rue).....		12
10. Site dominated by Alpine bluegrass.....	<b>SACMA8</b>	
Site dominated by rocky mtn. fescue, alpine timothy, sedge, slender wheatgrass, fringed brome (moister sites).....		11
11. Site dominated by rocky mtn. fescue, alpine timothy, sedge.....	<b>SACFA6</b>	
Site dominated by slender wheatgrass, sedge, fringed brome.....	<b>SACFA7</b>	
12. Heavily grazed site dominated by Kentucky bluegrass and dandelion.....	<b>SACFA16</b>	
Abandoned airstrips dominated by creeping red fescue, or invaded strips dominated by forbs (fireweed, veiny meadow).....		13
13 Site dominated by creeping red fescue.....	<b>SACFA15</b>	
Moister disturbed site dominated by fireweed and veiny meadow rue... <b>SACFA17</b>		
14. Moist sites dominated by sedge and tufted hairgrass.....		15
Drier sites dominated by rough fescue, hairy wildrye, bog sedge, yellow dryas or California oatgrass.....		16
15. Site dominated by tufted hairgrass, sedge co-dominant.....	<b>SACFA4</b>	
Site dominated by sedge, tufted hairgrass co-dominant.....	<b>SACFA5</b>	
16. Sites dominated by rough fescue and hairy wildrye.....	<b>SACFA9</b>	
Sites dominated by California oatgrass, bog sedge, or yellow dryas.....		17
17. Gravelly river flats dominated by yellow dryas.....	<b>SACMA9</b>	
Meadow areas dominated by California oatgrass and/or bog sedge.....		18
18. Site dominated by bog sedge and california oatgrass.....		19
Site dominated by California oatgrass and sedge, bog sedge not present.....	<b>SACFA8</b>	
19. Site dominated by bog sedge, california oatgrass, drier sites.....	<b>SACMA1</b>	
Site co-dominated by bog sedge, tufted hairgrass, and sedge moister sites..	<b>SACFA13</b>	
20. Lower elevation grasslands in the Foothills of the Subalpine.....		21
Higher elevation grasslands in the mountains of the Subalpine.....		22
21 Northern wheatgrass dominated site.....	<b>SACMA7</b>	

Fringed sage, sedge and junegrass dominated slope.....	SACFA12	
22. Avalanche slopes dominated by hairy wildrye, juniper, and bearberry....	SACMA6	
Drier sites or windswept ridges dominated hairy wildrye, juniper, bearberry, shrubby cinquefoil, white mtn. avens.....		23
23. Windswept ridges dominated by white mtn. avens.....	SACFA14	
South facing slopes dominated by hairy wildrye.....		24
24. Shallow rocky soils with little grass cover, site dominated by bearberry..	SACMA4	
Deeper soils, good grass cover dominated by hairy wildrye, junegrass.....		25
25. Shrubby cinquefoil dominant in stand.....	SACMA3	
Grass cover extensive, dominated by hairy wildrye, junegrass, and brome....	SACMA5	

### Shrub dominated communities

1. Timberline communities dominated by whitebark pine, subalpine fir, grouseberry, or willow communities with marsh marigold, wandering daisy or globeflower in understory.....		2
Riparian communities adjacent to streams or rivers.....		5
2. Trees present in community (whitebark pine, subalpine fir) or grouseberry dominated....		3
Moist seepage areas at treeline dominated by globeflower, wandering daisy or marsh marigold in understory.....	SACMB6	
3. Trees (subalpine fir, whitebark pine) on site.....		4
Grouseberry dominated shrubland.....	SACMB7	
4. Whitebark pine present.....	SACMB3	
Subalpine fir present.....	SACMB8	
5. Very wet sites with water sedge or horsetail dominated understories.....		6
Drier sites with tufted hairgrass, california oatgrass, bog sedge, hairy wildrye, rough fescue Kentucky bluegrass, dandelion dominated understories.....		7
6. Water sedge dominated understory.....	SACFB1, SACMB1	
Horsetail dominated understory.....	SACFB2	
7. Grazed communities dominated by clover and dandelion in understory.....	SACFB5	
Ungrazed sites dominated by native forbs and grasses in understory.....		8
8. Shrubland communities on seepage areas on south facing slopes with shallow soils, dominated by bog birch and juniper.....	SACMB5	
Meadows and lowland shrublands dominated by rough fescue, bog sedge, california oatgrass, tufted hairgrass, hairy wildrye or sedge in the understory.....		9
9. Rough fescue dominates the understory.....		10
Tufted hairgrass, california oatgrass, bog sedge, sedge, hairy wildrye dominate.....		11
10. Rough fescue and bog sedge dominate understory higher elevations.....	SACFB10	
Rough fescue dominates, bog sedge not present lower elevations.....	SACFB9	
11. Moister sites with deep fluvial deposits dominated by tufted hairgrass, sedge, or fringed brome in understory.....		12
Drier sites which are well drained at the surface dominated by hairy wildrye, sedge, bog sedge or california oatgrass in understory.....		14
12. Tufted hairgrass or sedge dominated understory.....		13
Fluvial areas with Fringed brome dominated understory, lower elevation.....	SACFB11	
13. Tufted hairgrass dominates understory.....	SACFB4	
Graceful sedge and other sedge species dominate understory.....	SACFB3, SASMB2	
14. Modal sites with hairy wildrye and sedge dominating understory.....		15
Sites dominated by california oatgrass or bog sedge in understory.....		16
15. Hairy wildrye dominates understory.....	SACFB7, SACMB3	
Graceful sedge and other sedge species dominate understory.....	SACFB3, SASMB2	
16. California oatgrass dominates understory.....	SACFB6	
Bog sedge dominates understory.....	SACFB8, SACMB4	

## SACFA1. Water sedge-Beaked sedge meadows (*Carex aquatilis*-*Carex rostrata*)

**n=11** This community type is found in all ecodistricts of the subalpine. Wet conditions and periodic flooding result in the formation of water sedge-beaked sedge meadows. Bog birch and willow will invade into the drier edges of these meadows to form the Willow-bog birch/Water sedge community type.

These community types are quite productive producing nearly 2000 kg/ha of forage, but the high water table in the spring and summer when these meadows are most palatable limits livestock use. A study in the Yukon found that crude protein on these meadows declined from a high of 10% in May to less than 5% in September (Bailey et al. 1992). As a result, these meadows would be rated as secondary or non-use range.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
SMOOTH WILLOW ( <i>Salix glauca</i> )	1	0-7	27
BOG BIRCH ( <i>Betula glandulosa</i> )	3	0-20	46
<b>FORBS</b>			
ARROW LEAVED COLTSFOOT ( <i>Petasites sagittatus</i> )	7	0-14	18
<b>GRASSES</b>			
WATER SEDGE ( <i>Carex aquatilis</i> )	23	0-62	64
BEAKED SEDGE ( <i>Carex rostrata</i> )	11	0-42	55
ROCKY-GROUND SEDGE ( <i>Carex saxatilis</i> )	9	0-70	18
TUFTED HAIRGRASS ( <i>Deschampsia cespitosa</i> )	2	0-5	64

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:  
HYGRIC

NUTRIENT REGIME  
PERMESOTROPHIC

ELEVATION:  
1750M

SOIL DRAINAGE:  
POORLY

### FORAGE PRODUCTION IN KG/HA

GRASS 1215  
FORBS 774  
TOTAL 1721 \* ESTIMATE

SUGGESTED GRAZING CAPACITY  
NON-USE

## SACFA2. Tufted bulrush

*(Scirpus cespitosus)*

**n=12** This community type occurs on subhydryc to hydric, lower subalpine sites (1490-1870 m) on nearly level slopes. Corns and Achuff (1982), described this community type on poorly drained soils in the valleys of Banff and Jasper National Park. The soils are dominated by Terric Mesisols and Orthic Gleysols. They felt this community type was successional mature.

This community type and the Water sedge-Beaked sedge both occupy poorly drained sites with free standing water, but the dominance of tufted bulrush appears to indicate acidic boggy and peaty sites (Scoggan 1978).

### PLANT COMPOSITION CANOPY COVER(%)

MEAN    RANGE    CONST.

#### SHRUBS

SHRUBBY CINQUEFOIL <i>(Potentilla fruticosa)</i>	6	0-35	75
BOG BIRCH <i>(Betula glandulosa)</i>	3	0-10	50

#### FORBS

ELEPHANT'S HEAD <i>(Pedicularis groenlandicum)</i> 1		0-5	58
WOOLY EVERLASTING <i>(Antennaria lanata)</i>	1	0-5	42

#### GRASSES

TUFTED BULRUSH <i>(Scirpus cespitosus)</i>	61	18-85	100
WATER SEDGE <i>(Carex aquatilis)</i>	2	0-10	42
RUSH-LIKE SEDGE <i>(Carex scirpoidea)</i>	1	0-10	17

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:  
HYGRIC-SUBHYGRIC

NUTRIENT REGIME:  
PERMESOTROPHIC

ELEVATION:  
1678(1490-1870) M

SOIL DRAINAGE:  
POORLY

### FORAGE PRODUCTION KG/HA

NOT AVAILABLE

SUGGESTED GRAZING CAPACITY

NON-USE



### SACFA3. Sedge-Cottongrass

(*Carex spp.-Eriophorum spp.*)

n=7 Corns and Achuff (1982), described a cottongrass dominated community on hydric sites in the Upper subalpine on level to gentle slopes. They found the cottongrass communities to form on depressional areas where the snow melts late and seepage is received throughout the growing season. Cottongrass is also characteristic of muskogs and boggy marshes. It appears that this community is located on better drained areas adjacent to tufted bulrush dominated community type. The high acidity of the soil appears to favour the growth of cottongrass, rush-like sedge and rocky ground sedge over water sedge.

#### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

##### SHRUBS

WILLOW SPP. ( <i>Salix spp.</i> )	4	1-11	100
BOG BIRCH ( <i>Betula glandulosa</i> )	1	0-5	43

##### FORBS

LANCED -LEAVED PAINT BRUSH ( <i>Castilleja occidentalis</i> )	2	0-10	57
WOOLLY EVERLASTING ( <i>Antennaria lanata</i> )	1	0-4	57
COMMON HORSETAIL ( <i>Equisetum arvense</i> )	3	0-22	14

##### GRASSES

RUSH-LIKE SEDGE ( <i>Carex scirpoidea</i> )	8	0-25	57
ROCKY-GROUND SEDGE ( <i>Carex saxatilis</i> )	12	0-80	29
SHORT SEDGE ( <i>Carex curta</i> )	7	0-50	14
SHEATHED COTTONGRASS ( <i>Eriophorum vaginatum</i> )	1	0-10	14
TALL COTTONGRASS ( <i>Eriophorum polystachion</i> )	5	0-18	57

#### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:

HYDRIC-SUBHYGRIC

NUTRIENT REGIME:

EUTROPHIC

ELEVATION:

1586M

SOIL DRAINAGE:

POORLY

#### FORAGE PRODCUTION KG/HA

NOT AVAILABLE

SUGGESTED GRAZING CAPACITY

NON-USE

## SACFA4. Tufted hairgrass-Sedge (*Deschampsia cespitosa*-*Carex* spp.)

**n=18** This community type is located on moist sites that are better drained and slightly drier than the pure sedge meadows. Willoughby(2001), found that tufted hairgrass is a common plant species on lowland sites in the valley bottoms of the Upper Foothills subregion . Willoughby (1992), found when this community type is protected from grazing for 25-30 years, willow and bog birch expand and tufted hairgrass and sedge decline. The decline in graminoid cover also results in a decline in available forage production . Continuous heavy grazing causes hairgrass to decline and the site will be invaded by Kentucky bluegrass and dandelion.

Bork (1994), found this community type to be the most productive type described in Willmore Wilderness park. Forage production averages over 2000 kg/ha and can vary from (800-3300 kg/ha). It is interesting to note that forage production on this community type declines from an average of 2200 kg/ha in the Upper Foothills to 1555 kg/ha in the Subalpine subregion. The shorter growing season and colder climate may account for this change in forage productivity. These community types when located next to backcountry campsites will be utilized by horses.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
BARCLAY'S WILLOW ( <i>Salix barclayi</i> )	2	0-7	22
<b>FORBS</b>			
SLENDER BLUE BEARDSTONGUE ( <i>Penstemon procerus</i> )	2	0-9	72
VEINY MEADOW RUE ( <i>Thalictrum venulosum</i> )	5	0-14	50
YARROW ( <i>Achillea millefolium</i> )	4	0-14	94
SMOOTH LEAVED CINQUEFOIL ( <i>Potentilla diversifolia</i> )	12	0-43	72
OLD MAN'S WHISKERS ( <i>Geum triflorum</i> )	1	0-6	22
ALPINE GOLDENROD ( <i>Solidago multiradiata</i> )	1	0-13	72
<b>GRASSES</b>			
GRACEFUL SEDGE ( <i>Carex praegracilis</i> )	38	0-85	94
TUFTED HAIRGRASS ( <i>Deschampsia cespitosa</i> )	31	12-57	100
SLENDER WHEATGRASS ( <i>Agropyron trachycaulum</i> ) <sup>2</sup>		0-12	33
SHEEP FESCUE ( <i>Festuca saximontana</i> )	3	0-16	33
HAIRY WILD RYE ( <i>Elymus innovatus</i> )	3	0-20	50

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:  
MESIC TO HYGRIC

NUTRIENT REGIME:  
MESOTROPHIC TO PERMESOTROPHIC

ELEVATION:  
1896(1630-2130) M

SOIL DRAINAGE:  
MOD. WELL TO VERY POORLY

SLOPE:  
3(1-5)%

ASPECT:  
VARIABLE

### FORAGE PRODUCTION KG/HA

GRASS 997 (532-1923)  
FORBS 288(0-928)  
TOTAL 1284 (532-2118)

SUGGESTED GRAZING CAPACITY 0.7 HA/AUM
--

## SACFA5. Sedge-Tufted hairgrass

(*Carex* spp.-*Deschampsia cespitosa*)

n=8 This community type was described in the Job Lake, Blackstone-Wapiabi Forest Land Use Zones. These areas are extensively utilized by equestrian backcountry users. This community type appears to develop from moderate to heavy grazing pressure on a Tufted hairgrass-Sedge community. Continued heavy grazing pressure appears to cause a further decline in tufted hairgrass to form the Sedge-Rocky Mtn. fescue-Alpine timothy and Sedge-Slender wheatgrass-Fringed brome/Forb community types. If the seed source becomes available these sites have the potential to be invaded by Kentucky bluegrass, timothy, clover and dandelion if the grazing pressure continues.

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

#### SHRUBS

##### BOG BIRCH

(*Betula glandulosa*) 1 0-2 13

#### FORBS

##### ALPINE BISTORT

(*Polygonum viviparum*) 1 0-4 75

##### VEINY MEADOW RUE

(*Thalictrum venulosum*) 4 0-16 50

##### YARROW

(*Achillea millefolium*) 7 0-6 88

##### SMOOTH LEAVED CINQUEFOIL

(*Potentilla diversifolia*) 9 0-18 75

##### PURPLE AVENS

(*Geum rivale*) 7 0-42 63

##### GREEN SORREL

(*Rumex acetosa*) 2 0-8 75

#### GRASSES

##### GRACEFUL SEDGE

(*Carex praegracilis*) 58 19-98 100

##### TUFTED HAIRGRASS

(*Deschampsia cespitosa*) 18 3-34 100

##### KENTUCKY BLUEGRASS

(*Poa pratensis*) 4 0-15 75

##### ALPINE FESCUE

(*Festuca brachyphylla*) 3 0-15 63

##### CALIFORNIA OATGRASS

(*Danthonia californica*) 3 0-9 50

### ENVIRONMENTAL VARIABLES

#### MOISTURE REGIME:

MESIC -SUBHYGRIC

#### NUTRIENT REGIME:

MESOTROPHIC TO PERMESOTROPHIC

#### ELEVATION:

1866(1832-1895) M

#### SOIL DRAINAGE:

WELL TO MOD. WELL

### FORAGE PRODUCTION KG/HA

GRASS 785(341-1369)

FORBS 146(0-443)

TOTAL 931(412-1738)

SUGGESTED GRAZING CAPACITY

1.0 HA/AUM

**SACFA6. Sedge-Rocky Mountain fescue-Alpine timothy**  
*(Carex praegracilis-Festuca saximontana-Phleum alpina)*

n=6 This community type appears to arise from grazing of a modal Tufted hairgrass-Sedge community type. The six sites described in this community were all located next to outfitter campsites and had been heavily grazed by horses. The heavy grazing pressure causes tufted hairgrass to decline and allows non-native plants such as Kentucky bluegrass and dandelion to invade onto the site. The heavy grazing pressure also appears to change the moisture regime of the site. Many of the plant species on the site, Rocky mountain fescue, Alpine timothy, Alpine bluegrass, junegrass and hairy wildrye are better adapted to well drained, drier conditions. Perhaps, the removal of litter causes the water to drain away from the site more rapidly. Bork (1994), noticed this on similar sites in Willmore Wilderness park. This community type maybe grazed heavier than the previously described Sedge-Tufted hairgrass community type or it could be drier and the grazing pressure shifts the community to one dominated by more drought resistant species

**PLANT COMPOSITION CANOPY COVER(%)**  
 MEAN RANGE CONST.

**SHRUB**

BOG BIRCH  
*(Betula glandulosa)* 3 0-12 33

**FORBS**

SLENDER BLUE BEARDTONGUE  
*(Penstemon procerus)* 2 0-3 83  
 YARROW  
*(Achillea millefolium)* 6 1-9 100  
 GRACEFUL CINQUEFOIL  
*(Potentilla gracilis)* 4 0-13 83  
 CHICKWEED  
*(Cerastium arvense)* 1 0-3 83  
 DANDELION  
*(Taraxacum officinale)* 2 0-7 83  
 NORTHERN VALERIAN  
*(Valeriana dioica)* 2 0-5 50

**GRASSES**

ROCKY MOUNTAIN FESCUE  
*(Festuca saximontana)* 15 0-30 83  
 TUFTED HAIRGRASS  
*(Deschampsia cespitosa)* 4 0-20 50  
 KENTUCKY BLUEGRASS  
*(Poa pratensis)* 5 0-14 67  
 GRACEFUL SEDGE  
*(Carex praegracilis)* 9 0-42 50  
 ALPINE TIMOTHY  
*(Phleum commutatum)* 6 0-15 50

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
 SUBMESIC-SUBHYGRIC  
 NUTRIENT REGIME:  
 SUBMESOTROPHIC TO MESOTROPHIC  
 ELEVATION:  
 1886(1634-2130) M  
 SOIL DRAINAGE:  
 WELL TO IMPERFECTLY  
 SLOPE: 6(2-10)%  
 ASPECT: VARIABLE

**FORAGE PRODUCTION KG/HA**

GRASS 661 (294-1121)  
 FORB 326 (0-524)  
 SHRUB 7 (0-14)  
 TOTAL 994 (729-1341)

SUGGESTED GRAZING CAPACITY  
 0.9 HA/AUM

## SACFA7. Sedge-Slender wheatgrass-Fringed brome/Forbs

(*Carex praeegracilis*-*Agropyron trachycaulum*-*Bromus ciliatus*/Forbs)

**n=18** This community type appears to arise from grazing of a modal Tufted hairgrass-Sedge community type. All the sites described in this community were found adjacent to outfitter campsites in the Job Lake, Blackstone-Wapiabi forest land use zones. The heavy grazing pressure causes tufted hairgrass to decline and allows sedges, slender whatgrass and fringed brome to increase. This community type maybe slightly moister than the previously described Sedge-Rocky mountain fescue-Alpine timothy community. As a result there is succession to more mesic loving plants rather than the drought tolerant plants described in the previous community type.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
WILLOW SPP. ( <i>Salix spp.</i> )	1	0-10	33
<b>FORBS</b>			
ALPINE BISTORT ( <i>Polygonum viviparum</i> )	7	0-59	67
YARROW ( <i>Achillea millefolium</i> )	4	0-21	83
SMOOTH LEAVED CINQUEFOIL ( <i>Potentilla diversifolia</i> )	11	0-37	72
ALPINE GOLDENROD ( <i>Solidago multiradiata</i> )	4	0-28	72
STRAWBERRY ( <i>Fragaria virginiana</i> )	6	0-29	61
FALSE-DANDELION ( <i>Agoseris glauca</i> )	3	0-22	72
<b>GRASSES</b>			
SLENDER WHEATGRASS ( <i>Agropyron trachycaulum</i> )	4	0-15	50
TUFTED HAIRGRASS ( <i>Deschampsia cespitosa</i> )	4	0-20	33
KENTUCKY BLUEGRASS ( <i>Poa pratensis</i> )	6	0-26	78
GRACEFUL SEDGE ( <i>Carex praeegracilis</i> )	50	2-94	100
FRINGED BROME ( <i>Bromus ciliatus</i> )	14	0-50	78
HAIRY WILD RYE ( <i>Elymus innovatus</i> )	8	0-46	44

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:

MESIC-HYGRIC

NUTRIENT REGIME:

MESOTROPHIC-PERMESOTROPHIC

ELEVATION:

1926(1830-2076) M

SOIL DRAINAGE:

WELL TO MODERATELY WELL

### FORAGE PRODUCTION KG/HA

GRASS 539(162-944)

FORB 156(0-370)

SHRUB 7(0-44)

TOTAL 702(354-1094)

SUGGESTED GRAZING CAPACITY

1.3 HA/AUM OR 2.8 ACRES/AUM

**SACFA8. California oatgrass-Sedge**  
(*Danthonia californica*-*Carex* spp.)

**n=5** This community is very similar to a community type described by Willoughby(2001) in the Upper Foothills subregion. Corns and Achuff (1982), described a Willow/California oatgrass dominated community type in Banff and Jasper National Park. It appears dry, gravelly or stony soils, with a fluctuating water table support this moderately productive grassland. Small pockets of this community type occur throughout the Subalpine subregion. In the Yukon these small meadows were found to form in depressions which appeared to act as pronounced frost pockets (Bailey et al. 1992). The cold air drainage and poor nutrient quality of the soil limits the forage productivity of these sites.

**PLANT COMPOSITION CANOPY COVER(%)**

**MEAN RANGE CONST.**

**SHRUBS**

WILLOW SPP. ( <i>Salix</i> spp.)	3.8	0-13	60
SHRUBBY CINQUEFOIL ( <i>Potentilla fruticosa</i> )	3	0-10	60

**FORBS**

TALL LARKSPUR ( <i>Delphinium glaucum</i> )	4	0-11	80
WILD STRAWBERRY ( <i>Fragaria virginiana</i> )	6	0-25	60
SMOOTH LEAVED CINQUEFOIL ( <i>Potentilla diversifolia</i> )	11	0-31	80
YARROW ( <i>Achillea millefolium</i> )	4	0-20	80
VEINY MEADOW RUE ( <i>Thalictrum venulosum</i> )	10	0-25	60
SLENDER BLUE BEARDTONGUE ( <i>Penstemon procerus</i> )	9	0-30	60

**GRASSES**

CALIFORNIA OATGRASS ( <i>Danthonia californica</i> )	37	15-60	100
GRACEFUL SEDGE ( <i>Carex praegracilis</i> )	24	0-63	40
SLENDER WHEATGRASS ( <i>Agropyron trachycaulum</i> )	3	0-15	20
HAIRY WILD RYE ( <i>Elymus innovatus</i> )	11	0-29	60
SPIKED TRisetum ( <i>Trisetum spicatum</i> )	1	0-5	60
BOG SEDGE ( <i>Kobresia myosuroides</i> )	4	0-19	20

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
SUBMESIC

NUTRIENT REGIME:  
MESOTROPHIC

ELEVATION:  
1830(1380-2100)M

SOIL DRAINAGE:  
MODERATELY WELL TO WELL

**FORAGE PRODUCTION KG/HA**

GRASS	921
FORBS	352
TOTAL	1273 * ESTIMATE

**SUGGESTED GRAZING CAPACITY**  
0.7 HA/AUM OR 1.6 ACRES/AUM

## SACFA9. Rough fescue-Hairy wildrye-Sedge

(*Festuca scabrella*-*Elymus innovatus*-*Carex spp.*)

n=13 This community was described in the Panther Corners Forest Land Use Zone and in Willmore Wilderness Park on level to undulating ridges, terraces and lower slope positions with Orthic Eutric Brunisolic soils. It is very similar to the Rough fescue-Hairy wildrye community described by Willoughby(2001) in the Upper Foothills subregion and the Rough fescue-Wheatgrass-Hairy wildrye community described by Morgantini and Russell (1983) on Ribbon flats just north of the Panther Corners. This community type is moderately productive and one of the most important communities for wintering elk (Morgantini and Russell 1983). An examination of winter elk diets found that rough fescue made up 45 to 60% of their food intake during the months of December, January and March (Morgantini and Russell 1983). Care must be taken that this community type is not over-utilized by horses and that sufficient forage is left for overwintering elk.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
SHRUBBY CINQUEFOIL ( <i>Potentilla fruticosa</i> )	8	0-31	92
<b>FORBS</b>			
TALL LARKSPUR ( <i>Delphinium glaucum</i> )	1	0-7	62
WILD STRAWBERRY ( <i>Fragaria virginiana</i> )	2	0-7	77
GRACEFUL CINQUEFOIL ( <i>Potentilla gracilis</i> )	4	0-11	62
YARROW ( <i>Achillea millefolium</i> )	3	0-7	92
OLD MAN'S WHISKERS ( <i>Geum triflorum</i> )	5	0-12	62
AMERICAN VETCH ( <i>Vicia americana</i> )	3	0-7	77
<b>GRASSES</b>			
ROUGH FESCUE ( <i>Festuca scabrella</i> )	18	10-34	100
SEDGE ( <i>Carex spp.</i> )	8	1-14	100
SLENDER WHEATGRASS ( <i>Agropyron trachycaulum</i> )	3	0-13	69
HAIRY WILDRYE ( <i>Elymus innovatus</i> )	6	1-18	100
JUNEGRASS ( <i>Koeleria macrantha</i> )	1	0-4	85

MOISTURE REGIME:

SUBXERIC TO MESIC

NUTRIENT REGIME:

MESOTROPHIC -PERMESOTROPHIC

ELEVATION:

1786(1600-2150)M

SOIL DRAINAGE:

WELL TO RAPIDLY

SLOPE:

32(10-60)%

ASPECT:

SOUTHERLY

### FORAGE PRODUCTION KG/HA

GRASS	1487(454-3056)
FORBS	689(302-1792)
SHRUB	167(0-968)
TOTAL	2343(1284-4060)

SUGGESTED GRAZING CAPACITY  
0.4 HA/AUM OR 0.9 ACRES/AUM

### ENVIRONMENTAL VARIABLES

## SACFA10. Sedge-Hairy wildrye

(*Carex spp.-Elymus innovatus*)

**n=13** This community was described in the Panther Corners Forest Land Use Zone on steep west and south facing slopes at higher elevations. It occupies sites that are similar to the Fringed sage/Sedge-Junegrass community, but this community type is found at elevations averaging over 2000 meters. This community type is similar to the hairy wildrye dominated communities described by Corns and Achuff (1982) at higher elevations in the subalpine of the Central Mountains ecodistrict (SACMA,4,5,6). This community type was only lightly utilized by elk. In contrast the lower elevation Fringed sage/Sedge-Junegrass exhibited signs of heavy elk use. The higher elevation of this community may limit access to wildlife in this area. It is possible that if this community type was heavily grazed it may resemble the Fringed sage/Sedge-Junegrass community described at lower elevations.

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

#### SHRUBS

SHRUBBY CINQUEFOIL <i>(Potentilla fruticosa)</i>	4	0-14	77
---	---	------	----

#### FORBS

TALL LARKSPUR <i>(Delphinium glaucum)</i>	1	0-5	54
WILD STRAWBERRY <i>(Fragaria virginiana)</i>	2	0-16	39
GRACEFUL CINQUEFOIL <i>(Potentilla gracilis)</i>	2	0-5	100
YARROW <i>(Achillea millefolium)</i>	1	0-4	77
OLD MAN'S WHISKERS <i>(Geum triflorum)</i>	2	0-18	23
AMERICAN VETCH <i>(Vicia americana)</i>	1	0-3	39
BEARBERRY <i>(Arctostaphylos uva-ursi)</i>	4	0-61	8

#### GRASSES

ROUGH FESCUE <i>(Festuca scabrella)</i>	2	0-9	31
SEDGE <i>(Carex spp.)</i>	20	3-48	84
POA <i>(Poa spp.)</i>	2	0-5	54
HAIRY WILDRYE <i>(Elymus innovatus)</i>	8	0-29	100
JUNEGRASS <i>(Koeleria macrantha)</i>	1	0-3	46
SMOOTH BROME <i>(Bromus inermis)</i>	3	0-6	85

### ENVIRONMENTAL VARIABLES

#### MOISTURE REGIME:

XERIC TO MESIC

#### NUTRIENT REGIME:

SUBMESOTROPHIC

#### ELEVATION:

2029( 1650-2300)M

#### SOIL DRAINAGE:

WELL TO RAPIDLY

#### ASPECT:

SOUTHERLY AND WESTERLY

#### SLOPE:

36(0-50)%

### FORAGE PRODUCTION KG/HA

GRASS	771 (460-1168)
FORBS	390 (160-1010)
SHRUB	284 (0-1712)
TOTAL	1444 (676-3150)

SUGGESTED GRAZING CAPACITY

NON-USE



**SACFA11. Blunt sedge-Junegrass/Bearberry**  
*(Carex obtusata-Koeleria macrantha/Arctostaphylos uva-ursi)*

**n=3** This community type occurs on steep south facing slopes, with shallow soils, overlying sandstone bedrock. The majority of the vegetation are composed of drought tolerant species bearberry and junegrass. The inaccessibility and fragile nature of the soils make this community type unsuitable for grazing. This community is very similar to the Junegrass/Sage community described by Willoughby(2001) in the Upper Foothills subregion and the Low northern Sedge/Bearberry community described by Lane et al. (2000) in the Lower Foothills subregion on the south facing slopes of the Athabasca River valley.

**PLANT COMPOSITION CANOPY COVER(%)**  
**MEAN RANGE CONST.**

<b>SHRUBS</b>			
SHRUBBY CINQUEFOIL <i>(Potentilla fruticosa)</i>	3	2-3	100
<b>FORBS</b>			
<b>BEARBERRY</b>			
<i>(Arctostaphylos uva-ursi)</i>	11	0-16	67
SHOWY LOCOWEED <i>(Oxytropis splendens)</i>	10	9-10	100
SWEET FLOWERED ANDROSACE <i>(Androsace chamaejasme)</i>	7	0-19	67
ALPINE GOLDENROD <i>(Solidago multiradiata)</i>	1	0-2	67
COMMON YARROW <i>(Achillea millefolium)</i>	7	2-13	100
<b>GRASSES</b>			
<b>JUNEGRASS</b>			
<i>(Koeleria macrantha)</i>	10	1-23	100
SHEEP FESCUE <i>(Festuca saximontana)</i>	7	0-12	67
BLUNT SEDGE <i>(Carex obtusata)</i>	3	1-7	100
HAIRY WILDRYE <i>(Elymus innovatus)</i>	3	0-6	67
SLENDER WHEATGRASS <i>(Agropyron trachycaulum)</i>	3	1-5	100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
XERIC TO SUBXERIC

NUTRIENT REGIME:  
MESOTROPHIC

ELEVATION:  
1990(1950-2070)M

SOIL DRAINAGE: RAPIDLY

SLOPE: 40(30-60)%

ASPECT: SOUTHERLY

**FORAGE PRODUCTION KG/HA**

GRASS 1235(1196-1274)  
 FORB 264(148-380)  
 SHRUB 13(0-26)  
 TOTAL 1512(1370-1654)

SUGGESTED GRAZING CAPACITY  
 NON-USE

## SACFA12. Fringed sage/White scaled sedge-Junegrass

*(Artemisia frigida/Carex xerantica-Koeleria macrantha)*

n=5 This community type occurs on steep south facing slopes, with shallow soils. It is very similar to the previously described Sedge-Junegrass/Bearberry community type, but lacks the cover of bearberry. The lack of bearberry cover in this community type may indicate that the soils of this type are better developed and slightly moister. The inaccessibility and fragile nature of the soils make this community type unsuitable for domestic livestock grazing. This community type is important winter habitat for migrating elk. The steepness of the slope and the southerly aspect limit snow accumulation and allows access to the forage supply. This community is very similar to the Junegrass/Sage community described by Willoughby(2001) in the Upper Foothills subregion.

### PLANT COMPOSITION CANOPY COVER(%) MEAN RANGE CONST.

#### SHRUBS

SHRUBBY CINQUEFOIL <i>(Potentilla fruticosa)</i>	2	0-4	80
PRICKLY ROSE <i>(Rosa acicularis)</i>	3	0-8	100

#### FORBS

BEARBERRY <i>(Arctostaphylos uva-ursi)</i>	1	0-2	20
EARLY YELLOW LOCOWEED <i>(Oxytropis sericea)</i>	3	1-4	100
FRINGED SAGE <i>(Artemisia frigida)</i>	6	1-15	100
AMERICAN VETCH <i>(Vicia americana)</i>	3	0-9	60
NORTHERN BEDSTRAW <i>(Galium boreale)</i>	5	1-14	100

#### GRASSES

JUNEGRASS <i>(Koeleria macrantha)</i>	5	3-7	100
WESTERN WHEATGRASS <i>(Agropyron smithii)</i>	4	0-9	80
WHITE SCALED SEDGE <i>(Carex xerantica)</i>	10	6-12	100
ROUGH FESCUE <i>(Festuca scabrella)</i>	3	0-6	80
SLENDER WHEATGRASS <i>(Agropyron trachycaulum)</i>	3	0-7	60

MOISTURE REGIME:

XERIC TO SUBXERIC

NUTRIENT REGIME:

SUBMESOTROPHIC

SLOPE: 53(45-60)%

ASPECT: SOUTHERLY

ELEVATION:

1790(1650-1900)M

SOIL DRAINAGE: RAPIDLY

### FORAGE PRODUCTION KG/HA

Grass	1133(650-2206)
FORB	545(202-890)
SHRUB	250(0-556)
TOTAL	1928(936-3096)

SUGGESTED GRAZING CAPACITY  
NON-USE

### ENVIRONMENTAL VARIABLES

**SACFA13. Sedge-Bog sedge-Tufted hairgrass**  
(*Carex praegracilis*-*Kobresia myosuroides*-*Deschampsia cespitosa*)

**n=6** This community type is found on moist lowland sites at higher elevations in the Central and Northern Foothills. The presence of bog sedge appears to indicate the transition to the higher Alpine subregion. Indeed, Ogilvie (1969) described bog sedge dominated community types at higher elevations in the Alpine subregion.

The presence of bog sedge may also represent the transition between the foothills ecodistricts to the rocky mountain ecodistricts. Corns and Achuff (1982), described bog sedge dominated community types in the Subalpine subregion of Banff and Jasper National Parks.

The forage production on this community type is only moderate. Perhaps, the higher elevation and colder climate which favours the growth of bog sedge limits the total productivity of the site. Camping and grazing of these communities by horses should be restricted.

**PLANT COMPOSITION CANOPY COVER(%)**

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
SHRUBBY CINQUEFOIL ( <i>Potentilla fruticosa</i> )	1	0-4	67
<b>FORBS</b>			
ALPINE GOLDENROD ( <i>Solidago multiradiata</i> )	7	0-15	100
STRAWBERRY ( <i>Fragaria virginiana</i> )	1	0-4	50
GRACEFUL CINQUEFOIL ( <i>Potentilla gracilis</i> )	3	0-10	33
YARROW ( <i>Achillea millefolium</i> )	5	0-15	100
ALPINE BISTORT ( <i>Polygonum viviparum</i> )	6	0-13	100
SMOOTH LEAVED CINQUEFOIL ( <i>Potentilla diversifolia</i> )	12	0-31	83
<b>GRASSES</b>			
GRACEFUL SEDGE ( <i>Carex praegracilis</i> )	40	14-60	100
BOG SEDGE ( <i>Kobresia myosuroides</i> )	20	10-28	100
TUFTED HAIRGRASS ( <i>Deschampsia cespitosa</i> )	13	5-37	83
ROCKY MOUNTAIN FESCUE ( <i>Festuca saximontana</i> )	3	0-10	33
ALPINE BLUEGRASS ( <i>Poa alpina</i> )	2	0-13	50
HAIRY WILDRYE ( <i>Elymus innovatus</i> )	7	0-30	100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
MESIC-SUBHYGRIC

NUTRIENT REGIME:  
MESOTROPHIC

ELEVATION:  
1900(1832-2438)M

SOIL DRAINAGE:  
IMPERFECTLY

SLOPE:  
1%

ASPECT:  
NORTHEAST

**FORAGE PRODUCTION KG/HA**

GRASS 582 (310-1002)  
FORB 158 (58-272)  
TOTAL 740 (582-1060)

SUGGESTED GRAZING CAPACITY  
NON-USE

## SACFA14. White mountain avens/Bog sedge

*(Dryas integrifolia/Kobresia myosuroides)*

**n=3** This community type occupies shallow, stoney, wind exposed sites. It represents the transitional community between the bog sedge and white mountain avens community types described by Ogilvie (1969) and Corns and Achuff (1982) on windswept ridges in the Alpine and Subalpine subregions of the Rocky Mountains. The microsite conditions are very similar to higher elevation sites in the Rocky Mountains allowing this community to form in the lower Central Foothills.

The poor soil conditions limits the forage productivity and amount of regrowth after grazing. Often this community type is important winter range for bighorn sheep, because this community type remains snow free for much of the winter.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>FORBS</b>			
WHITE MOUNTAIN AVENS			
<i>(Dryas integrifolia)</i>	46	0-41	67
ALPINE BISTORT			
<i>(Polygonum viviparum)</i>	6	3-9	100
ALPINE HEDYSARUM			
<i>(Hedysarum alpinum)</i>	2	0-4	100
ALPINE MILKVETCH			
<i>(Astragalus alpinum)</i>	1	0-3	33
<b>GRASSES</b>			
BOG SEDGE			
<i>(Kobresia myosuroides)</i>	11	1-32	100
GRACEFUL SEDGE			
<i>(Carex praegracilis)</i>	6	0-17	33
HAIRY WILDRYE			
<i>(Elymus innovatus)</i>	5	0-9	67
<b>LICHENS</b>			
REINDEER LICHEN			
<i>(Cladina spp.)</i>	17	0-40	67

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:  
XERIC-SUBXERIC

NUTRIENT REGIME:  
SUBMESOTROPHIC

ELEVATION:  
1912(1878-1981)M

SOIL DRAINAGE: RAPIDLY

SLOPE: 30%

ASPECT: SOUTHERLY

### FORAGE PRODUCTION KG/HA

GRASS 517(89-945)  
FORB 245(200-290)  
TOTAL 762(289-1235)

SUGGESTED GRAZING CAPACITY  
NON-USE

**SACFA15. Creeping red fescue-Sedge**  
*(Festuca rubra-Carex spp.)*

**n=1** This community type was described in the Blackstone-Wapiabi forest land use zone. It represents an old wellsite road that has been seeded to creeping red fescue. The creeping red fescue was probably used as the reclamation seed to stabilize the road from erosion. Presently, it is the recommendation of the forest service to use native seed in reclamation of these areas in the backcountry. Many of these agronomic mixes are highly invasive on the surrounding vegetation and there is the potential to introduce noxious weeds.

**PLANT COMPOSITION CANOPY COVER(%)**  
**MEAN RANGE CONST.**

**SHRUBS**

WILLOW  
*(Salix barclayi)* 1 - 100

**FORBS**

PALMATE LEAVED COLTSFOOT  
*(Petasites palmatus)* 1 - 100  
 FIREWEED  
*(Epilobium angustifolium)* 1 - 100

**GRASSES**

CREEPING RED FESCUE  
*(Festuca rubra)* 63 - 100  
 GRACEFUL SEDGE  
*(Carex praegracilis)* 38 - 100  
 TUFTED HAIRGRASS  
*(Deschampsia cespitosa)* 2 - 100  
 TIMOTHY  
*(Phleum pratense)* 1 - 100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
 MESIC

NUTRIENT REGIME:  
 MESOTROPHIC

ELEVATION:  
 1832M

SOIL DRAINAGE:  
 WELL

SLOPE:  
 LEVEL

**FORAGE PRODUCTION KG/HA**

GRASS 1705  
 FORB 15  
 TOTAL 1720

SUGGESTED GRAZING CAPACITY  
 0.6 HA/AUM OR 1.2 AC/AUM

## SACFA16. Kentucky bluegrass-Sedge/Dandelion

*(Poa pratensis-Carex spp./Taraxacum officinale)*

n=1 This community type was described in the Job Lake forest land use zone. It represents a Tufted hairgrass-Sedge meadow that has been heavily grazed for a prolonged period of time. As a result there has been a decline in tufted hairgrass and other native plant species and an invasion of Kentucky bluegrass and dandelion. This community type is uncommon in the backcountry areas because of the lack of seed source for Kentucky bluegrass and dandelion. However, once established it is very competitive and will likely remain on the site. Kentucky bluegrass is very productive, but it quickly loses its nutrient quality in the dormant season. This loss of nutrient quality will impact wildlife utilizing the area.

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

#### SHRUBS

WILLOW ( <i>Salix barclayi</i> )	1	-	100
-------------------------------------	---	---	-----

#### FORBS

ALPINE BISTORT ( <i>Polygonum viviparum</i> )	18	-	100
--	----	---	-----

GRACEFUL CINQUEFOIL ( <i>Potentilla gracilis</i> )	8	-	100
---	---	---	-----

SWEET FLOWERED ANDROSACE ( <i>Androsace chamaejasme</i> )	3	-	100
--	---	---	-----

ALPINE MILKVETCH ( <i>Astragalus alpinus</i> )	3	-	100
---	---	---	-----

MOUSE EARED CHICKWEED ( <i>Cerastium arvense</i> )	2	-	100
---	---	---	-----

#### GRASSES

KENTUCKY BLUEGRASS ( <i>Poa pratensis</i> )	36	-	100
--	----	---	-----

GRACEFUL SEDGE ( <i>Carex praegracilis</i> )	32	-	100
---	----	---	-----

SLENDER WHEATGRASS ( <i>Agropyron trachycaulum</i> )	5	-	100
---	---	---	-----

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:

MESIC

NUTRIENT REGIME:

PERMESOTROPHIC

ELEVATION:

1832M

SOIL DRAINAGE: WELL

SLOPE: LEVEL

### FORAGE PRODUCTION KG/HA

GRASS 380

FORB 224

TOTAL 604

SUGGESTED GRAZING CAPACITY

1.5 HA/AUM OR 3.4 AC/AUM

**SACFA17. Fireweed-Meadow rue/Sedge**  
(*Epilobium angustifolium*-*Thalictrum venulosum*/*Carex spp.*)

n=1 This community type was described in the Panther Corners forest land use zone adjacent to an abandoned air strip. The site was a moist site with a higher nutrient regime making it highly productive. This community type had not been extensively utilized by horses or wildlife and appears to be undergoing succession to a shrub dominated community type. Some invasion of agronomic species (smooth brome, bluegrass spp.) has occurred off the old airstrip into this community type.

**PLANT COMPOSITION** **CANOPY COVER(%)**  
MEAN RANGE CONST.

**SHRUBS**

WILLOW  
(*Salix spp.*) T - 100

**FORBS**

FIREWEED  
(*Epilobium angustifolium*)37 - 100

VEINY MEADOW RUE  
(*Thalictrum venulosum*) 22 - 100

YARROW  
(*Achillea millefolium*) 13 - 100

AMERICAN VETCH  
(*Vicia americana*) 5 - 100

NORTHERN BEDSTRAW  
(*Galium boreale*) 4 - 100

**GRASSES**

BLUEGRASS  
(*Poa spp.*) 1 - 100

SEDGE  
(*Carex spp.*) 21 - 100

SLENDER WHEATGRASS  
(*Agropyron trachycaulum*)1 - 100

SMOOTH BROME  
(*Bromus inermis*) 3 - 100

HAIRY WILDRYE  
(*Elymus innovatus*) 13 - 100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME: SUBHYGRIC

NUTRIENT REGIME:  
PERMESOTROPHIC

ELEVATION: 1650 M

SOIL DRAINAGE:  
MODERATELY WELL

SLOPE: LEVEL

**FORAGE PRODUCTION**

GRASS 1006

FORB 2134

TOTAL 3140

SUGGESTED GRAZING CAPACITY  
0.3 HA/AUM OR 0.7 AC/AUM

## SACFB1. Willow-Bog birch/Water sedge

(*Salix spp.*-*Betula glandulosa*/*Carex aquatilis*)

n=9 This shrub community appears on areas with very poor drainage. It is found in association with the wetter water sedge meadows. These sites are fairly productive but difficult to graze due to the moist ground conditions and heavy shrub cover which reduces access and mobility within the area. Increased flooding and prolonged waterlogging may result in the disappearance of willow and a transition to a water sedge meadow.

This community is similar to the water sedge-beaked sedge community in that it is found throughout the foothills and into the mountains. It maybe found in the Upper Foothills, Subalpine and lower Alpine subregions.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
WILLOW			
( <i>Salix spp</i> )	30	1-67	100
BOG BIRCH			
( <i>Betula glandulosa</i> )	18	0-44	100
<b>FORBS</b>			
ELEPHANT'S HEAD			
( <i>Pedicularis groenlandicum</i> )	1	0-6	78
SMOOTH ASTER			
( <i>Aster laevis</i> )	2	0-8	56
ALPINE BISTORT			
( <i>Polygonum viviparum</i> )	1	0-6	78
SMOOTH LEAVED CINQUEFOIL			
( <i>Potentilla diversifolia</i> )	2	0-15	33
ENTIRE LEAVED GROUNDSEL			
( <i>Senecio lugens</i> )	2	0-13	22
<b>GRASSES</b>			
WATER SEDGE			
( <i>Carex aquatilis</i> )	32	0-91	67
TUFTED HAIRGRASS			
( <i>Deschampsia cespitosa</i> )	4	0-26	89
GRACEFUL SEDGE			
( <i>Carex praegracilis</i> )	2	0-20	11
BALTIC RUSH			
( <i>Juncus balticus</i> )	5	0-17	56

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME :  
HYGRIC

NUTRIENT REGIME:  
PERMESOTROPHIC

ELEVATION:  
1760(1600-1950) M

SOIL DRAINAGE:  
MODERATELY WELL TO POORLY

SLOPE:  
6(2-5)%

ASPECT:  
SOUTHEASTERLY

### FORAGE PRODUCTION KG/HA

GRASS	1234(500-2320)
FORBS	129(0-354)
SHRUB	1150(0-2990)
TOTAL	2514(1870-3848)

SUGGESTED GRAZING CAPACITY  
NON-USE



## SACFB2. Willow/Horsetail

(*Salix spp./ Equisetum spp.*)

n=4 This community type occupies level to gently sloping, fluvial landforms at lower elevations in the subalpine. The sites are hygric and imperfectly to poorly drained. This community borders rivers and streams and is transitional to the spruce, subalpine fir, horsetail dominated forest.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
WILLOW SPP. ( <i>Salix spp.</i> )	32	5-56	100
BOG BIRCH ( <i>Betula glandulosa</i> )	5	0-20	60
<b>FORBS</b>			
VARIEGATED HORSETAIL ( <i>Equisetum variegatum</i> )	10	0-40	60
COMMON HORSETAIL ( <i>Equisetum arvense</i> )	10	0-25	60
WOOLLY EVERLASTING ( <i>Antennaria lanata</i> )	1	0-3	40
<b>GRASSES</b>			
SEDGE SPP. ( <i>Carex spp.</i> )	14	3-25	100
TUFTED HAIRGRASS ( <i>Deschampsia cespitosa</i> )	1	0-3	60

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME :  
SUBHYGRIC

NUTRIENT REGIME:  
PERMESOTROPHIC

ELEVATION:  
1512(1260-1810) M

SOIL DRAINAGE:  
POORLY

### FORAGE PRODUCTION KG/HA

NONE AVAILABLE

SUGGESTED GRAZING CAPACITY  
NON-USE

## SACFB3. Willow/Graceful sedge

(*Salix spp./ Carex praegracilis*)

n=6 This community type appears to represent a stage of succession onto tufted hairgrass meadows. When these communities are protected from disturbance (fire and grazing) willow and bog birch expand and tufted hairgrass declines. Willow growth also appears to favour the growth of tall forbs (veiny meadow rue, fireweed, aster) and slender wheatgrass. Fire has played a dominant role in controlling brush encroachment in the past and continued protection will allow continued shrub expansion, resulting in a decline in forage production.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
WILLOW SPP. ( <i>Salix spp.</i> )	35	18-55	100
BOG BIRCH ( <i>Betula glandulosa</i> )	1	0-3	50
<b>FORBS</b>			
SMOOTH LEAVED CINQUEFOIL ( <i>Potentilla diversifolia</i> )	8	0-44	50
STRAWBERRY ( <i>Fragaria virginiana</i> )	4	0-17	50
YARROW ( <i>Achillea millefolium</i> )	2	0-8	83
ALPINE BISTORT ( <i>Polygonum viviparum</i> )	2	0-10	50
SMALL LEAVED EVERLASTING ( <i>Antennaria parviflora</i> )	10	0-56	33
MEADOW RUE ( <i>Thalictrum venulosum</i> )	3	0-15	17
<b>GRASSES</b>			
GRACEFULSEDGE ( <i>Carex praegracilis</i> )	50	35-73	100
TUFTED HAIRGRASS ( <i>Deschampsia cespitosa</i> )	9	0-41	67
SLENDER WHEATGRASS ( <i>Agropyron trachycaulum</i> )	3	0-13	33
SPIKED TRisetum ( <i>Trisetum spicatum</i> )	3	0-16	33

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:  
SUBHYGRIC

NUTRIENT REGIME:  
PERMESOTROPHIC

ELEVATION:  
1876(1832-1985) M

SOIL DRAINAGE:  
POORLY

### FORAGE PRODUCTION KG/HA

GRASS 806(380-1369)  
FORBS 109(52-224)  
SHRUBS 3(0-10)  
TOTAL 919(604-1421)

SUGGESTED GRAZING CAPACITY  
1.0 HA/AUM OR 2.2 AC/AUM

## SACFB4. Willow-Bog birch/Tufted hairgrass

(*Salix glauca*-*Betula glandulosa*/*Deschampsia cespitosa*)

**n=14** This community type is found in association with the Tufted hairgrass-Sedge c.t.. Willow encroachment into a tufted hairgrass meadow eventually results in this community type. Historically fire has played an important role in the maintenance of the grassland community type in this ecoregion. Continued fire suppression will eventually allow willow and bog birch to invade many of these grassy meadows.

Willoughby (1998) found that the encroachment of willow into the Tufted hairgrass-Sedge c.t. caused a decline in forage production from 2200 kg/ha to 1800 kg/ha in the Upper Foothills subregion. This community has a high cover of willow and very little forage for domestic livestock.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
SMOOTH WILLOW . ( <i>Salix glauca</i> )	26	0-65	86
BARRET'S WILLOW ( <i>Salix barrattiana</i> )	6	0-55	43
BOG BIRCH ( <i>Betula glandulosa</i> )	10	0-25	79
<b>FORBS</b>			
YARROW ( <i>Achillea millefolium</i> )	1	0-3	43
WILD STRAWBERRY ( <i>Fragaria virginiana</i> )	2	0-20	36
LINDLEY'S ASTER ( <i>Aster ciliolatus</i> )	2	0-20	36
MOUNTAIN HELIOTROPE ( <i>Valeriana sitchensis</i> )	3	0-35	43
TALL LARKSPUR ( <i>Delphinium glaucum</i> )	1	0-5	43
WANDERING DAISY ( <i>Erigeron peregrinus</i> )	3	0-15	29
<b>GRASSES</b>			
TUFTED HAIRGRASS ( <i>Deschampsia cespitosa</i> )	19	2-35	100
HAIRY WILD RYE ( <i>Elymus innovatus</i> )	2	0-5	36
SEDGE ( <i>Carex spp.</i> )	10	0-26	86

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME :  
MESIC TO SUBHYGRIC

NUTRIENT REGIME :  
PERMESOTROPHIC

ELEVATION:  
1828(1220-2210) M

SOIL DRAINAGE :  
MOD. WELL

### FORAGE PRODUCTION KG/HA

GRASS 950  
FORBS 493  
SHRUBS 265  
TOTAL 1803 \*ESTIMATE

SUGGESTED GRAZING CAPACITY  
0.5 HA/AUM OR 1.1 ACRES/AUM

**SACFB5. Willow-Bog birch/Clover-Dandelion**  
*(Salix glauca-Betula glandulosa/Trifolium repens-Taraxacum officinale)*

**n=1** This community type represents a Willow-Bog birch/Tufted hairgrass community that has been extensively grazed by horses. This community was described near a historic campsite in the South Ram river drainage. Long-term moderate grazing pressure or heavy grazing pressure over a couple of years causes tufted hairgrass to decline and allows sedge, slender wheatgrass, Kentucky bluegrass, clover and dandelion to increase (Willoughby 2001) These community types are highly productive for domestic livestock throughout the growing season, but the poor quality of Kentucky bluegrass, clover and dandelion, particularly, in the dormant season limits the use of these community types for wildlife.

**PLANT COMPOSITION** **CANOPY COVER(%)**  
**MEAN RANGE CONST.**

<b>SHRUBS</b>			
SMOOTH WILLOW . <i>(Salix glauca)</i>	19	-	100
<b>BOG BIRCH</b>			
<i>(Betula glandulosa)</i>	10	-	100
<b>FORBS</b>			
<b>YARROW</b>			
<i>(Achillea millefolium)</i>	11	-	100
<b>WILD STRAWBERRY</b>			
<i>(Fragaria virginiana)</i>	4	-	100
<b>DANDELION</b>			
<i>(Taraxacum officinale)</i>	15	-	100
<b>CLOVER</b>			
<i>(Trifolium repens)</i>	16	-	100
<b>GRACEFUL CINQUEFOIL</b>			
<i>(Potentilla gracilis)</i>	18	-	100
<b>GRASSES</b>			
<b>TUFTED HAIRGRASS</b>			
<i>(Deschampsia cespitosa)</i>	10	-	100
<b>SLENDER WHEATGRASS</b>			
<i>(Agropyron trachycaulum)</i>	6	-	100
<b>SEDGE</b>			
<i>(Carex spp.)</i>	12	-	100
<b>KENTUCKY BLUEGRASS</b>			
<i>(Poa pratensis)</i>	7	-	100
<b>HAIRY WILD RYE</b>			
<i>(Elymus innovatus)</i>	8	-	100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
MESIC

NUTRIENT REGIME :  
MESOTROPHIC

ELEVATION:  
1966 M

SOIL DRAINAGE :  
MOD. WELL

SLOPE:  
2%

ASPECT:  
SOUTHEAST

**FORAGE PRODUCTION KG/HA**

GRASS 1728  
 FORBS 199  
 TOTAL 1927

**SUGGESTED GRAZING CAPACITY**  
 0.5 HA/AUM OR 1.1 ACRES/AUM

## SACFB6. Willow-Bog birch/California oatgrass

(*Salix glauca*-*Betula glandulosa*/*Danthonia californica*)

**n=19** This community type likely develops from willow encroaching onto an oatgrass dominated meadow. The oatgrass meadows are found on dry, gravelly soils. These meadows may also form in frost pockets. The spread of willow is likely caused by lack of natural disturbance, such as fire. The cover of willow on this community type is fairly extensive. This will restrict access to domestic livestock. This community type would be rated as secondary range.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
SMOOTH WILLOW ( <i>Salix glauca</i> )	22	0-75	90
BARRET'S WILLOW ( <i>Salix barratiana</i> )	8	0-50	47
BOG BIRCH ( <i>Betula glandulosa</i> )	7	0-50	58
<b>FORBS</b>			
YARROW ( <i>Achillea millefolium</i> )	2	0-9	68
MOUNTAIN CINQUEFOIL ( <i>Potentilla diversifolia</i> )	1	0-6	63
WILD STRAWBERRY ( <i>Fragaria virginiana</i> )	9	0-33	90
WANDERING DAISY ( <i>Erigeron peregrinus</i> )	1	0-6	42
GLOBEFLOWER ( <i>Trollius albiflorus</i> )	2	0-25	26
NORTHERN VALERAIN ( <i>Valeriana dioica</i> )	1	0-8	21
<b>GRASSES</b>			
CALIFORNIA OATGRASS ( <i>Danthonia californica</i> )	24	10-70	100
SEDGE ( <i>Carex spp.</i> )	4	0-21	68
MOUNTAIN TIMOTHY ( <i>Phleum commutatum</i> )	4	0-35	68
HAIRY WILD RYE ( <i>Elymus innovatus</i> )	1	0-15	32
SPIKED TRISTEUM ( <i>Trisetum spicatum</i> )	3	0-20	68

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME :

MESIC-SUBHYGRIC

NUTRIENT REGIME:

MESOTROPHIC

ELEVATION:

1888(1360-2340) M

SOIL DRAINAGE:

MODERATELY WELL TO WELL

### FORAGE PRODUCTION KG/HA

GRASS	598
FORBS	418
SHRUBS	300
TOTAL	1316 *ESTIMATE

SUGGESTED GRAZING CAPACITY

0.7 HA/AUM OR 1.5 ACRES/AUM

## SACFB7. Willow-Bog birch/Hairy wildrye

*(Salix glauca-Bog birch/Elymus innovatus)*

n=20 This community is typical of the valley bottoms where the low temperatures prohibit the growth of trees. Corns and Achuff (1982) described a similar community in the Banff and Jasper National Parks. They found this community type occupied coarse stream deposits which had repeated flooding.

Bork (1994) felt this community type developed from the invasion of willow and bog birch onto grasslands in the absence of disturbance in Willmore Wilderness park. Willow cover has increased, shading the growth of grasses and allowing tall-growing forbs, such as fireweed, aster and veiny meadow rue to increase. He felt continued protection from disturbance will allow succession to shrub and eventually tree species, which will increase shading of the understory vegetation and eventually lower forage production.

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

#### TREES

WHITE SPRUCE <i>(Picea glauca)</i>	T	0-3	5
---------------------------------------	---	-----	---

#### SHRUBS

WILLOW SPP. <i>(Salix glauca)</i>	30	8-52	100
BOG BIRCH <i>(Betula glandulosa)</i>	24	2-58	90

#### FORBS

NORTHERN VALERIAN <i>(Valeriana dioica)</i>	4	0-11	75
YARROW <i>(Achillea millefolium)</i>	3	0-8	95
FIREWEED <i>(Epilobium angustifolium)</i> 3	3	0-11	65
STRAWBERRY <i>(Fragaria virginiana)</i>	4	0-12	80
TALL LARKSPUR <i>(Delphinium glaucum)</i>	3	0-10	85

#### GRASSES

BOG SEDGE <i>(Kobriesia myosuroides)</i>	5	0-16	55
HAIRY WILDRYE <i>(Elymus innovatus)</i>	10	0-30	95
GRACEFUL SEDGE <i>(Carex praeegracilis)</i>	3	0-13	55
SLENDER WHEATGRASS <i>(Agropyron trachycaulum)</i> 3	3	0-13	85

### ENVIRONMENTAL VARIABLES

#### MOISTURE REGIME:

XERIC TO SUBHYGRIC

#### NUTRIENT REGIME:

MESOTROPHIC

#### ELEVATION:

1926(1560-2250) M

#### SOIL DRAINAGE:

RAPIDLY TO MODERATELY WELL

#### SLOPE:

3(1-10)%

#### ASPECT:

EASTERLY

### FORAGE PRODUCTION KG/HA

GRASS	868(756-1003)
FORBS	713(85-2120)
SHRUB	135(0-540)
TOTAL	1716(1088-2898)

#### SUGGESTED GRAZING CAPACITY

0.5 HA/AUM 1.2 AC/AUM

**SACFB8. Bog birch/Bog sedge-Sedge**  
(*Betula glandulosa*/*Kobresia myosuroides*-*Carex spp.*)

**n=5** This community type was described on moist lowland sites at higher elevations in the Central and Northern foothills ecodistricts. It appears this community type originated from recent shrub encroachment onto sedge-bog sedge-tufted hairgrass community type. The presence of bog sedge may represent the transition between the foothills ecodistricts and the rocky mountain ecodistricts. Corns and Achuff (1982) described bog sedge dominated community types in the Central and Northern Rocky Mountains of the Subalpine subregion of Banff and Jasper National Parks. Camping and grazing of these communities by horses should be restricted.

**PLANT COMPOSITION CANOPY COVER(%)**  
MEAN RANGE CONST.

<b>SHRUBS</b>			
WILLOW SPP. ( <i>Salix spp.</i> )	3	0-10	60
BOG BIRCH ( <i>Betula glandulosa</i> )	27	12-40	100
<b>FORBS</b>			
SLENDER BLUE BEARDTONGUE ( <i>Penstemon procerus</i> )	3	1-7	100
BEARBERRY ( <i>Arctostaphylos uva-ursi</i> )	4	0-16	60
SMOOTH LEAVED CINQUEFOIL ( <i>Potentilla diversitolia</i> )	5	0-19	80
OLD MAN'S WHISKERS ( <i>Geum triflorum</i> )	3	0-9	80
ALPINE GOLDENROD ( <i>Solidago multiradiata</i> )	2	0-5	60
<b>GRASSES</b>			
BOG SEDGE ( <i>Kobresia myosuroides</i> )	26	16-45	100
GRACEFUL SEDGE ( <i>Carex praegracilis</i> )	16	0-41	80
SLENDER WHEATGRASS ( <i>Agropyron trachycaulum</i> )	9	1-23	100
CALIFORNIA OATGRASS ( <i>Danthonia californica</i> )	5	0-12	80

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
MESIC-SUBHYGRIC

NUTRIENT REGIME:  
MESOTROPHIC

ELEVATION:  
1791(1530-2286) M

SOIL DRAINAGE:  
MODERATELY WELL

SLOPE: 2(1-2)%

ASPECT: EASTERLY

**FORAGE PRODUCTION KG/HA**

GRASS	1333(391-2848)
FORBS	390(88-695)
SHRUBS	202(0-807)
TOTAL	1925(683-3416)

SUGGESTED GRAZING CAPACITY  
0.5 HA/AUM OR 1.1 ACRES/AUM

## SACFB9. Bog birch-Willow/Rough fescue

*(Betula glandulosa-Salix spp./Festuca scabrella)*

**n=4** This community type is very similar to the Bog birch/Rough fescue-Sedge community described by Willoughby (1992) in the Upper Foothills subregion. Willoughby found that the rough fescue grasslands were located upslope of tufted hairgrass meadows on slightly drier, gravelly soils. Bork (1994), also described rough fescue dominated grasslands in Willmore Wilderness Park. This community type is also similar to the Bog birch/Rough fescue-Bog sedge community type but lacks the cover of bog sedge. Bog sedge tends to grow at higher elevations and appears to indicate the transition from the Upper Foothills subregion to the Subalpine subregion.

It appears the lack of fire on this community type has allowed the shrub cover to expand, reducing forage productivity for wildlife and domestic livestock. In one study, burning a Bog birch/Rough fescue community type twice in 3 year intervals controlled birch growth and increased total forage production by over 40% compared to the unburned control in the Upper Foothills subregion (Bork 1990).

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
BOG BIRCH <i>(Betula glandulosa)</i>	39	24-62	100
WILLOW SPP. <i>(Salix spp.)</i>	13	0-26	75
<b>FORBS</b>			
AMERICAN VETCH <i>(Vicia americana)</i>	4	1-8	100
SMOOTH ASTER <i>(Aster laevis)</i>	2	0-5	100
TALL LARKSPUR <i>(Delphinium glaucum)</i>	2	0-6	75
OLD MAN'S WHISKERS <i>(Geum triflorum)</i>	4	0-7	75
FIREWEED <i>(Epilobium angustifolium)</i> 3		0-6	75
<b>GRASSES</b>			
ROUGH FESCUE <i>(Festuca scabrella)</i>	19	10-31	100
GRACEFUL SEDGE <i>(Carex praegracilis)</i>	6	1-12	100
SLENDER WHEATGRASS <i>(Agropyron trachycaulum)</i> 9		0-33	60
HAIRY WILD RYE <i>(Elymus innovatus)</i>	3	0-6	50

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:  
MESIC

NUTRIENT REGIME:  
MESOTROPHIC

ELEVATION:  
1675(1600-1750) M

SOIL DRAINAGE:  
WELL

SLOPE:  
5%

ASPECT:  
VARIABLE

### FORAGE PRODUCTION KG/HA

GRASS	1807(642-3564)
FORBS	705(492-902)
SHRUBS	559(170-800)
TOTAL	3071(2070-4226)

SUGGESTED GRAZING CAPACITY  
0.4 HA/AUM OR 0.8 ACRES/AUM



**SACFB10. Bog birch/Rough fescue-Bog sedge**  
*(Betula glandulosa/Festuca scabrella-Kobresia myosuroides)*

n=1 This community type is very similar to the Bog birch-Willow/Rough fescue community previously described. Willoughby(2001) found that the rough fescue grasslands were located upslope of tufted hairgrass meadows on slightly drier, gravelly soils. Bork (1994), also described rough fescue dominated grasslands in Willmore Wilderness Park. The presence of bog sedge in this community type appears to indicate the transition from the Upper Foothills and lower Subalpine subregions to the Upper subalpine subregion.

It appears the lack of fire on this community type has allowed the shrub cover to expand, reducing forage productivity for wildlife and domestic livestock. In one study, burning a Bog birch/Rough fescue community type twice in 3 year intervals controlled birch growth and increased total forage production by over 40% compared to the unburned control in the Upper Foothills subregion (Bork 1990).

**PLANT COMPOSITION CANOPY COVER(%)**

MEAN RANGE CONST.

**SHRUBS**

BOG BIRCH <i>(Betula glandulosa)</i>	30	-	100
WILLOW SPP. <i>(Salix barclayi)</i>	5	-	100

**FORBS**

FALSE DANDELION <i>(Agoseris glauca)</i>	6	-	100
GRACEFUL CINQUEFOIL <i>(Potentilla gracilis)</i>	6	-	100
TALL LARKSPUR <i>(Delphinium glaucum)</i>	6	-	100
ALPINE GOLDENROD <i>(Solidago multiradiata)</i>	5	-	100
SHOW LOCOWEED <i>(Oxytropis splendens)</i>	5	-	100

**GRASSES**

ROUGH FESCUE <i>(Festuca scabrella)</i>	85	-	100
GRACEFUL SEDGE <i>(Carex praegracilis)</i>	10	-	100
SLENDER WHEATGRASS <i>(Agropyron trachycaulum)</i> 34		-	100
BOG SEDGE <i>(Kobresia myosuroides)</i>	13	-	100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:

MESIC

NUTRIENT REGIME:

MESOTROPHIC

ELEVATION:

1981 M

SOIL DRAINAGE:

WELL

**FORAGE PRODUCTION KG/HA**

GRASS	1201
FORBS	147
SHRUBS	0
TOTAL	1348

SUGGESTED GRAZING CAPACITY  
 0.7 HA/AUM OR 1.6 ACRES/AUM

**SACFB11. Willow/Fringed brome-Sedge**  
*(Salix barclayi/Bromus ciliatus-Carex spp.)*

**n=1** This community was described on the banks of Forbidden Creek west of Rocky Mtn. House where the water table is high but flooding is rare. It occupies the fluvial terraces along the creek. A similar community type Willow/Fringed brome-Slender wheatgrass was described in the Lower Foothills subregion (Lane et al. 2000). The production of the Lower Foothills type averaged over 1700 kg/ha. This community type had only half the production (760 kg/ha). The more extreme climatic conditions of this site in the subalpine likely limits the growth of forage.

**PLANT COMPOSITION CANOPY COVER(%)**

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
WILLOW SPP. <i>(Salix barclayi)</i>	50	-	100
<b>FORBS</b>			
YARROW <i>(Achillea millefolium)</i>	8	-	100
GRACEFUL CINQUEFOIL <i>(Potentilla gracilis)</i>	9	-	100
ALPINE BISTORT <i>(Polygonum viviparum)</i>	43	-	100
OLD MAN'S WHISKERS <i>(Geum triflorum)</i>	8	-	100
<b>GRASSES</b>			
FRINGED BROME <i>(Bromus ciliatus)</i>	61	-	100
GRACEFUL SEDGE <i>(Carex praegracilis)</i>	62	-	100
ROUGH FESCUE <i>(Festuca scabrella)</i>	2	-	100
SLENDER WHEATGRASS <i>(Agropyron trachycaulum)</i>	15	-	100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
MESIC

NUTRIENT REGIME:  
MESOTROPHIC

ELEVATION:  
2286 M

SOIL DRAINAGE:  
IMPEPECTLY

**FORAGE PRDUCTION KG/HA**

GRASS	472
FORBS	288
TOTAL	760

SUGGESTED GRAZING CAPACITY  
1.1 HA/AUM OR 2.4 ACRES/AUM

**SUBALPINE SUBREGION**  
**CENTRAL AND NORTHERN ROCKY MOUNTAINS**  
**NATIVE GRASSLANDS AND SHRUBLANDS**



**Figure 3.** The presence of bog sedge in this Willow-Bog birch/Bog sedge community indicates the transition to the higher and steeper Central and Northern Rocky Mountains of the Subalpine subregion. Note the transition from the Subalpine to Alpine subregions in the background.

## Native grass and shrubland ecology Mountain ecodistricts

The native grass and shrubland community types in the Central and Northern Rocky Mountains of the Subalpine subregion (Table 3) are found in the valley bottoms adjacent to streams and rivers. The tufted hairgrass, California oatgrass and rough fescue dominated community types described previously in the foothills can also be found in the mountains, but these grassland community types are more common in the foothills and therefore were described in that section of the guide.

There are a number of grassland community types in the Central and Northern Rocky Mountains which are unique and appear to represent a transition from the lower subalpine mountain ecodistricts to the Alpine subregion. These include the hairy wildrye, junegrass and shrubby cinquefoil dominated community types (SACMA3,4,5,6) which are found on steep south facing slopes, at higher elevations throughout the mountains (Figure 2). Near timberline there is a unique forb dominated (globeflower, wandering daisy, mountain marigold, mountain heliotrope) community type found on imperfectly to well drained sites. The presence of bog sedge (*Kobresia myosuroides*) in the Bog sedge-California oatgrass community type appears to indicate the transition from the lower Central and Northern Rocky Mountains to the Alpine subregion. Ogilvie (1969) and Corns and Achuff (1982), described bog sedge dominated community types in the higher elevations of the subalpine and alpine of the Rocky Mountains of Banff and Jasper National Parks.

The maintenance of the grassland community types in the mountains is extremely fire dependent. The lack of fire quickly allows bog birch and willow to expand shading the modal grassland community types. Prolonged shading causes the understory composition to shift from a tufted hairgrass-California oatgrass dominated understory to a slender wheatgrass-hairy wildrye dominated understory. Under a heavy shrub cover there is little forb or grass cover. The sequence of the grassland and shrubland community types unique to the mountain ecodistricts is outlined in figures 2 and 4. These figures represent the transition from willow, bog birch dominated communities in the valley bottoms to the grass and dwarf shrublands in the upper Subalpine and Alpine subregions.


Many of these subalpine grass and shrublands are very fragile because of exposure and cold climate. The forage productivity is generally only half of what is found in the lower Upper Foothills subregion and recovery from overgrazing will likely take some time because of the poor growing conditions. As a result grazing by domestic livestock should be done with caution.

**Table 3.** Native grass and shrublands of the Central and Northern Rocky Mountain eodistricts of the Subalpine subregion

Community number	Community type	Grass	Forb	Shrub	Productivity (kg/ha)	Moisture	Drainage	Carrying capacity (ha/AUM)
<b>A. GRASSLANDS</b>								
SACMA1.	Bog sedge-California oatgrass					Mesic	Well	Non-use
SACMA2.	Forb meadows				N/A	Mesic to Hygric	Poorly	Non-use
SACMA3.	Shrubby cinquefoil/Hairy wildrye				N/A	Xeric	Rapidly	Non-use
SACMA4.	Bearberry-Juniper				N/A	Xeric	Rapidly	Non-use
SACMA5.	Junegrass-Hairy wildrye-Brome				N/A	Subxeric	Rapidly	Non-use
SACMA6.	Hairy wildrye/Bearberry-Juniper				N/A	Mesic	Well	Non-use
SACMA7.	Northern wheatgrass				N/A	Subxeric	Rapidly	Non-use
SACMA8.	Alpine bluegrass				N/A	Submesic	Well	Non-use
SACMA9.	Yellow mountain avens				N/A	Subxeric	Rapidly	Non-use
<b>B. SHRUBLANDS</b>								
SACMB1.	Willow-Bog birch/Water sedge	2320	24		2344	Hygric	Poorly	Non-use
SACMB2.	Willow-Bog birch/Sedge				N/A	Subhygric	Poorly	Non-use
SACMB3.	Willow-Bog birch/Hairy wildrye				N/A	Subhygric	Mod. well	Non-use
SACMB4.	Willow-Bog birch/Bog sedge				N/A	Subhygric	Mod. well	Non-use
SACMB5.	Bog birch-Juniper				N/A	Subxeric	Rapidly	Non-use
SACMB6.	Willow/Forb				N/A	Mesic	Imperfectly	Non-use

Table 3. cont'd

Community number	Community type	Production(kg/ha)			Moisture	Drainage	Carrying capacity (ac/AUM)
		Grass	Forb	Shrub			
SACMB7.	Grouseberry/Juniper			N/A	Mesic	Well	Non-use
SACMB8.	Subalpine fir			N/A	Mesic	Well	Non-use



Adapted from (Ogilvie 1969)

Figure 4. Sequence of plant communities in the Alpine subregion of Alberta.

## Native grass and shrublands in Central and Northern Mountain and Foothills areas

1. Community above timberline (Alpine subregion).....	<b>Alpine section of guide</b>	
Community not above timberline, meadows, shrublands or south facing slopes dominated by shrubs and grass.....		2
2. Moist sites, community dominated by shrubs >20% cover (willow, bog birch) or timberline communities with dwarf trees or grouseberry .....	<b>see shrub key pg 34</b>	
Drier to moist sites shrub cover <10% site dominated by grasses and forbs.....		3
3. Community very wet free standing water, dominated by sedge, cottongrass or tufted bulrush .....		4
Community drier, dominated by tufted hairgrass, rough fescue, california oatgrass, hairy wildrye, juniper or bearberry.....		6
4. Very wet nutrient rich sites dominated by water or beaked sedge.....	<b>SACFA1</b>	
Very wet nutrient poor, acidic sites dominated by tufted bulrush and cottongrass.....		5
5. Site dominated by tufted bulrush.....	<b>SACFA2</b>	
Site dominated by cottongrass and sedge.....	<b>SACFA3</b>	
6. Grasslands of meadows and lowland areas.....		7
Grasslands of south facing slopes, or windswept ridges, hairy wildrye, bearberry, white mtn. avens dominated .....		20
7. Higher elevation sites near or at timberline, moist sites dominated by wandering daisy, globeflower, mountain marigold.....	<b>SACMA2</b>	
Lower elevation sites, valley bottoms adjacent to streams or rivers or if higher elevation dry sites dominated by bog sedge.....		8
8. Disturbed or grazed community types dominated by Kentucky bluegrass, creeping red fescue, dandelion, alpine bluegrass, alpine timothy or fireweed.....		9
Undisturbed community types dominated by rough fescue, California oatgrass, bog sedge, tufted hairgrass, and sedge species.....		14
9. Moderately grazed site native increasers dominant (slender wheatgrass, sedge, fringed brome, rocky mtn. fescue, alpine timothy), Alpine or Kentucky bluegrass increasing in cover.....		10
Heavily grazed sites dominated by Kentucky bluegrass or abandoned airstrips dominated by creeping red fescue or native forb species (fireweed, veiny meadow rue).....		12
10. Site dominated by Alpine bluegrass.....	<b>SACMA8</b>	
Site dominated by rocky mtn. fescue, alpine timothy, sedge, slender wheatgrass, fringed brome (moister sites).....		11
11. Site dominated by rocky mtn. fescue, alpine timothy, sedge.....	<b>SACFA6</b>	
Site dominated by slender wheatgrass, sedge, fringed brome.....	<b>SACFA7</b>	
12. Heavily grazed site dominated by Kentucky bluegrass and dandelion.....	<b>SACFA16</b>	
Abandoned airstrips dominated by creeping red fescue, or invaded strips dominated by forbs (fireweed, veiny meadow).....		13
13. Site dominated by creeping red fescue.....	<b>SACFA15</b>	
Moister disturbed site dominated by fireweed and veiny meadow rue... <b>SACFA17</b>		
14. Moist sites dominated by sedge and tufted hairgrass.....		15
Drier sites dominated by rough fescue, hairy wildrye, bog sedge, yellow dryas or California oatgrass.....		16
15. Site dominated by tufted hairgrass, sedge co-dominant.....	<b>SACFA4</b>	
Site dominated by sedge, tufted hairgrass co-dominant.....	<b>SACFA5</b>	
16. Sites dominated by rough fescue and hairy wildrye.....	<b>SACFA9</b>	
Sites dominated by California oatgrass, bog sedge, or yellow dryas.....		17
17. Gravelly river flats dominated by yellow dryas.....	<b>SACMA9</b>	
Meadow areas dominated by California oatgrass and/or bog sedge.....		18
18. Site dominated by bog sedge and california oatgrass.....		19
Site dominated by california oatgrass and sedge, bog sedge not present.....	<b>SACFA8</b>	
19. Site dominated by bog sedge, california oatgrass, drier sites.....	<b>SACMA1</b>	
Site co-dominated by bog sedge, tufted hairgrass, and sedge moister sites... <b>SACFA13</b>		
20. Lower elevation grasslands in the Foothills of the Subalpine.....		21
Higher elevation grasslands in the mountains of the Subalpine.....		22
21. Northern wheatgrass dominated site.....	<b>SACMA7</b>	



Fringed sage, sedge and junegrass dominated slope.....	SACFA12	
22. Avalanche slopes dominated by hairy wildrye, juniper, and bearberry....	SACMA6	
Drier sites or windswept ridges dominated hairy wildrye, juniper, bearberry, shrubby cinquefoil, white mtn. avens.....		23
23. Windswept ridges dominated by white mtn. avens.....	SACFA14	
South facing slopes dominated by hairy wildrye.....		24
24. Shallow rocky soils with little grass cover, site dominated by bearberry..	SACMA4	
Deeper soils, good grass cover dominated by hairy wildrye, junegrass.....		25
25. Shrubby cinquefoil dominant in stand.....	SACMA3	
Grass cover extensive, dominated by hairy wildrye, junegrass, and brome....	SACMA5	

## Shrub dominated communities

1. Timberline communities dominated by whitebark pine, subalpine fir, grouseberry, or willow communities with marsh marigold, wandering daisy or globeflower in understory.....		2
Riparian communities adjacent to streams or rivers.....		5
2. Trees present in community (whitebark pine, subalpine fir) or grouseberry dominated....		3
Moist seepage areas at treeline dominated by globeflower, wandering daisy or marsh marigold in understory.....	SACMB6	
3. Trees (subalpine fir, whitebark pine) on site.....		4
Grouseberry dominated shrubland.....	SACMB7	
4. Whitebark pine present.....	SACMB3	
Subalpine fir present.....	SACMB8	
5. Very wet sites with water sedge or horsetail dominated understories.....		6
Drier sites with tufted hairgrass, california oatgrass, bog sedge, hairy wildrye, rough fescue Kentucky bluegrass, dandelion dominated understories.....		7
6. Water sedge dominated understory.....	SACFB1, SACMB1	
Horsetail dominated understory.....	SACFB2	
7. Grazed communities dominated by clover and dandelion in understory.....	SACFB5	
Ungrazed sites dominated by native forbs and grasses in understory.....		8
8. Shrubland communities on seepage areas on south facing slopes with shallow soils, dominated by bog birch and juniper.....	SACMB5	
Meadows and lowland shrublands dominated by rough fescue, bog sedge, california oatgrass, tufted hairgrass, hairy wildrye or sedge in the understory.....		9
9. Rough fescue dominates the understory.....		10
Tufted hairgrass, california oatgrass, bog sedge, sedge, hairy wildrye dominate.....		11
10. Rough fescue and bog sedge dominate understory higher elevations.....	SACFB10	
Rough fescue dominates, bog sedge not present lower elevations.....	SACFB9	
11. Moister sites with deep fluvial deposits dominated by tufted hairgrass, sedge, or fringed brome in understory.....		12
Drier sites which are well drained at the surface dominated by hairy wildrye, sedge, bog sedge or california oatgrass in understory.....		14
12. Tufted hairgrass or sedge dominated understory.....		13
Fluvial areas with Fringed brome dominated understory, lower elevation.....	SACFB11	
13. Tufted hairgrass dominates understory.....	SACFB4	
Graceful sedge and other sedge species dominate understory.....	SACFB3, SASMB2	
14. Modal sites with hairy wildrye and sedge dominating understory.....		15
Sites dominated by california oatgrass or bog sedge in understory.....		16
15. Hairy wildrye dominates understory.....	SACFB7, SACMB3	
Graceful sedge and other sedge species dominate understory.....	SACFB3, SASMB2	
16. California oatgrass dominates understory.....	SACFB6	
Bog sedge dominates understory.....	SACFB8, SACMB4	

## SACMA1. Bog sedge-California oatgrass

*(Kobresia myosuroides-Danthonia californica)*

n=1 This community type appears to represent the transition from the foothills ecodistricts to the mountain ecodistricts of the subalpine. It appears that tufted hairgrass, california oatgrass and rough fescue all decline and bog sedge increases as there is an increase in elevation and change from the foothills to the mountains.

This community type is found on level to gently sloping valley bottoms with mesic moisture regimes. The presence of California oatgrass maybe indicative of a well drained, gravelly site. In the Yukon the California oatgrass dominated community types were found to form in depressions which appeared to act as pronounced frost pockets (Bailey et al. 1992). Bog sedge also appears to be adapted to these site condition (Oglivie 1969).

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
WILLOW <i>(Salix spp.)</i>	2	-	100
SHRUBBY CINQUEFOIL <i>(Potentilla fruticosa)</i>	25	-	100
<b>FORBS</b>			
BEARBERRY <i>(Arctostaphylos uva-ursi)</i>	15	-	100
YARROW <i>(Achillea millefolium)</i>	1	-	100
SMALL LEAVED EVERLASTING <i>(Antennaria parviflora)</i>	1	-	100
<b>GRASSES</b>			
CALIFORNIA OATGRASS <i>(Danthonia californica)</i>	35	-	100
BOG SEDGE <i>(Kobresia myosuroides)</i>	25	-	100

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:

MESIC

NUTRIENT REGIME:

MESOTROPHIC

ELEVATION:

1850 M

SOIL DRAINAGE:

WELL

### FORAGE PRODUCTION KG/HA

SUGGESTED GRAZING CAPACITY NON-USE
---------------------------------------

## SACMA2. Forb meadows

(*Trollius albiflorus*, *Erigeron peregrinus*, *Anemone occidentalis*, *Caltha leptosepala*)

n=17 These forb dominated meadows include both the *Caltha leptosepala*-*Trollius albiflorus* and *Erigeron peregrinus*-*Valeriana sitchensis* community types described by Corns and Achuff (1982). These meadows occupy mesic to hygric, gently sloping, upper subalpine to alpine areas. The soils are imperfectly to well drained Gleysols on fluvial and morainal landforms. On the poorly to imperfectly drained sites in areas where snow melts late and seepage is received throughout the growing season mountain marigold and globeflower predominate. In contrast on better drained, drier sites fleabane and mountain heliotrope predominate.

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

#### SHRUBS

ARCTIC WILLOW  
(*Salix arctica*) 1 0-5 59

WESTERN MOUNTAIN HEATHER  
(*Cassiope mertensiana*) 3 0-10 47

#### FORBS

WANDERING DAISY  
(*Erigeron peregrinus*) 9 0-10 88

WOOLY EVERLASTING  
(*Antennaria lanata*) 4 0-35 65

GLOBEFLOWER  
(*Trollius albiflorus*) 13 0-40 82

MOUNTAIN HELIOTROPE  
(*Valeriana sitchensis*) 9 0-20 82

MOUNTAIN MARIGOLD  
(*Caltha leptosepala*) 5 0-23 53

CHALICEFLOWER  
(*Anemone occidentalis*) 7 0-20 70

#### GRASSES

BLACKENING SEDGE  
(*Carex nigrescens*) 2 0-20 35

MOUNTAIN TIMOTHY  
(*Phleum commutatum*) 1 0-2 41

SEDGE  
(*Carex spectabilis*) 1 0-10 23

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:  
MESIC TO HYGRIC

NUTRIENT REGIME:  
PERMESOTROPHIC

ELEVATION:  
2024 (1850-2300) M

SOIL DRAINAGE:  
IMPERFECTLY TO WELL

SLOPE:  
27(3-55)%

ASPECT:  
VARIABLE

### FORAGE PRODUCTION KG/HA

SUGGESTED GRAZING CAPACITY  
NON-USE

## SACMA3. Shrubby cinquefoil/Hairy wildrye

(*Potentilla fruticosa*/*Elymus innovatus*)

n=5    Corns and Achuff (1982), described this community type on subxeric to xeric, south facing slopes in Banff and Jasper. The soils are rapidly to well drained Regosols on colluvial, eolian and glacial landforms. On more stable sites they felt succession would be to a Lodgepole pine/Juniper/Bearberry community type.

### PLANT COMPOSITION CANOPY COVER(%)

MEAN    RANGE    CONST.

#### SHRUBS

SHRUBBY CINQUEFOIL. ( <i>Potentilla fruticosa</i> )	20	4-40	100
BOG BIRCH ( <i>Betula glandulosa</i> )	2	0-5	40

#### FORBS

BEARBERRY ( <i>Arctostaphylos uva-ursi</i> )	1	0-3	20
WHITE MOUNTAIN AVENS ( <i>Dryas integrifolia</i> )	3	0-15	20
STRAWBERRY ( <i>Fragaria virginiana</i> )	6	0-10	60
ALPINE FORGET-ME-KNOT ( <i>Myosotis alpestris</i> )	5	0-15	40
SWEET-FLOWERED ANDROSACE ( <i>Androsace chamaejasme</i> ) <sup>4</sup>	4	0-15	40
WANDERING DAISY ( <i>Erigeron peregrinus</i> )	2	0-11	20
SPOTTED SAXIFRAGE ( <i>Saxifraga bronchialis</i> )	3	0-13	20

#### GRASSES

HAIRY WILDRYE ( <i>Elymus innovatus</i> )	3	0-10	60
BLUNT SEDGE ( <i>Carex obtusata</i> )	1	0-4	20
NORWAY SEDGE ( <i>Carex norvegica</i> )	8	0-40	20
BROAD GLUMED WHEATGRASS ( <i>Agropyron violaceum</i> )	5	0-25	20

### ENVIRONMENTAL VARIABLES

#### MOISTURE REGIME:

XERIC-SUBXERIC

#### NUTRIENT REGIME:

SUBMESOTROPHIC

#### ELEVATION:

1720(1500-1800) M

#### SOIL DRAINAGE:

RAPIDLY

#### SLOPE:

28 (15-55)%

#### ASPECT:

SOUTHERLY

### FORAGE PRODUCTION KG/HA

NOT AVAILABLE

SUGGESTED GRAZING CAPACITY

NON-USE

## SACMA4. Bearberry-Juniper

(*Arctostaphylos uva-ursi*-*Juniperus communis*)

**n=13** This community type is very similar to the previously describe Shrubby cinquefoil/Hairy wildrye dominated community type. Both community types occupy rapidly drained, steep south facing slopes. This community type is distinguished from the Shrubby cinquefoil community type by the presence of a high cover of bearberry and juniper and a low cover of shrubby cinquefoil. This community type is much drier than the shrubby cinquefoil type and is located on much steeper slopes.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
SHRUBBY CINQUEFOIL ( <i>Potentilla fruticosa</i> )	4	0-18	77
GROUND JUNIPER ( <i>Juniperus communis</i> )	9	0-25	92
SMOOTH WILLOW ( <i>Salix glauca</i> )	2	0-10	46
<b>FORBS</b>			
STRAWBERRY ( <i>Fragaria virginiana</i> )	1	0-3	62
WHITE CAMUS ( <i>Zigadenus elegans</i> )	2	0-8	46
MOUNTAIN VALERIAN ( <i>Valeriana sitchensis</i> )	1	0-7	15
HARE BELL ( <i>Campanula rotundifolia</i> )	1	0-5	39
TWINFLOWER ( <i>Linnaea borealis</i> )	1	0-5	15
BEARBERRY ( <i>Arctostaphylos uva-ursi</i> )	16	8-25	100
<b>GRASSES</b>			
HAIRY WILDRYE ( <i>Elymus innovatus</i> )	2	0-10	46
SPIKED TRisetum ( <i>Trisetum spicatum</i> )	T	0-1	23
SLENDER WHEATGRASS ( <i>Agropyron trachycaulum</i> )	T	0	8
SHEEP FESCUE ( <i>Festuca saximontana</i> )	T	0-1	15
SEDGE SPP. ( <i>Carex spp.</i> )	1	0-6	54

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:  
XERIC-SUBXERIC

NUTRIENT REGIME:  
SUBMESOTROPHIC

ELEVATION:  
1901(1700-2000) M

SOIL DRAINAGE:  
RAPIDLY

SLOPE:  
60(55-71)%

ASPECT:  
SOUTHERLY

### FORAGE PRODUCTION KG/HA

SUGGESTED GRAZING CAPACITY NON-USE
---------------------------------------

## SACMA5. Junegrass-Hairy wildrye-Brome

*(Koeleria macrantha-Elymus innovatus-Bromus inermis)*

**n=22** This community type was described by Corns and Achuff (1982) on subxeric, steep south facing slopes in the Front ranges east of Banff and Jasper. It is very similar to the bearberry and shrubby cinquefoil community types previously described, but this community type has better developed soils (Brunisols) than the Regosolic soils of the other community types.

This community type is also similar to the Pasture sagewort/Junegrass and Junegrass-Plains reed grass community types described by Willoughby et al (2003) and Stringer (1973) in the lower Montane subregion and the Junegrass/Sage community type described in the Upper Foothills subregion (Willoughby 2001). The high elevations of this community distinguishes this community type from the lower elevation grasslands (Corns and Achuff 1982).

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

#### SHRUBS

SHRUBBY CINQUEFOIL  
*(Potentilla fruticosa)* 1 0-6 59

#### FORBS

SLENDER BLUE BEARDTONGUE  
*(Penstemon procerus)* T 0-2 27

YARROW  
*(Achillea millefolium)* 2 0-15 82

SHOWY LOCOWEED  
*(Oxytropis splendens)* 2 0-15 59

BEARBERRY  
*(Arctostaphylos uva-ursi)* 1 0-10 41

YELLOW HEDYSARUM  
*(Hedysarum sulphurcens)* 5 0-15 50

SMALL LEAVED EVERLASTING  
*(Antennaria parviflora)* 3 0-15 36

#### GRASSES

ROCKY MOUNTAIN FESCUE  
*(Festuca saximontana)* T 0-1 18

JUNEGRASS  
*(Koeleria macrantha)* 15 0-35 96

HAIRY WILDRYE  
*(Elymus innovatus)* 17 0-55 82

SEDGE  
*(Carex spp.)* 1 0-10 54

BROME  
*(Bromus inermis)* 10 0-30 64

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:  
XERIC-SUBXERIC

NUTRIENT REGIME:  
SUBMESOTROPHIC

ELEVATION:  
1964(1910-2100) M

SOIL DRAINAGE:  
RAPIDLY

SLOPE:  
49(40-56)%

ASPECT:  
SOUTHERLY

### FORAGE PRODUCTION KG/HA

SUGGESTED GRAZING CAPACITY  
NON-USE

**SACMA6. Hairy wildrye/Bearberry-Juniper**  
*(Elymus innovatus/Arctostaphylos uva-ursi-Juniperus communis)*

**n=36** Ogilvie (1969) and Corns and Achuff (1982), described this community type on steep south-facing slopes, with stoney, black soils. This community type is also subjected to frequent snow avalanching, particularly during the spring when the snow is melting. This community type is moister than the junegrass, bearberry and shrubby cinquefoil communities previously described. This is likely the result of the increased snow cover. The soils are Melanic Brunisols and Humic Regosols, which are better developed than the drier community types previously described.

**PLANT COMPOSITION CANOPY COVER(%)**

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
JUNIPER			
<i>(Juniperus communis)</i>	6	0-30	63
SHRUBBY CINQUEFOIL			
<i>(Potentilla fruticosa)</i>	3	0-24	44
<b>FORBS</b>			
BEARBERRY			
<i>(Arctostaphylos uva-ursi)</i>	16	0-60	76
WILD STRAWBERRY			
<i>(Fragaria virginiana)</i>	2	0-15	74
WHITE CAMAS			
<i>(Zigadenus elegans)</i>	1	0-7	39
FIREWEED			
<i>(Epilobium angustifolium)</i>	1	0-5	46
TWINFLOWER			
<i>(Linnaea borealis)</i>	1	0-35	11
YELLOW HEDYSARUM			
<i>(Hedysarum sulphureum)</i>	3	0-25	39
<b>GRASSES</b>			
HAIRY WILDRYE			
<i>(Elymus innovatus)</i>	44	0-70	98
JUNEGRASS			
<i>(Koeleria macrantha)</i>	1	0-5	28
CALIFORNIA OATGRASS			
<i>(Danthonia californica)</i>	15	0-40	13
RED FESCUE			
<i>(Festuca rubra)</i>	1	0-30	2
SPIKED TRISETUM			
<i>(Trisetum spicatum)</i>	1	0-5	28

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
MESIC

NUTRIENT REGIME:  
PERMESOTROPHIC

ELEVATION:  
1942(1380-2300)M

SOIL DRAINAGE:  
WELL

SLOPE:  
52( 0-80)%

ASPECT:  
SOUTH

**FORAGE PRODUCTION KG/HA**

SUGGESTED GRAZING CAPACITY  
NON-USE

## SACMA7. Northern wheatgrass

*(Agropyron dasystachyum)*

**n=3** This community type occurs on steep south facing slopes, with shallow soils, at lower elevations in the subalpine. It is very similar to the northern wheatgrass community type described by Willoughby et al. (2003) and Corns and Achuff (1982) in the Montane subregion of Banff and Jasper. This community type is distinguished from the other hairy wildrye, junegrass, bearberry and shrubby cinquefoil community types by the presence of northern wheatgrass and the lower elevations. The inaccessibility and fragile nature of the soils make this community type unsuitable for grazing.

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

#### SHRUBS

PRICKLY ROSE  
*(Rosa acicularis)* 1 0-1 67

#### FORBS

BEARBERRY  
*(Arctostaphylos uva-ursi)* T 0-1 33

SHOWY LOCOWEED  
*(Oxytropis splendens)* 1 0-3 67

FIREWEED  
*(Epilobium angustifolium)* 2 0-5 67

WESTERN MEADOW RUE  
*(Thalictrum occidentale)* 6 0-18 33

COMMON YARROW  
*(Achillea millefolium)* 1 1-2 100

#### GRASSES

JUNEGRASS  
*(Koeleria macrantha)* 3 0-5 67

ALPINE TIMOTHY  
*(Phleum commutatum)* 1 0-2 67

BLUNT SEDGE  
*(Carex obtusata)* 1 0-1 67

HAIRY WILDRYE  
*(Elymus innovatus)* 3 0-10 33

NORTHERN WHEATGRASS  
*(Agropyron dasystachyum)* 30 0-15 67

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:  
SUBXERIC-XERIC

NUTRIENT REGIME:  
SUBMESOTROPHIC

SLOPE:  
31 (30-60)%

ASPECT:  
SOUTHERLY

ELEVATION:  
1720( 1220-1859)M

SOIL DRAINAGE:  
RAPIDLY

### FORAGE PRODUCTION KG/HA

TOTAL 400 \*ESTIMATE

SUGGESTED GRAZING CAPACITY  
0 HA/AUM OR 0 ACRES/AUM



## SACMA8. Alpine bluegrass

(*Poa alpina*)

n=1 The ecology of this community type is unclear. It was described on a gentle, easterly slope in the lower subalpine. Alpine bluegrass is known to grow in meadows, tundra and rocky slopes and is often abundant where the ground has been compacted (MacKinnon et al. 1992). It is possible that this community type could have been described adjacent to a game or hiking trail.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
DWARF BILBERRY ( <i>Vaccinium caespitosum</i> )	6	-	100
SHRUBBY CINQUEFOIL ( <i>Potentilla fruticosa</i> )	4	-	100
<b>FORBS</b>			
STRAWBERRY ( <i>Fragaria virginiana</i> )	20	-	100
FIREWEED ( <i>Epilobium angustifolium</i> )	4	-	100
BALSAM GROUNDSEL ( <i>Senecio pauperculus</i> )	3	-	100
<b>GRASSES</b>			
ALPINE BLUEGRASS ( <i>Poa alpina</i> )	45	-	100
ROCKY MOUNTAIN FESCUE ( <i>Fescue brachyphylla</i> )	1	-	100

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:

SUBMESIC

NUTRIENT REGIME:

SUBMESOTROPHIC

ELEVATION:

1800M

SOIL DRAINAGE:

WELL

SLOPE:

4%

ASPECT:

EASTERLY

### FORAGE PRODUCTION KG/HA

SUGGESTED GRAZING CAPACITY

NON-USE

**SACMA9. Yellow mountain avens**  
(*Dryas drummondii*)

n=4 Corns and Achuff (1982), described this community type on recent fluvial and glacialfluvial landforms with gentle slopes. The soils are rapidly drained. Willoughby et al. (2003), described a yellow mountain avens community type on dry, gravelly river flats with nutrient poor soils in the Montane subregion. They found this community type to be successionaly immature and succession would be to a Balsam poplar dominated community type.

**PLANT COMPOSITION CANOPY COVER(%)**

MEAN RANGE CONST.

**SHRUBS**

YELLOW MOUNTAIN AVENS ( <i>Dryas drummondii</i> )	50	25-50	100
WILLOW SPP. ( <i>Salix spp.</i> )	5	0-20	50

**FORBS**

ALPINE GOLDENROD ( <i>Solidago multiradiata</i> )	T	0-1	25
BROAD LEAVED FIREWEED ( <i>Epilobium latifolium</i> )	1	0-1	50
SHOWY LOCOWEED ( <i>Oxytropis splendens</i> )	1	0-3	25

**GRASSES**

SEDGE ( <i>Carex spp</i> )	T	0-2	25
-------------------------------	---	-----	----

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
SUBXERIC

NUTRIENT REGIME:  
MESOTROPHIC

ELEVATION:  
1542 (1450-1670)M

SOIL DRAINAGE:  
RAPIDLY

**FORAGE PRODUCTION KG/HA**

SUGGESTED GRAZING CAPACITY  
NON-USE

## SACMB1. Willow/Water sedge

(*Salix spp./Carex aquatilis*)

**n=23** This shrub community appears on areas with very poor drainage. It is found in association with the wetter water sedge meadows. These sites are fairly productive but difficult to graze due to the moist ground conditions and heavy shrub cover which reduces access and mobility within the area. Increased flooding and prolonged waterlogging may result in the disappearance of willow and a transition to a water sedge meadow.

This community is similar to the water sedge-beaked sedge community in that it is found throughout the foothills and into the mountains. It maybe found in the Upper Foothills, Subalpine and lower Alpine subregions.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
WILLOW			
( <i>Salix spp</i> )	27	0-75	96
BOG BIRCH			
( <i>Betula glandulosa</i> )	7	0-38	70
<b>FORBS</b>			
ELEPHANT'S HEAD			
( <i>Pedicularis groenlandicum</i> )1		0-7	30
DWARF RASPBERRY			
( <i>Rubus arcticus</i> )	1	0-4	30
ALPINE BISTORT			
( <i>Polygonum viviparum</i> )	T	0-1	17
WANDERING DAISY			
( <i>Erigeron peregrinus</i> )	1	0-3	22
<b>GRASSES</b>			
WATER SEDGE			
( <i>Carex aquatilis</i> )	50	15-80	100
TUFTED HAIRGRASS			
( <i>Deschampsia cespitosa</i> )	3	0-30	26
SEDEGE			
( <i>Carex spp.</i> )	9	0-20	4
BALTIC RUSH			
( <i>Juncus balticus</i> )	T	0-1	4

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:  
SUBHYGRIC-HYGRIC

NUTRIENT REGIME:  
PERMESOTROPHIC

ELEVATION:  
1747(1340-1970) M

SOIL DRAINAGE:  
POORLY

### FORAGE PRODUCTION KG/HA

GRASS 2320  
FORBS 24  
TOTAL 2344

SUGGESTED GRAZING CAPACITY

NON-USE

**SACMB2. Willow-Bog birch/Sedge**  
*(Salix glauca-Betula glandulosa/Carex spp.)*

**n=21** This community type is found in association with the Tufted hairgrass-Sedge or California oatgrass community type. Willow encroachment into grassland meadows eventually results in this community type. Historically fire has played an important role in the maintenance of the grassland community type in this ecoregion. Continued fire suppression will eventually allow willow and bog birch to invade many of the grassy meadows.

This community type is slightly drier than the Willow/Water sedge dominated community type.

**PLANT COMPOSITION CANOPY**

**COVER(%)**

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
SMOOTH WILLOW . <i>(Salix glauca)</i>	11	0-30	67
BARRET'S WILLOW <i>(Salix barrattiana)</i>	15	0-55	71
BOG BIRCH <i>(Betula glandulosa)</i>	12	0-30	81
<b>FORBS</b>			
YARROW <i>(Achillea millefolium)</i>	1	0-5	43
WILD STRAWBERRY <i>(Fragaria virginiana)</i>	3	0-23	57
WANDERING DAISY <i>(Erigeron peregrinus)</i>	1	0-8	38
MOUNTAIN HELIOTROPE <i>(Valeriana sitchensis)</i>	T	0-5	23
GLOBEFLOWER <i>(Trollius albiflorus)</i>	2	0-7	38
WOOLLY EVERLASTING <i>(Antennaria lanata)</i>	1	0-12	33
<b>GRASSES</b>			
TUFTED HAIRGRASS <i>(Deschampsia cespitosa)</i>	1	0-4	29
CALIFORNIA OATGRASS <i>(Danthonia californica)</i>	2	0-10	19
SEDGE <i>(Carex spp.)</i>	23	0-73	95

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
MESIC -HYGRIC

NUTRIENT REGIME:  
PERMESOTROPHIC

ELEVATION:  
1793(1370-2110) M

SOIL DRAINAGE:  
MOD. WELL

**FORAGE PRODUCTION KG/HA**

<p>SUGGESTED GRAZING CAPACITY NON-USE</p>
---

## SACMB3. Willow-Bog birch/Hairy wildrye

*(Salix glauca-Bog birch/Elymus innovatus)*

n=17 This community is typical of the valley bottoms where the low temperatures prohibit the growth of trees. Corns and Achuff (1982) described a similar community in the Banff and Jasper National Parks. They found this community type occupied coarse stream deposits which had repeated flooding.

Bork (1994) felt this community type developed from the invasion of willow and bog birch onto grasslands in the absence of disturbance in Willmore Wilderness park. Willow cover has increased, shading the growth of grasses and allowing tall-growing forbs, such as fireweed, aster and veiny meadow rue to increase. He felt continued protection from disturbance will allow succession to shrub and eventually tree species. This community is typical of the valley bottoms throughout the subalpine in both the foothills and mountain ecodistricts.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>TREES</b>			
ENGELMANN SPRUCE <i>(Picea engelmannii)</i>	T	0-1	18
<b>SHRUBS</b>			
WILLOW SPP. <i>(Salix glauca)</i>	36	0-85	36
BOG BIRCH <i>(Betula glandulosa)</i>	17	0-65	59
SHRUBBY CINQUEFOIL <i>(Potentilla fruticosa)</i>	2	0-12	47
<b>FORBS</b>			
NORTHERN VALERIAN <i>(Valeriana dioica)</i>	T	0-1	12
YARROW <i>(Achillea millefolium)</i>	1	0-4	71
FIREWEED <i>(Epilobium angustifolium)</i>	3	0-20	59
STRAWBERRY <i>(Fragaria virginiana)</i>	5	0-20	82
TALL LARKSPUR <i>(Delphinium glaucum)</i>	3	0-10	65
VEINY MEADOW RUE <i>(Thalictrum venulosum)</i>	T	0-2	6
<b>GRASSES</b>			
BOG SEDGE <i>(Kobriesia myosuroides)</i>	1	0-13	12
HAIRY WILDRYE <i>(Elymus innovatus)</i>	17	0-50	94
SEDGE <i>(Carex spp)</i>	1	0-5	23

TUFTED HAIR GRASS

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:  
SUBHYGRIC

NUTRIENT REGIME:  
MESOTROPHIC

ELEVATION:  
1859(1400-2240) M

SOIL DRAINAGE:  
WELL TO MODERATELY WELL

### FORAGE PRODUCTION KG/HA

SUGGESTED GRAZING CAPACITY NON-USE
---------------------------------------

**SACMB4. Willow-Bog birch/Bog sedge**  
*(Salix spp.-Betula glandulosa/Kobresia myosuroides)*

n=14 This community type was described on moist lowland sites at higher elevations in the Central and Northern Rocky Mountain ecodistricts. It appears this community type originated from recent shrub encroachment onto sedge-bog sedge-tufted hairgrass community type. The presence of bog sedge may represent the transition between the foothills ecodistricts and the rocky mountain ecodistricts. Corns and Achuff (1982) described bog sedge dominated community types in the Central and Northern Rocky Mtn. ecodistricts of the Subalpine subregion of Banff and Jasper National Parks.

**PLANT COMPOSITION** **CANOPY COVER(%)**  
**MEAN RANGE CONST.**

<b>SHRUBS</b>			
WILLOW SPP. <i>(Salix spp.)</i>	24	0-55	93
BOG BIRCH <i>(Betula glandulosa)</i>	15	0-55	71
SHRUBBY CINQUEFOIL <i>(Potentilla fruticosa)</i>	7	0-20	79
<b>FORBS</b>			
ALPINE BISTORT <i>(Polygonum viviparum)</i>	1	0-5	64
BEARBERRY <i>(Arctostaphylos uva-ursi)</i>	3	0-15	50
SMOOTH LEAVED CINQUEFOIL <i>(Potentilla diversitolia)</i>	1	0-10	29
ALPINE HEDYSARUM <i>(Hedysarum alpinum)</i>	3	0-15	71
STRAWBERRY <i>(Fragaria virginiana)</i>	3	0-15	71
<b>GRASSES</b>			
BOG SEDGE <i>(Kobresia myosuroides)</i>	26	7-45	100
SEDEGE <i>(Carex spp.)</i>	3	0-4	71
TUFTED HAIRGRASS <i>(Deschampsia cespitosum)</i>	2	0-10	29
CALIFORNIA OATGRASS <i>(Danthonia californica)</i>	2	0-10	21

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
MESIC-SUBHYGRIC

NUTRIENT REGIME:  
MESOTROPHIC

ELEVATION:  
1901(1700-2280) M

SOIL DRAINAGE:  
MODERATELY WELL

**FORAGE PRODUCTION KG/HA**

SUGGESTED GRAZING CAPACITY  
NON-USE

**SACMB5. Bog birch/Juniper**  
*(Betula glandulosa/Juniperus communis)*

**n=6** This community type is found on rocky exposures with westerly aspects in association with Engelmann spruce and lodgepole pine forests. The soils are very shallow and rapidly drained.

**PLANT COMPOSITION CANOPY COVER(%)**

MEAN RANGE CONST.

**SHRUBS**

**BOG BIRCH**

*(Betula glandulosa)* 38 0-65 83

**WILLOW SPP.**

*(Salix spp.)* 7 0-4 50

**FORBS**

**YARROW**

*(Achillea millefolium)* 1 0-2 33

**FIREWEED**

*(Epilobium angustifolium)* 2 1-2 100

**TALL LARKSPUR**

*(Delphinium glaucum)* T 0-1 17

**ALPINE GOLDENROD**

*(Solidago multiradiata)* T 0-1 17

**TWINFLOWER**

*(Linnaea borealis)* 1 0-3 33

**GRASSES**

**ROCKY MOUNTAIN FESCUE**

*(Festuca brachyphylla)* T 0-1 17

**SEDGE**

*(Carex spp.)* 2 0-3 80

**HAIRY WILD RYE**

*(Elymus innovatus)* 3 0-15 50

**ENVIRONMENTAL VARIABLES**

**MOISTURE REGIME:**

SUBXERIC

**NUTRIENT REGIME:**

MESOTROPHIC

**ELEVATION:**

1998(1900-2130) M

**SOIL DRAINAGE:**

RAPIDLY

**SLOPE:**

35(10-50)%

**ASPECT:**

WESTERLY

**FORAGE PRODUCTION KG/HA**

SUGGESTED GRAZING CAPACITY  
NON-USE

## SACMB6. Willow/Forb

(*Salix spp./Trollius albiflorus, Erigeron peregrinus, Mountain heliotrope*)

n=6 This community type results from the invasion of willow onto the forb dominated meadows (SACMA2) previously described. These meadows occupy mesic to subhygric, gently sloping, upper subalpine to alpine areas. The soils are imperfectly to well drained Gleysols on fluvial and morainal landforms (Corns and Achuff 1982). On the poorly to imperfectly drained sites in areas where snow melts late and seepage is received throughout the growing season mountain marigold and globeflower predominate. In contrast on better drained, drier sites wandering daisy and mountain heliotrope predominate.

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

#### SHRUBS

BARRET'S WILLOW . ( <i>Salix barrattiana</i> )	40	10-75	100
SMOOTH WILLOW ( <i>Salix glauca</i> )	17	0-40	67
BOG BIRCH ( <i>Betula glandulosa</i> )	2	0-5	33

#### FORBS

GLOBEFLOWER ( <i>Trollius albiflorus</i> )	8	2-15	100
WANDERING DAISY ( <i>Erigeron peregrinus</i> )	4	0-10	83
MOUNTAIN SAGE ( <i>Artemisia norvegica</i> )	6	0-15	50
MOUNTAIN HELIOTROPE ( <i>Valeriana sitchensis</i> )	4	0-15	50
SMOOTH LEAVED CINQUEFOIL ( <i>Potentilla diversifolia</i> )	4	0-10	50
MOUNTAIN MARIGOLD ( <i>Caltha leptosepala</i> )	2	0-2	33

#### GRASSES

ALPINE TIMOTHY ( <i>Phleum commutatum</i> )	1	0-1	67
SEDGE ( <i>Carex spp.</i> )	10	0-50	83
SIMPLE BOG SEDGE ( <i>Kobresia simpliciuscula</i> )	13	0-80	17

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:

MESIC

NUTRIENT REGIME:

PERMESOTROPHIC

ELEVATION:

2103(1960-2320) M

SOIL DRAINAGE:

IMPERFECTLY

### FORAGE PRODUCTION KG/HA

SUGGESTED GRAZING CAPACITY

NON-USE



**SACMB7. Grouseberry-Juniper**  
(*Vaccinium scoparium*-*Juniperus communis*)

n=4 This is a timberline community type found in conjunction with small patches of subalpine fir. Ogilvie (1969) described a heath-grouseberry community occurring among tree islands and krummholz colonies, on lee slopes with very deep snow accumulation.

**PLANT COMPOSITION CANOPY COVER(%)**

	MEAN	RANGE	CONST.
<b>TREES</b>			
SUBALPINE FIR ( <i>Abies lasiocarpa</i> )	1	0-4	25
<b>SHRUBS</b>			
SMOOTH WILLOW ( <i>Salix glauca</i> )	1	0-2	50
GROUSEBERRY ( <i>Vaccinium scoparium</i> )	14	7-20	100
GROUND JUNIPER ( <i>Juniperus communis</i> )	5	0-12	100
CROWBERRY ( <i>Emptrum nigrum</i> )	2	0-8	25
<b>FORBS</b>			
FIREWEED ( <i>Epilobium angustifolium</i> )	12	5-20	100
SMALL LEAVED EVERLASTING ( <i>Antennaria parviflora</i> )	3	1-8	100
STRAWBERRY ( <i>Fragaria virginiana</i> )	3	1-5	100
YARROW ( <i>Achillea millefolium</i> )	2	0-2	100
<b>GRASSES</b>			
SPIKED TRisetum ( <i>Trisetum spicatum</i> )	7	1-20	100
SEDGE ( <i>Carex spp.</i> )	1	0-3	75
CALIFORNIA OATGRASS ( <i>Danthonia californica</i> )	6	0-15	50

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
SUBXERIC-MESIC

NUTRIENT REGIME:  
MESOTROPHIC

ELEVATION:  
2157(2080-2260) M

SOIL DRAINAGE:  
WELL

**FORAGE PRODUCTION KG/HA**

SUGGESTED GRAZING CAPACITY  
NON-USE

## SACMB8. Subalpine fir

(*Abies lasiocarpa*)

**n=22** This is a timberline community type found in conjunction with small patches of the grouseberry-juniper community type. The trees tend to be very small and shrub like. Ogilvie (1969) described timberline as a gradual breaking -up of the forest into groves, tree islands, low stunted krummholz colonies, and finally dwarfed isolated trees. Ogilvie found timberline to occur high on lee slopes and low on wind-exposed slopes, south facing slopes, avalanche slopes, along stream bottoms and on unstable substrata such as scree and rubble. The major environmental factors controlling timberline are low temperature, wind dessication, avalanching and snow depth.

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

#### TREES

SUBALPINE FIR  
(*Abies lasiocarpa*)      29      0-20      39

#### SHRUBS

SMOOTH WILLOW  
(*Salix glauca*)      1      0-10      22

GROUSEBERRY  
(*Vaccinium scoparium*)      4      0-15      22

WHITE FLOWERED RHODODENDRON  
(*Rhododendron albiflorum*)<sup>4</sup>      0-30      22

GROUND JUNIPER  
(*Juniperus communis*)      2      0-15      44

#### FORBS

FIREWEED  
(*Epilobium angustifolium*)<sup>2</sup>      0-10      61

MOUNTAIN SAGE  
(*Artemisia norvegica*)      3      0-10      61

STRAWBERRY  
(*Fragaria virginiana*)      1      0-5      48

WANDERING DAISY  
(*Erigeron peregrinus*)      2      0-13      52

#### GRASSES

SPIKED TRisetum  
(*Trisetum spicatum*)      T      0-5      22

SEDGE  
(*Carex spp.*)      2      0-20      63

HAIry WILDrye  
(*Elymus innovatus*)      2      0-20      30

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:  
MESIC

NUTRIENT REGIME:  
MESOTROPHIC

ELEVATION:  
1945(1610-2340) M

SOIL DRAINAGE:  
WELL

### FORAGE PRODUCTION KG/HA

SUGGESTED GRAZING CAPACITY  
NON-USE

**SUBALPINE SUBREGION**  
**SOUTHERN ROCKY MOUNTAINS**  
**NATIVE GRASSLANDS AND SHRUBLANDS**



**Figure 5.** This figure is typical of Rough fescue-Sedge community on mesic sites in the Southern Rocky Mountains of the Subalpine subregion. On steeper slopes rough fescue and hairy wildrye predominate.

## **Native grass and shrubland ecology of the Southern Rocky Mountains**

The ecosites and ecosite phases of the native grass and shrubland community types in the Southern Rocky Mountains of the Subalpine subregion (Table 4) are found in the valley bottoms adjacent to streams and rivers and on south facing slopes. This also includes an area that was classified as Lower Foothills, west of Turner Valley. The grass and shrublands within this area more closely resemble the subalpine than the Lower Foothills.

The grassland and shrubland community types in this ecodistrict are strongly influenced by the lower Montane subregion. Many of the grass species associated with the Montane (rough fescue, Parry oatgrass, Idaho fescue) are associated with the grassland community types described in this ecodistrict. On the wet, imperfectly drained lower slope positions the grass and shrubland communities are very similar to the water sedge and willow/water sedge communities described in the northern ecodistricts. It is the grasslands of the south facing slopes that are different between the southern and northern ecodistricts. In the northern ecodistricts the grasslands of south facing slopes are dominated by hairy wildrye, junegrass and shrubby cinquefoil. In contrast, the grasslands of the southern ecodistrict are dominated by rough fescue, bearberry, hairy wildrye and sedge species.

On gentler south-facing slopes at lower elevations rough fescue and sedge dominate the grassland community types. On more mesic sites within this community Richardson needlegrass may become co-dominant with rough fescue. In contrast at higher elevations on steeper slopes hairy wildrye replaces sedge and Richardson needlegrass as the co-dominant species.

At higher elevations just north of Waterton Lakes National Park the windswept ridges are dominated by Idaho fescue-Junegrass, Bearberry and White mountain avens to form the Fescue-Junegrass/Bearberry and White mountain avens community types. These community types are important wintering areas for bighorn sheep.

Many of these subalpine grass and shrublands are very fragile because of exposure and cold climate. The forage productivity is generally only half of what is found in the lower Montane subregion and recovery from overgrazing will likely take some time because of the poor growing conditions. Grazing pressure causes rough fescue to decline and allows sedge and hairy wildrye to dominate the community. On moist sites heavy grazing pressure allows Kentucky bluegrass to invade.

The carrying capacity, moisture and nutrient regime of the grass and shrubland communities found in the Southern Rocky Mountains of the Subalpine subregion are outlined in Table 5.

Table 4. Ecosites, ecosite phases and community types for the Subalpine subregion in Southwestern Alberta (adapted from Archibald et al. 1996)(highlighted communities are described in this guide, non-highlighted communities are outlined in guide to Ecosites of Southwestern Alberta)

Ecosite	Ecosite Phase	Forested Plant Community Type	Range plant community type	Grazing Succession
a lichen (xeric/poor)	a1 lichen P1	a1.1 P1/juniper/lichen	SASME1 P1/Juniper	
	a2 grassland		SASMA13 Fescue-Junegrass/ Early yellow locoweed SASMA14 White Mtn. avens	
b bearberry/hairy wild rye (submesic/medium)	b1 bearberry/hairy wild rye P1	b1.1 P1/bearberry/hairy wild rye		
	b2 grassland		SASMA2 Rough fescue-Sedge SASMA3 Rough fescue-Hairy wildrye-Sedge SASMA9 Rough fescue-Sedge/Bearberry SASMA10 Parry oatgrass-Rough fescue-Sedge SASMA3a Hairy wildrye-R.fescue-Sedge	SASMA4 Sedge-Hairy wildrye-Slender wheatgrass SASMA11 Sedge/Bearberry SASMC2 Parry oatgrass-Kentucky bluegrass-Sedge SASMC4 Fringed sage/Kentucky bluegrass-Parry oatgrass SASMC9 Idaho fescue-Rough fescue/Bearberry
c subalpine larch/heather (submesic/poor)	b3 shrubland		SASMB2 Willow/Richardson needlegrass	
	c1 subalpine larch/heather La-Fa	c1.1 La-Fa/heather-grouse-berry		
	c2 yellow mountain avens		SASMA6 Yellow mountain avens	

cc rough fescue (mesic/rich)	cc1 rough fescue		SASMA8 Rough fescue-Idaho fescue-Parry oatgrass	SASMA5 Kentucky bluegrass/Dandelion SASMC1 Parry oatgrass- Rough fescue-Kentucky bluegrass SASMC3 Meadow foxtail- Kentucky bluegrass SASMC5 Rough fescue- Kentucky bluegrass SASMC6 Kentucky bluegrass- Rough fescue SASMC7 Timothy-Slender wheatgrass/Fireweed
	cc2 shrubland		SASMA12 Silverberry-Rose	
	cc3 forb meadow		SASMA16 Forb meadow	
d spruce/heather (mesic/poor)	d1 spruce/heather Se	d1.1 Se/heather		
e false azalea-grouse- berry (mesic/medium)	e1 false azalea-grouse- berry P1	e1.1 P1/green alder/armica		
		e1.2 P1/grouse-berry/feather moss		
		e1.3 P1/low bilberry		
		e1.4 P1/false azalea/feather moss	SASME3 P1-Se/Moss	
		e1.5 P1/pine grass	SASME2 P1/Pinegrass	
		e1.6 P1/Canada buffalo-berry		
	e2 false azalea-grouse- berry Pw	e2.1 Pw/false azalea	SASMB3 Whitebark pine	


e3 false azalea-grouse- berry Se	e3.1	Se/grouse-berry/feather moss			
	e3.2	Se/low bilberry/feather moss			
	e3.3	Se/green alder/feather moss	SASME4 Sw-Aw/Alder/Hairy wildrye		
	e3.4	Se/false azalea/feather moss			
	e3.5	Se/Canada buffalo- berry/feather moss			
	e3.6	Se/stair-step moss			
	e3.7	Se/wiry fern moss			
e4 false azalea-grouse- berry Fa	e4.1	Fa/grouse-berry/feather moss			
	e4.2	Fa/false azalea/feather moss			
e5 deciduous			SASMD1 Pb/Silverberry		
			SASMD2 Aw/Rose/Pinegrass	SASMD3 Aw/Fireweed/ Meadow foxtail SASMD4 Aw/Rose/ Canada bluegrass	
			SASMA15 Pinegrass-Hairy wildrye/Strawberry		
f thimbleberry (subhygric/rich)	e6 grassland				
	f1 thimbleberry P1	P1/thimbleberry			
	f2 thimbleberry Fa-Se	f2.1 Fa-Se/Thimbleberry			

	f3 thimbleberry Aw		SASMD2 Aw-Pb/Cow parsnip	SASMD6 Aw-Pb/Cow parsnip/Timothy
	f4 shrubby seepages		SASMA7a Marsh reedgrass/Cow parsnip	SASMC8 Marsh reedgrass-Timothy/Cow parsnip
g dwarf birch/tufted hair grass (hygric/rich)	g1 dwarf birch/tufted hair grass	g1.1 dwarf birch/tufted hair grass	SASMA7 Tufted hairgrass-Sedge SASMA1a Beaked sedge-Alpine foxtail-Tufted hairgrass	SASMB4 Willow-Bog birch/R. fescue- Kentucky bluegrass SASMA5 Kentucky bluegrass/Dandelion
h horsetail (subhydric/rich)	h1 horsetail Se	h1.1 Se/horsetail/feather moss		
	h2 horsetail fen	h2.1 dwarf birch/sedge/golden moss		
i fen (subhydric/rich)	i1 shrub fen		SASMB1 Willow/Sedge SASMB5 Willow/Marsh reedgrass	
	i2 graminoid fen		SASMA1 Water sedge	



**a2 grassland (n=29)**

---



**CHARACTERISTIC SPECIES**

---

**Shrubs**

[ 2 ] Shrubby cinquefoil

**Forbs**

- [ 15 ] White mtn. avens
- [ 1 ] Bearberry
- [ 1 ] False dandelion
- [ 2 ] Spotted saxifrage
- [ 1 ] Sandwort
- [ 1 ] Kittenail
- [ 2 ] Early yellow locoweed

**Graminoids**

- [ 3 ] Sedge
- [ 1 ] Junegrass
- [ 3 ] Idaho fescue
- [ 1 ] Rough fescue

**SITE CHARACTERISTICS**

---

**Moisture regime:** subxeric, submesic

**Nutrient regime:** poor, very poor

**Topographic position:** upper slope, crest

**Slope:** (6-40)

**Aspect:** variable

**SOIL CHARACTERISTICS**

---

**Organic thickness:** (0-2)

**Humus form:** not available

**Surface texture:** SL,S,L

**Effective texture:** SL,LS

**Depth to Mottles/Gley:** none

**Drainage:** rapid, well

**Parent material:** M,C/X

**Soil subgroup:** O,R, non-soil

**COMMUNITY TYPES**


---

SASMA13 Fescue-Junegrass/Early yellow

locoweed(n=17)

SASMA14 White mtn. avens(n=12)

**b2 grassland (n=52)**



**CHARACTERISTIC SPECIES**

**Shrubs**

[ 8 ] Shrubby cinquefoil

**Forbs**

- [ 4 ] Yellow beardtongue
- [ 17 ] Bearberry
- [ 2 ] Yellow hedysarum
- [ 2 ] Brown bracted everlasting
- [ 2 ] Yarrow
- [ 1 ] Kittentail
- [ 2 ] Strawberry

**Graminoids**

- [ 9 ] Sedge
- [ 1 ] Junegrass
- [ 2 ] Idaho fescue
- [ 20 ] Rough fescue
- [ 9 ] Parry oatgrass
- [ 2 ] Richardson needlegrass

**SITE CHARACTERISTICS**

**Moisture regime:** subxeric, submesic, mesic  
**Nutrient regime:** medium  
**Topographic position:** upper slope, crest  
**Slope:** (0-50)  
**Aspect:** south, west


**SOIL CHARACTERISTICS**

**Organic thickness:** (0-2)(2-10)  
**Humus form:** mull  
**Surface texture:** SL, LS  
**Effective texture:** SL,LS  
**Depth to Mottles/Gley:** none  
**Drainage:** rapid, well  
**Parent material:** M,C, GF  
**Soil subgroup:** O.R, O.EB, O.MB

**COMMUNITY TYPES**

- SASMA2 Rough fescue-Sedge(n=23)
- SASMA3 Rough fescue-Hairy wildrye-Sedge(n=9)
- SASMA3a Hairy wildrye-Rough fescue-Sedge(n=1)
- SASMA9 Rough fescue-Sedge/Bearberry(n=7)
- SASMA10 Parry oatgrass-Rough fescue-Sedge(n=12)

**b2a grazed grassland (n=33)**



**CHARACTERISTIC SPECIES**

**Shrubs**

- [ 4 ] Shrubby cinquefoil
- [ 1 ] Saskatoon
- [ 1 ] Juniper

**Forbs**

- [ 6 ] Old man's whiskers
- [ 8 ] Bearberry
- [ 2 ] Yellow hedysarum
- [ 1 ] Brown bracted everlasting
- [ 4 ] Yarrow
- [ 1 ] Sandwort
- [ 2 ] Strawberry

**Graminoids**

- [ 9 ] Sedge
- [ 1 ] Junegrass
- [ 2 ] Idaho fescue
- [ 5 ] Rough fescue
- [ 4 ] Parry oatgrass
- [ 3 ] Kentucky bluegrass
- [ 1 ] Timothy

**SITE CHARACTERISTICS**

**Moisture regime:** subxeric, submesic, mesic  
**Nutrient regime:** medium  
**Topographic position:** upper slope, crest  
**Slope:** (0-50)  
**Aspect:** south, west

**SOIL CHARACTERISTICS**


**Organic thickness:** (0-2)(2-10)  
**Humus form:** mull  
**Surface texture:** SL, LS  
**Effective texture:** SL,LS  
**Depth to Mottles/Gley:** none  
**Drainage:** rapid, well  
**Parent material:** M,C, GF  
**Soil subgroup:** O.R, O.EB, O.MB

**COMMUNITY TYPES**

SASMA4 Sedge-Hairy wildrye-Slender wheatgrass(n=15)  
 SASMA11 Sedge/Bearberry(n=11)  
 SASMC2 Parry oatgrass-Kentucky bluegrass-Sedge(n=4)  
 SASMC4 Fringed sage/Kentucky bluegrass-Parry oatgrass(n=1)  
 SASMC9 Idaho fescue-Rough fescue/Bearberry (n=2)

**b3 shrubland (n=2)**

**CHARACTERISTIC SPECIES**



**Moisture regime:** submesic, mesic

**Nutrient regime:** medium

**Topographic position:** lower slope

**Slope:** (0-5)

**Aspect:** south, west

**SOIL CHARACTERISTICS**

**Organic thickness:** (0-2)(2-10)

**Humus form:** mull

**Surface texture:** SL, LS

**Effective texture:** SL,LS

**Depth to Mottles/Gley:** none

**Drainage:** well

**Parent material:** F, GF, C

**Soil subgroup:** O.R, O.EB

**COMMUNITY TYPES**

SASMB2 Willow/Richardson needlegrass(n=2)

**Shrubs**

[ 30 ] Willow

[ 5 ] Bog birch

[ 5 ] Shrubby cinquefoil

**Forbs**

[ 1 ] Northern valerian

[ 1 ] Yarrow

[ 1 ] Fireweed

[ 2 ] Strawberry

**Graminoids**

[ 3 ] Sedge

[ 8 ] Rocky mtn. fescue


[ 45 ] Richardson needlegrass

[ 1 ] Slender wheatgrass

**SITE CHARACTERISTICS**

**c2 yellow mtn. avens (n=1)**

---



**SITE CHARACTERISTICS**

---

Moisture regime: subseric, submesic  
 Nutrient regime: poor, medium  
 Topographic position: floodplain  
 Slope: (0-5)  
 Aspect: variable

**SOIL CHARACTERISTICS**

---

Organic thickness: (0-2)  
 Humus form: mor  
 Surface texture: SL,SiL  
 Effective texture: SL,  
 Depth to Mottles/Gley: none  
 Drainage: rapid, well  
 Parent material: F,GF  
 Soil subgroup: O.R

**COMMUNITY TYPES**

---

SASMA6 Yellow mtn. avens (n=1)

**CHARACTERISTIC SPECIES**

---

**Shrubs**

[ 4 ] Shrubby cinquefoil

**Forbs**

- [ 13 ] Yellow mtn. avens
- [ 5 ] Late yellow locoweed
- [ 3 ] Silvery cinquefoil
- [ 3 ] Low goldenrod
- [ 3 ] Yellow hedysarum


**Graminoids**

- [ 67 ] Sedge
- [ 6 ] Hairy wildrye
- [ 5 ] Fringed brome
- [ 1 ] Slender wheatgrass

cc rough fescue (n=51)

GENERAL DESCRIPTION

This ecosite is typical of south and west facing slopes and lower slope positions throughout the southern mountains of the Subalpine subregion from an elevation of 1500 m to 1900 m. This ecosite is usually dominated by grass species because of the dry site conditions and westerly winds. The soils of this ecosite are dominated by deep black soils. A number of rough fescue dominated sites have not had the species composition change in over 30 years of no disturbance in the Montane subregion indicating the climax nature of this ecosite.




SUCCESSIONAL RELATIONSHIPS

Due to the nature of the site grasslands often remain the climax vegetation on these sites. On moister sites shrubs and trees such as saskatoon, snowberry, chokecherry and aspen often invade the site. Heavy grazing pressure on these grasslands can often lead to a degraded site that is dominated by Kentucky bluegrass, timothy and clover species.

INDICATOR SPECIES

rough fescue	Kentucky bluegrass
Parry oatgrass	Timothy
Idaho fescue	Dandelion
Old man's whiskers	Cut leaved anemone
Silverberry	Rose

mesic/rich



SITE CHARACTERISTICS

**Moisture regime:** submesic, mesic  
**Nutrient regime:** rich, medium  
**Topographic position:** crest, midslope, lower slope  
**Slope:** (0-2%)<sup>3</sup>(16-30%)<sup>3</sup> (31-45%)<sup>1</sup>(45-70%)<sup>1</sup>  
**Aspect:** south, southwest, west


SOIL CHARACTERISTICS

**Organic thickness:** (0-2)(2-5)  
**Humus form:** mull  
**Surface texture:** CL, SiL, L  
**Effective texture:** CL, SiL, SL,  
**Depth to Mottles/Gley:** none  
**Drainage:** well  
**Parent material:** F, GF, M  
**Soil subgroup:** O.BL, O.DG, O.MB

ECOSITE PHASES

cc1 rough fescue (n=4)  
 cc1a grazed rough fescue(n=46)  
 cc2 shrubland(n=1)

**cc1 rough fescue (n=4)**



**SITE CHARACTERISTICS**

**Moisture regime:** submesic, mesic

**Nutrient regime:** rich, medium

**Topographic position:** crest, midslope, lower slope

**Slope:** (0-2%)<sup>3</sup>(16-30%)<sup>3</sup> (31-45%)<sup>1</sup>(45-70%)<sup>1</sup>

**Aspect:** south, southwest, west

**SOIL CHARACTERISTICS**

**Organic thickness:** (0-2)(2-5)

**Humus form:** mull

**Surface texture:** CL, SiL, L

**Effective texture:** CL, SiL, SL,

**Depth to Mottles/Gley:** none

**Drainage:** well

**Parent material:** F, GF, M

**Soil subgroup:** O.BL, O.DG, O.MB

**COMMUNITY TYPES**

SASMA8 Rough fescue-Idaho fescue-Parry oatgrass(n=4)

**CHARACTERISTIC SPECIES**

**Shrubs**

- [ 6 ] Shrubby cinquefoil
- [ 4 ] Rose
- [ 3 ] Saskatoon


**Forbs**

- [ 15 ] Showy aster
- [ 10 ] Strawberry
- [ 7 ] Sticky purple geranium
- [ 1 ] White mtn. avens
- [ 3 ] Northern bedstraw

**Graminoids**

- [ 23 ] Rough fescue
- [ 10 ] Idaho fescue
- [ 4 ] Parry oatgrass
- [ 3 ] Sedge

**cc1a grazed rough fescue (n=46)**



**CHARACTERISTIC SPECIES**

**Shrubs**

- [ 4 ] Shrubby cinquefoil
- [ 2 ] Rose
- [ 1 ] Saskatoon

**Forbs**

- [ 4 ] Dandelion
- [ 1 ] Strawberry
- [ 2 ] Old man's whiskers
- [ 2 ] Yarrow
- [ 2 ] Clover
- [ 3 ] Graceful cinquefoil

**Graminoids**

- [ 8 ] Rough fescue
- [ 1 ] Idaho fescue
- [ 2 ] Parry oatgrass
- [ 1 ] Sedge
- [ 15 ] Kentucky bluegrass
- [ 8 ] Timothy

**SITE CHARACTERISTICS**

**Moisture regime:** submesic, mesic  
**Nutrient regime:** rich, medium  
**Topographic position:** crest, midslope, lower slope  
**Slope:** (0-2%)<sup>5</sup>(16-30%)<sup>3</sup> (31-45%)<sup>1</sup>(45-70%)<sup>1</sup>  
**Aspect:** south, southwest, west

**SOIL CHARACTERISTICS**

**Organic thickness:** (0-2)(2-5)  
**Humus form:** mull  
**Surface texture:** CL, SiL, L  
**Effective texture:** CL, SiL, SL,  
**Depth to Mottles/Gley:** none  
**Drainage:** well  
**Parent material:** F, GF, M  
**Soil subgroup:** O.BL, O.DG, O.MB

**COMMUNITY TYPES**


- SASMA5 Kentucky bluegrass/Dandelion(n=14)
- SASMC1 Parry oatgrass-Rough fescue-Kentucky bluegrass(n=7)
- SASMC3 Meadow foxtail-Kentucky bluegrass(n=3)
- SASMC5 Rough fescue-Kentucky bluegrass(n=1)
- SASMC6 Kentucky bluegrass-Rough fescue(n=13)
- SASMC7 Timothy-Slender wheatgrass/Fireweed(n=8)



**cc2 shrubland (n=1)**

---

**CHARACTERISTIC SPECIES**



**Topographic position:** crest, midslope

**Slope:** (16-30%)

**Aspect:** south

**SOIL CHARACTERISTICS**

---

**Organic thickness:** (0-2)(2-5)

**Humus form:** mull

**Surface texture:** CL, SiL, L

**Effective texture:** CL, SiL, SL,

**Depth to Mottles/Gley:** none

**Drainage:** well

**Parent material:** F, GF, M

**Soil subgroup:** O.BL, O.DG, O.MB

**COMMUNITY TYPES**

---

SASMA15 Silverberry-Rose (n=1)

**Trees**

[ 8 ] Aspen

**Shrubs**

[ 15 ] Silverberry

[ 5 ] Rose

**Forbs**

[ 6 ] Chickweed

[ 2 ] False dandelion

[ 1 ] Silver plant

**Graminoids**

[ 1 ] Rocky mtn. fescue

[ 9 ] Bluebunch wheatgrass


**SITE CHARACTERISTICS**

---

**Moisture regime:** mesic

**Nutrient regime:** rich

**cc3 Forb meadow (n=1)**



**SITE CHARACTERISTICS**

**Moisture regime:** mesic, subhygric

**Nutrient regime:** rich, medium

**Topographic position:** lower slope

**Slope:** (0-8%)

**Aspect:** south, level

**SOIL CHARACTERISTICS**

**Organic thickness:** (6-15)

**Humus form:** moder

**Surface texture:** SL, SiL, L, C

**Effective texture:** CL, SiL, SCL,

**Depth to Mottles/Gley:** none, (0-25)

**Drainage:** well, mod. well

**Parent material:** F, C, M, X

**Soil subgroup:** O.EB, E.DYB, O.EB, BR.GL

**COMMUNITY TYPES**

SASMA16 Forb meadow(n=1)

**CHARACTERISTIC SPECIES**

**Shrubs**

- [ 3 ] Shrubby cinquefoil
- [ 2 ] Rose


**Forbs**

- [ 3 ] Graceful cinquefoil
- [ 4 ] Yellow hedsyarum
- [ 2 ] American vetch
- [ 6 ] Fireweed
- [ 1 ] Strawberry

**Graminoids**

- [ 2 ] Parry oatgrass
- [ 3 ] Hairy wildrye
- [ 2 ] Rough fescue

## e5 deciduous (n=21)



## SITE CHARACTERISTICS

**Moisture regime:** mesic

**Nutrient regime:** medium

**Topographic position:** midslope, lower slope, floodplain

**Slope:** (0-5%)

**Aspect:** south, level

## SOIL CHARACTERISTICS

**Organic thickness:** (0-2)(2-5)

**Humus form:** mor

**Surface texture:** SL, SiL, L

**Effective texture:** CL, SiL, SCL,

**Depth to Mottles/Gley:** none

**Drainage:** well, mod. well

**Parent material:** F, C, M

**Soil subgroup:** O.EB, E.DYB, O.EB, BR.GL

## COMMUNITY TYPES

SASMD1 Pb/Silver berry(n=1)

SASMD2 Aw/Rose/Pinegrass(n=20)

## CHARACTERISTIC SPECIES

### Trees

[ 20 ] Aspen

[ 10 ] Balsam poplar

### Shrubs

[ 2 ] Silverberry

[ 7 ] Rose

[ 3 ] White meadowsweet

### Forbs

[ 5 ] American vetch

[ 6 ] Yellow peavine

[ 4 ] Strawberry

[ 4 ] Fireweed

[ 4 ] Showy aster

[ 1 ] Alpine hedysarum


### Graminoids

[ 14 ] Pinegrass

[ 6 ] Hairy wildrye

[ 1 ] Junegrass

## e5a grazed deciduous (n=2)



## CHARACTERISTIC SPECIES

### Trees

- [ 35 ] Aspen
- [ 3 ] White spruce

### Shrubs

- [ 5 ] Gooseberry
- [ 6 ] Rose

### Forbs

- [ 1 ] American vetch
- [ 3 ] Yellow peavine
- [ 3 ] Strawberry
- [ 6 ] Fireweed
- [ 13 ] Lindley aster
- [ 2 ] Dandelion

### Graminoids

- [ 4 ] Pinegrass
- [ 2 ] Kentucky bluegrass
- [ 18 ] Canada bluegrass
- [ 16 ] Meadow foxtail
- [ 9 ] Orchardgrass
- [ 3 ] Timothy

## SITE CHARACTERISTICS

**Moisture regime:** mesic

**Nutrient regime:** medium

**Topographic position:** midslope, lower slope

**Slope:** (0-5%)

**Aspect:** south, level

## SOIL CHARACTERISTICS

**Organic thickness:** (0-2)(2-5)

**Humus form:** mor

**Surface texture:** SL, SiL, L

**Effective texture:** CL, SiL, SCL,

**Depth to Mottles/Gley:** none

**Drainage:** well, mod. well

**Parent material:** F, C, M


**Soil subgroup:** O.EB, E.DYB, O.EB, BR.GL

## COMMUNITY TYPES

SASMD3 Aw/Fireweed/Meadow foxtail(n=1)

SASMD4 Aw/Rose/Canada bluegrass(n=1)

**e6 grassland (n=4)**



**SITE CHARACTERISTICS**

Moisture regime: mesic

Nutrient regime: medium

Topographic position: crest, midslope, lower slope

Slope: (16-30%)

Aspect: southerly

**SOIL CHARACTERISTICS**

Organic thickness: (0-2)(2-5)

Humus form: mull

Surface texture: CL, SiL, L

Effective texture: CL, SiL, SL,

Depth to Mottles/Gley: none

Drainage: well, rapid

Parent material: C, M

Soil subgroup: O.EB, O.MB, O.DG

**COMMUNITY TYPES**

SASMA15 Pinegrass-Hairy

wildrye/Strawberry(n=4)

**CHARACTERISTIC SPECIES**

**Shrubs**

[ 1 ] Shrubby cinquefoil

[ 9 ] Rose

**Forbs**

[ 4 ] Lupine

[ 2 ] Yellow peavine

[ 3 ] Dandelion

[ 1 ] Graceful cinquefoil

[ 3 ] Showy aster

[ 1 ] False mtn. dandelion

**Graminoids**


[ 11 ] Pinegrass

[ 2 ] Junegrass

[ 4 ] Hairy wildrye

[ 3 ] Kentucky bluegrass

### f3 thimbleberry Aw (n=3)



### SITE CHARACTERISTICS

**Moisture regime:** subhygric  
**Nutrient regime:** rich  
**Topographic position:** lower slope  
**Slope:** (0-8%)  
**Aspect:** south, level

### SOIL CHARACTERISTICS

**Organic thickness:** (6-15)  
**Humus form:** moder  
**Surface texture:** SL, SiL, L, C  
**Effective texture:** CL, SiL, SCL,  
**Depth to Mottles/Gley:** none, (0-25)  
**Drainage:** well, mod. well  
**Parent material:** F, C, M, X  
**Soil subgroup:** O.EB, E.DYB, O.EB, BR.GL

### COMMUNITY TYPES

SASMD5 Aw-Pb/Cow parsnip(n=3)

### CHARACTERISTIC SPECIES

#### **Trees**

- [ 30 ] Aspen
- [ 10 ] Balsam poplar

#### **Shrubs**

- [ 6 ] Gooseberry
- [ 1 ] Rose


#### **Forbs**

- [ 48 ] Cow parsnip
- [ 2 ] Yellow peavine
- [ 15 ] Canada violet
- [ 6 ] Fireweed
- [ 22 ] Lindley aster
- [ 12 ] Tall lungwort
- [ 12 ] Tall larkspur

#### **Graminoids**

- [ 1 ] Marsh reedgrass
- [ 2 ] Hairy wildrye
- [ 2 ] Sedge

### f3a grazed thimbleberry Aw (n=1)



### SITE CHARACTERISTICS

**Moisture regime:** subhygic

**Nutrient regime:** rich

**Topographic position:** lower slope, midslope

**Slope:** (0-8%)

**Aspect:** south, level, north

### SOIL CHARACTERISTICS

**Organic thickness:** (6-15)

**Humus form:** moder

**Surface texture:** SL, SiL, L, C

**Effective texture:** CL, SiL, SCL,

**Depth to Mottles/Gley:** none, (0-25)

**Drainage:** well, mod. well

**Parent material:** F, C, M, X

**Soil subgroup:** O.EB, E.DYB, O.EB, BR.GL

### COMMUNITY TYPES

SASMD6 Aw-Pb/Cow\_parsnip/Timothy(n=1)

### CHARACTERISTIC SPECIES

#### Trees

[ 26 ] Aspen

[ 16 ] Balsam poplar

#### Shrubs

[ 4 ] Raspberry

[ 1 ] Rose

#### Forbs

[ 2 ] Cow parsnip

[ 2 ] Yellow peavine

[ 2 ] Canada violet

[ 6 ] Fireweed

[ 19 ] Lindley aster

[ 12 ] White geranium

[ 6 ] Dandelion

#### Graminoids


[ 4 ] Marsh reedgrass

[ 18 ] Timothy

[ 4 ] Kentucky bluegrass

[ 7 ] Smooth wildrye

**f4 shrubby seepage (n=1)**



**Moisture regime:** subhygric

**Nutrient regime:** rich

**Topographic position:** lower slope, midslope

**Slope:** (0-8%)

**Aspect:** south, level, north

**SOIL CHARACTERISTICS**

**Organic thickness:** (6-15)

**Humus form:** moder

**Surface texture:** SL, SiL, L, C

**Effective texture:** CL, SiL, SCL,

**Depth to Mottles/Gley:** none, (0-25)

**Drainage:** well, mod. well

**Parent material:** F, C, M, X

**Soil subgroup:** O.EB, E.DYB, O.EB, BR.GL

**COMMUNITY TYPES**

SASMA7a Marsh reedgrass/Cow parsnip/(n=1)

**CHARACTERISTIC SPECIES**

**Shrubs**

[ 1 ] Raspberry

[ 4 ] Rose

**Forbs**

[ 3 ] Cow parsnip

[ 10 ] Western meadow rue

[ 4 ] Canada violet

[ 25 ] Fireweed

[ 4 ] Horsetail

[ 4 ] Sticky purple geranium

[ 1 ] Dandelion

**Graminoids**

[ 24 ] Marsh reedgrass

[ 15 ] Sedge


[ 5 ] Idaho fescue

[ 2 ] Brome

**SITE CHARACTERISTICS**



**f4a grazed shrubby seepage (n=1)**



**SITE CHARACTERISTICS**

**Moisture regime:** subhygric  
**Nutrient regime:** rich  
**Topographic position:** lower slope, midslope  
**Slope:** (0-8%)  
**Aspect:** south, level, north

**SOIL CHARACTERISTICS**

**Organic thickness:** (6-15)  
**Humus form:** moder  
**Surface texture:** SL, SiL, L, C  
**Effective texture:** CL, SiL, SCL,  
**Depth to Mottles/Gley:** none, (0-25)  
**Drainage:** well, mod. well  
**Parent material:** F, C, M, X  
**Soil subgroup:** O.EB, E.DYB, O.EB, BR.GL

**COMMUNITY TYPES**

SASMC8 Marsh reedgrass-Timothy/Cow  
parsnip/(n=1)

**CHARACTERISTIC SPECIES**

**Shrubs**

[ 6 ] Raspberry

**Forbs**

- [ 28 ] Cow parsnip
- [ 15 ] Western meadow rue
- [ 20 ] Canada violet
- [ 3 ] Fireweed
- [ 1 ] Horsetail
- [ 10 ] White geranium
- [ 3 ] Dandelion

**Graminoids**

- [ 37 ] Marsh reedgrass
- [ 1 ] Sedge
- [ 29 ] Timothy
- [ 2 ] Mountain brome

**i fen** (n=8)(taken from Ecosites of West-Central Alberta)

**GENERAL DESCRIPTION**

The rich and poor fen are combined in this ecosite. The fen ecosite is generally characterized by flowing oxygenated water and alkaline, nutrient-rich conditions. This ecosite occupies level, depressional and lower slope positions where impeded drainage or high water tables enhance the accumulation of organic matter consisting of sedges, golden moss, tufted moss, and brown moss. Black and/or Engelmann spruce dominate the canopy of the treed phase, while dwarf birch or willow form the canopy of the shrubby phase and sedges dominate the graminoid phase.



**SUCCESSIONAL RELATIONSHIPS**


The fen is an early stage in hydarch succession. Species composition, direction, and rate of succession changes with changing hydrologic regime. As with other wetlands, rich fens have slow successional rates so recovery from disturbance may also be slow.

**INDICATOR SPECIES**

- Black spruce
- Engelmann spruce
- Willow
- Labrador tea
- Dwarf birch
- Horsetail
- Sedge
- Golden moss
- Brown moss

Tufted moss

**subhydic/rich**



**SITE CHARACTERISTICS**

- Moisture regime:** subhygric, hygric, subhydic, hydric
- Nutrient regime:** rich, very rich, medium
- Topographic position:** level, depression, toe
- Slope:** level, (0-1%)
- Aspect:** level, southerly, easterly


**SOIL CHARACTERISTICS**

- Organic thickness:** (>80)(60-79)
- Humus form:** mor, peaty mor
- Surface texture:** fibric, mesic
- Effective texture:** fibric, mesic
- Depth to Mottles/Gley:** not applicable
- Drainage:** imperfect, poor, very poor
- Parent material:** O
- Soil subgroup:** R.G, TY.M, TY.F, T.M, T.F, R.HG, O.HG, O.G, F.I.M

**ECOSITE PHASES**

- i1 shrub fen (n=4)
- i2 graminoid fen(n=4)

**ii shrubby fen (n=4)**



**CHARACTERISTIC SPECIES**

**Shrub**

- [ 25 ] Willow
- [ 5 ] Shrubby cinquefoil
- [ 3 ] Dwarf birch

**Forb**

- [ 3 ] Lindley's aster
- [ 6 ] Strawberry
- [ 4 ] Arrow leaved coltsfoot
- [ 2 ] Horsetail
- [ 1 ] Cow parsnip

**Grass**

- [ 25 ] Sedge
- [ 2 ] Tufted hairgrass
- [ 6 ] Baltic rush
- [ 2 ] Tufted hairgrass

**SITE CHARACTERISTICS**

**Moisture regime:** subhygric, subhydic

**Nutrient regime:** very rich, rich medium

**Topographic position:** level, depression

**Slope:** level, (0-2%)

**Aspect:** variable

**SOIL CHARACTERISTICS**

**Organic thickness:** (>80)(60-79)

**Humus form:** mor, peaty mor

**Surface texture:** fibric, mesic

**Effective texture:** fibric, mesic

**Depth to Mottles/Gley:** not applicable

**Drainage:** imperfect, poor, very poor

**Parent material:** O

**Soil subgroup:** R.G, TY.M, TY.F, T.M, T.F, R.HG, O.HG, O.G, F.I.M


**PLANT COMMUNITY TYPES**

SASMB1 Willow/Sedge (n=2)

SASMB5 Willow/Marsh reedgrass(n=2)

**i2**      **graminoid fen (n=4)**

---



**CHARACTERISTIC SPECIES**

---

**Shrub**

[ 1 ] Willow

**Forb**

[ 1 ] Northern bedstraw  
 [ 1 ] Fireweed

**Grass**

[ 81 ] Water sedge  
 [ 4 ] Tufted hairgrass

**SITE CHARACTERISTICS**

---

**Moisture regime:** hygric, subhydric  
**Nutrient regime:** very rich, rich  
**Topographic position:** level, depression  
**Slope:** level, (0-2%)  
**Aspect:** variable

**SOIL CHARACTERISTICS**

---

**Organic thickness:** (>80)(60-79)  
**Humus form:** mor, peaty mor  
**Surface texture:** fibric, mesic  
**Effective texture:** fibric, mesic  
**Depth to Mottles/Gley:** not applicable  
**Drainage:** imperfect, poor  
**Parent material:** O  
**Soil subgroup:** R.G, TY.M, TY.F, T.M, T.F, R.HG, O.HG, O.G, FLM

**PLANT COMMUNITY TYPES**

---

SASMA1 Sedge meadows (n=4)


Table 5. Native grass and shrublands of the Southern Rocky Mountains of the Subalpine subregion

Community number	Community type	Productivity (kg/ha)		Moisture	Drainage	Carrying capacity (ha/AUM)		
		Grass	Forb					
<b>A. GRASSLANDS</b>								
SASMA1.	Water sedge	1636	109	0	1745	Subhydryc	Poorly	Non-use
SASMA1a	Beaked sedge-Alpine foptail-Tufted hairgrass				2000*	Subhygric	Mod. Well	0.5
SASMA2.	Rough fescue-Sedge	1380	676	49	1837	Mesic	Well	0.5
SASMA3.	Rough fescue-Hairy wildrye	-	-	-	1480*	Submesic	Rapidly	0.6
SASMA3a	Hairy wildrye-Rough fescue-Sedge	-	-	-	1225	Submesic	Well	0.7
SASMA4.	Sedge-Hairy wildrye-Slender wheatgrass	814	505	514	1674	Submesic	Well	0.5
SASMA5.	Kentucky bluegrass/Dandelion	1673	593	75	2341	Subhygric	Mod. Well	0.4
SASMA6.	Yellow mountain avens	572	602	0	1372*	Mesic	Rapidly	Non-use
SASMA7	Tufted hairgrass-Sedge	1164	162	0	1326	Subhygric	Mod. Well	0.7
SASMA7a	Marsh reedgrass/Cow parnsip	-	-	-	2000*	Subhygric	Mod. Well	0.5
SASMA8	Rough fescue-Idaho fescue-Parry oatgrass	1146	614	55	1815	Submesic	Well	0.5
SASMA9	Rough fescue-Sedge/Bearberry	725	385	1475	2585	Subxeric	Rapidly	0.3
SASMA10	Parry oatgrass-Sedge	732	702	96	1452	Subxeric	Rapidly	0.6
SASMA11	Sedge/Bearberry	733	391	1051	2175	Xeric	Rapidly	Non-use
SASMA12	Sliverberry-Rose				500*	Subxeric	Rapidly	1.8
SASMA13	Fescue-Junegrass/E.yellow locoweed	512	291	192	994	Subxeric	Very rapidly	Non-use
SASMA14	White mountain avens	36	198	392	626	Very xeric	Rapidly	Non-use
SASMA15	Pinegrass-Hairy wildrye/Strawberry	758	1170	110	2037	Subxeric	Rapidly	0.4
SASMA16	Forb meadow	554	734	125	1413	Submesic	Well	0.6

Community number	Community type	Productivity (kg/ha)			Moisture	Drainage	Carrying capacity (ha/AUM)	
		Grass	Forb	Shrub Total				
<b>B. SHRUBLANDS</b>								
SASMB1.	Willow/Sedge	1695	373	149	2478	Hygric	Poorly	Non-use
SASMB2.	Willow/Richardson needlegrass	-	-	-	N/A	Submesic	Well	Non-use
SASMB3.	Whitebark pine	-	-	-	N/A	Subxeric	Rapidly	Non-use
SASMB4	Willow-Birch/R.fescue-K.bluegrass	600	200	150	950*	Subhygric	Mod. Well	1.1
SASMB5	Willow/Marsh reedgrass	-	-	-	N/A	Subhygric	Imperfectly	Non-use
<b>C. GRAZING MODIFIED</b>								
SASMC1.	Parry oatgrass-Rough fescue-Kentucky bluegrass	1160	712	24	1574	Xeric	Rapidly	0.6
SASMC2.	Parry oatgrass-Kentucky bluegrass-Sedge	919	930	60	1908	Subxeric	Rapidly	0.5
SASMC3.	Meadow foxtail-Kentucky bluegrass	2775	507	-	3282	Mesic	Well	0.3
SASMC4.	Fringed sage/Kentucky bluegrass-Parry oatgrass	704	490	236	1430	Xeric	Rapidly	0.6
SASMC5.	Rough fescue-Kentucky bluegrass	-	-	-	2000*	Mesic	Well	0.5*
SASMC6.	Kentucky bluegrass-Rough fescue	1382	887	14	2258	Mesic	Rapidly	0.4
SASMC7	Timothy-K. bluegrass/Dandelion	3300	940	-	4240	Submesic	Rapidly	0.2
SASMC8	Marsh reedgrass-Timothy/Cow parsnip	4030	863	33	4926	Subhygric	Mod. Well	0.2
SASMC9	Idaho fescue-Rough fescue/Bearberry	1408	862	43	2313	Submesic	Well	0.4

\*Estimate

Windswept ridges



(adapted from Willoughby 1992)

Figure 6. Sequence of grass and shrubland community types from the Montane to the Subalpine subregions of the Southern Rocky Mountain ecodistrict.

**Key to grass and shrubland dominated communities in the Southern mountains of the Subalpine subregion.**

1. Timberline communities or windswept ridges at timberline.....	2
Meadow communities or south facing grasslands at lower elevations.....	4
2. Plant community dominated by Whitebark pine.....SASMB3	
Windswept ridges dominated by white mtn. avens, Idaho fescue or junegrass.....	3
3. Community dominated by white mtn. avens.....SASMA14	
Grass dominated community dominated by Idaho fescue and Junegrass....SASMA13	
4. Moist meadows or gravelly river flats dominated by shrubs, tufted hairgrass, sedge, yellow mtn. avens or silverberry.....	5
Drier south and north facing slopes dominated by rough fescue, parry oatgrass, upland sedges, hairy wildrye, pinegrass or forb dominated meadows with fireweed.....	13
5. Dry gravelly river flats dominated by yellow mtn. avens or silverberry.....	6
Moist sites dominated by willow, bog birch or grassy areas dominated by tufted hairgrass, sedge, marsh reedgrass, kentucky bluegrass and dandelion.....	7
6. Site dominated by yellow mtn. avens.....SASMA6	
Site dominated by silverberry.....SASMA12	
7. Willow or bog birch dominated community types.....	8
Grass dominated meadows (tufted hairgrass, marsh reedgrass, alpine foxtail, beaked sedge, water sedge, kentucky bluegrass).....	11
8. Wetter sites dominated by sedge or marsh reedgrass in the understory.....	9
Drier, sites dominated by richardson needlegrass or rough fescue in understory.....	10
9. Understory dominated by marsh reedgrass.....SASMB5	
Understory dominated by sedge.....SASMB1	
10. Dry south facing slopes in Athabasca river valley near Jasper, dominated by Richardson needlegrass in understory.....SASMB2	
Understory dominated by rough fescue.....SASMB4	
11. Very wet sites dominated by water sedge.....SASMA1	
Drier sites dominated by tufted hairgrass, graceful sedge, marsh reedgrass, alpine foxtail or Kentucky bluegrass.....	12
12. Ungrazed sites dominated by tufted hairgrass, beaked sedge, alpine foxtail, marsh reedgrass, or graceful sedge.....	12a
Grazed sites dominated by Kentucky bluegrass, timothy or dandelion.....	12b
12a Marsh reedgrass and cow parsnip dominated grassy meadows.....SASMA7a	
Tufted hairgrass, beaked sedge, alpine foxtail or graceful sedge dominated meadow.....	12c
12b Marsh reedgrass, timothy, cow parsnip dominated meadows.....SASMC8	
Heavy grazing pressure site dominated by Kentucky bluegrass and dandelion....SASMA5	
12c Tufted hairgrass, graceful sedge dominated meadows.....SASMA7	
Beaked sedge, Alpine foxtail dominated meadow.....SASMA1a	
13. Ungrazed rough fescue, hairy wildrye, pinegrass or forb dominated grasslands.....	14
Moderately to heavily grazed grasslands .....	20
14. Pinegrass or forb (fireweed) dominated meadows .....	15



Rough fescue, parry oatgrass, hairy wildrye, bearberry dominated grasslands.....	16
15 Site dominated by forbs, moist seepage area.....	<b>SASMA16</b>
Pinegrass, hairy wildrye dominated grassland.....	<b>SASMA15</b>
16. Rough fescue dominated grasslands, south and west of Turner valley, hairy wildrye and sedge dominate or co-dominate grassland.....	<b>17</b>
Rough fescue dominated grasslands, west of Porcupine Hills and in Castle area, co- dominated by Parry oatgrass, sedge or bearberry.....	<b>18</b>
17. Grasslands of south facing slopes, sedge co-dominated.....	<b>SASMA2</b>
Grasslands of lower slope positions or north aspects, dominated or co-dominated by hairy wildrye.....	<b>17a</b>
17a Grassland dominated by hairy wildrye, north aspects on steep ridges.....	<b>SASMA3a</b>
Grasslands co-dominated by hairy wildrye, south facing.....	<b>SASMA3</b>
18. Grasslands of lower slope positions dominated by Rough fescue.....	<b>SASMA8</b>
Grasslands of mid to upper slope positions dominated by Parry oatgrass, Idaho fescue or bearberry.....	<b>19</b>
19. Grasslands of midslope position dominated by Parry oatgrass, Idaho fescue....	<b>SASMA10</b>
Grasslands of upper slope positions or hillcrests co-dominated by bearberry....	<b>SASMA9</b>
20. Moderately grazed grasslands native grass species still dominate the site.....	<b>21</b>
Heavily grazed grasslands non-native species (Kentucky bluegrass, timothy, meadow foxtail) dominate or co-dominate the site.....	<b>22</b>
21 Drier sites with bearberry .....	<b>21a</b>
Moister sites dominated by sedge, hairy wildrye and slender wheatgrass..	<b>SASMA4</b>
21a Grassy areas with a high cover of Idaho fescue and Rough fescue.....	<b>SASMC9</b>
Sedge dominates the grass layer.....	<b>SASMA11</b>
22. Old range improvement dominated by meadow foxtail.....	<b>SASMC3</b>
Meadow foxtail not present grazed sites.....	<b>23</b>
23 Kentucky bluegrass or timothy dominated sites.....	<b>24</b>
Kentucky bluegrass only co-dominate, Parry oatgrass or rough fescue or fringed sage dominates.....	<b>25</b>
24. Timothy dominated community.....	<b>SASMC7</b>
Kentucky bluegrass, dandelion dominated.....	<b>SASMC6</b>
25. Rough fescue dominates the site.....	<b>SASMC5</b>
Parry oatgrass or Fringed sage dominate the site.....	<b>26</b>
26. Hillcrests and south facing slopes dominated by fringed sage.....	<b>SASMC4</b>
Parry oatgrass dominated community types.....	<b>27</b>
27. Parry oatgrass with Rough fescue and Kentucky bluegrass, lower slope.....	<b>SASMC1</b>
Little rough fescue present, midslope position.....	<b>SASMC2</b>

## SASMA1. Water sedge (*Carex aquatilis*)

n=4 This community type is found in all ecodistricts of the subalpine. Wet conditions and periodic flooding result in the formation of water sedge meadows. Willow will invade into the drier edges of these meadows to form the Willow/Water sedge community type. These community types are quite productive producing nearly 2000 kg/ha of forage, but the high water table in the spring and summer when these meadows are most palatable limits livestock use. A study in the Yukon found that crude protein on these meadows declined from a high of 10% in May to less than 5% in September (Bailey et al. 1992). As a result, these meadows would be rated as secondary or non-use range.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
WILLOW ( <i>Salix spp.</i> )	T	-	25
<b>FORBS</b>			
FIREWEED ( <i>Epilobium angustifolium</i> )T		0-1	50
NORTHERN BEDSTRAW ( <i>Galium boreale</i> )	1	0-5	25
<b>GRASSES</b>			
WATER SEDGE ( <i>Carex aquatilis</i> )	74	52-93	100
TUFTED HAIRGRASS ( <i>Deschampsia cespitosum</i> )3		0-9	75

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME :  
SUBHYDRIC

NUTRIENT REGIME:  
MESOTROPHIC

ELEVATION:  
1600-1981(1859) M

SOIL DRAINAGE:  
POORLY

### FORAGE PRODUCTION

GRASS 1636 (636-2636)  
FORB 109 (0-218)  
TOTAL 1745(636-2854)

SUGGESTED GRAZING CAPACITY  
NON-USE

## SASMA1a. Beaked sedge-Alpine foxtail-Tufted hairgrass

(*Carex atherodes*-*Alopecurus occidentalis*-*Deschampsia cespitosa*)

n=1 This community type was described in a meadow adjacent to a small creek. It is similar to the previously described water sedge community, but this community type is better drained which favours the growth of beaked sedge, alpine foxtail and tufted hairgrass. On the drier edges of this community type upland grass species like rough fescue and Idaho fescue can be found. Willoughby (1992) has found that beaked sedge is palatable to livestock, which causes these meadows to be regularly grazed. Heavy grazing will allow Kentucky bluegrass, timothy and dandelion to invade onto these sites.

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

#### SHRUBS

WILLOW  
(*Salix spp.*) 1 - 100

#### FORBS

FIREWEED  
(*Epilobium angustifolium*) 3 - 100

MARSH CINQUEFOIL  
(*Poa palustre*) 40 - 100

SMOOTH ASTER  
(*Aster laevis*) 18 - 100

MOUNTAIN CINQUEFOIL  
(*Potentilla diversifolia*) 15 - 100

#### GRASSES

BEAKED SEDGE  
(*Carex atherodes*) 25 - 100

TUFTED HAIRGRASS  
(*Deschampsia cespitosum*) 10 - 100

ALPINE FOXTAIL  
(*Alopecurus occidentalis*) 22 - 100

IDAHO FESCUE  
(*Festuca idahoensis*) 7 - 100

TIMOTHY  
(*Phleum pratense*) 5 - 100

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME :  
SUBHYGRIC

NUTRIENT REGIME:  
PERMESOTROPHIC

ELEVATION:  
1640 M

SOIL DRAINAGE:  
MODERATELY WELL

### FORAGE PRODUCTION

TOTAL 2000 \*ESTIMATE

SUGGESTED GRAZING CAPACITY  
0.5 HA/AUM

## SASMA2. Rough fescue-Sedge

(*Festuca scabrella*-*Carex* spp.)

n=23 This community type appears to be the modal grassland community type found on level and gentle south-facing slopes in the Southern Rocky Mountains of the Subalpine subregion. The dominance of rough fescue indicates that this grassland is transitional from the lower Montane subregion. As one moves up in elevation there is a shift away from a Rough fescue-Parry oatgrass dominated community type to a Rough fescue Sedge dominated community type. Willoughby (1992), found that blunt sedge replaced Parry oatgrass as dominant or codominant on steep south-facing slopes, and Richardson needlegrass replaced Parry oatgrass as codominant on more mesic sites in this area. This community type includes both the Rough fescue-Sedge, Rough fescue-Richardson needlegrass and Rough fescue-Parry oatgrass/Shrubby cinquefoil community types described by Willoughby (1992).

Jaques (1976), described a similar community type from Plateau Mountain to Mount Allan. He felt this community type represented critical wildlife habitat because it remained snow-free for a majority of the winter. Grazing by livestock on these community types should be managed carefully in order to maintain a carryover for wildlife in the winter.

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

#### SHRUBS

SHRUBBY CINQUEFOIL  
(*Potentilla fruticosa*) 5 0-19 84

#### FORBS

YARROW  
(*Achillea millefolium*) 5 0-17 96  
YELLOW BEARDSTONGUE  
(*Penstemon confertus*) 5 0-39 52  
MOUSE EARED CHICKWEED  
(*Cerastium arvense*) 2 0-6 100  
BEARBERRY  
(*Arctostaphylos uva-ursi*) 3 0-25 32  
STRAWBERRY  
(*Fragaria virginiana*) 5 0-11 74  
OLD MANS WHISKER'S  
(*Geum triflorum*) 3 0-18 74

#### GRASSES

ROUGH FESCUE  
(*Festuca scabrella*) 18 2-47 100  
BLUNT SEDGE  
(*Carex obtusata*) 12 0-28 91  
CALIFORNIA OATGRASS  
(*Danthonia californica*) 11 0-43 78  
RICHARDSON NEEDLEGRASS  
(*Stipa richardsonii*) 4 0-22 48

#### HAIRY WILDRYE

(*Elymus innovatus*) 5 0-34 61

### ENVIRONMENTAL VARIABLES

#### MOISTURE REGIME:

SUBMESIC-MESIC

#### NUTRIENT REGIME :

MESOTROPHIC

#### ELEVATION:

1460-2024(1751) M

#### SOIL DRAINAGE:

WELL

#### SLOPE: 0-48(6)%

ASPECT SOUTHERLY

### FORAGE PRODUCTION KG/HA

GRASS 1380 (748-1972)

FORB 676 (224-1412)

SHRUB 49(0-296)

TOTAL 1837(1138-3103)

#### SUGGESTED GRAZING CAPACITY

0.5 HA/AUM, 1.2 AC/AUM

## SASMA3. Rough fescue-Hairy wildrye

(*Festuca scabrella*-*Elymus innovatus*)

**n=9** This community is similar to the Rough fescue-Sedge community type (SASMA2) previously described, but it is found on drier, steeper slopes with poorer soils than the other rough fescue dominated community type. As one moves upslope there is a shift in codominance of sedge to hairy wildrye and an increase in cover of bearberry and juniper.

Corns and Achuff (1982), described hairy wildrye dominated community types on south facing slopes in the more northern ecodistricts. They felt these grasslands occurred on areas with frequent snow avalanching. It is possible that this community type is associated with deeper snow accumulation than the Rough fescue-Sedge dominated type previously described.

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

#### SHRUBS

SHRUBBY CINQUEFOIL.

(*Potentilla fruticosa*) 5 0-23 88

CREeping JUNIPER

(*Juniperus horizontalis*) 2 0-9 33

#### FORBS

BEARBERRY

(*Arctostaphylos uva-ursi*)9 0-16 75

YARROW

(*Achillea millefolium*) 3 2-6 100

STRAWBERRY

(*Fragaria virginiana*) 4 1-9 100

YELLOW HEDYSARUM

(*Hedysarum sulphureum*)3 0-10 56

SILVERY PERENNIAL LUPINE

(*Lupinus argenteus*) 2 0-9 44

#### GRASSES

HAIry WILDrye

(*Elymus innovatus*) 24 15-57 100

BLUNT SEDGE

(*Carex obtusata*) 6 3-13 67

ROUGH FESCUE

(*Festuca scabrella*) 37 8-57 100

PARRY OATGRASS

(*Danthonia parryii*) 5 0-24 57

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:

SUBMESIC

NUTRIENT REGIME :

SUBMESOTROPHIC-MESOTROPHIC

ELEVATION:

1620-2042(1859)

SOIL DRAINAGE:

RAPIDLY

SLOPE:

0-50(22)%

ASPECT:

SOUTHERLY

### FORAGE PRODUCTION KG/HA

TOTAL 1480(900-2502)

SUGGESTED GRAZING CAPACITY

0.6 HA/AUM OR 1.4 AC/AUM

## SASMA3a. Hairy wildrye- Rough fescue -*Carex* spp.

(*Elymus innovatus*- *Festuca scabrella*-*Sedge*)

n=1 This community is similar to the Rough fescue-Hairy wildrye community type (SASMA3) previously described, but this type was described on a site with a north aspect. Corns and Achuff (1982), described hairy wildrye dominated community types on south facing slopes in the more northern ecodistricts. They felt these grasslands occurred on areas with frequent snow avalanching. It would appear that this community type represents the transition between the northerly hairy wildrye dominated grasslands and the southerly rough fescue-hairy wildrye dominated grasslands.

This community type was described in an area that is difficult for livestock to access. It should likely be rated as non-use.

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:  
SUBMESIC

NUTRIENT REGIME :  
SUBMESOTROPHIC

ELEVATION:  
1951M

SOIL DRAINAGE:  
WELL

SLOPE:  
30%

ASPECT:  
NORTHERLY

### FORAGE PRODUCTION KG/HA

TOTAL 1225(900-1500)

SUGGESTED GRAZING CAPACITY  
0.7 HA/AUM OR 1.6 AC/AUM

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

#### SHRUBS

SHRUBBY CINQUEFOIL. ( <i>Potentilla fruticosa</i> )	2	-	100
COMMON JUNIPER ( <i>Juniperus communis</i> )	1	-	100

#### FORBS

YELLOW BEARDTONGUE ( <i>Penstemon confertus</i> )	4	-	100
MISSOURI GOLDENROD ( <i>Solidago missouriensis</i> )	2	-	100
FIREWEED ( <i>Epilobium angustifolium</i> )	4	-	100
ALPINE HEDYSARUM ( <i>Hedysarum alpinus</i> )	5	-	100
MOUNTAIN DANDELION ( <i>Agoseris glauca</i> )	2	-	100

#### GRASSES

HAIRY WILDRYE ( <i>Elymus innovatus</i> )	20	-	100
BLUNT SEDGE ( <i>Carex obtusata</i> )	1	-	100
ROUGH FESCUE ( <i>Festuca scabrella</i> )	11	-	100

## SASMA4. Sedge-Hairy wildrye-Slender wheatgrass

(*Carex obtusata*-*Elymus innovatus*-*Agropyron trachycaulum*)

**n=15** This community type appears to represent the result of moderate to heavy grazing pressure on a Rough fescue-Sedge dominated community type. Heavy grazing appears to cause rough fescue to decline and allows sedge to increase. Indeed Willoughby et al. (2003), described similar species composition changes with grazing on south facing slopes in the Montane subregion.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
SHRUBBY CINQUEFOIL ( <i>Potentilla fruticosa</i> )	3	0-13	55
<b>FORBS</b>			
STRAWBERRY ( <i>Fragaria virginiana</i> )	11	1-17	100
WHITE CAMUS ( <i>Zigadenus elegans</i> )	T	0-1	9
YARROW ( <i>Achillea millefolium</i> )	12	1-15	100
SMOOTH LEAVED CINQUEFOIL ( <i>Potentilla diversifolia</i> )	6	0-23	46
BEARBERRY ( <i>Arctostaphylos uva-ursi</i> )	2	0-18	18
YELLOW BEARDTONGUE ( <i>Penstemon confertus</i> )	1	0-14	36
OLD MAN'S WHISKERS ( <i>Geum triflorum</i> )	7	0-11	46
<b>GRASSES</b>			
HAIRY WILDRYE ( <i>Elymus innovatus</i> )	10	0-26	91
ROUGH FESCUE ( <i>Festuca scabrella</i> )	3	0-13	55
SLENDER WHEATGRASS ( <i>Agropyron trachycaulum</i> )	5	0-21	64
IDAHO FESCUE ( <i>Festuca idahoensis</i> )	1	0-7	64
BLUNT SEDGE ( <i>Carex obtusata</i> )	14	0-31	55

MOISTURE REGIME:  
SUBMESIC

NUTRIENT REGIME:  
SUBMESOTROPIC

ELEVATION:  
1768-1981(1823) M

SOIL DRAINAGE:  
WELL

SLOPE:  
0-35(12)%

ASPECT:  
SOUTHERLY

### FORAGE PRODUCTION KG/HA

GRASS 814 (163-1364)  
FORB 505 (97-140)  
SHRUB 514 (0-1743)  
TOTAL 1674 (1000-2318)

SUGGESTED GRAZING CAPACITY  
0.5 HA/AUM

### ENVIRONMENTAL VARIABLES

## SASMA5. Kentucky bluegrass/Dandelion

(*Poa pratensis*/*Taraxacum officinale*)

n=14 This community type represents a moist, tufted hairgrass-sedge dominated community type that has been heavily grazed. Heavy grazing pressure causes tufted hairgrass to decline and allows sedge and Kentucky bluegrass to increase. Continuous heavy grazing pressure will eventually cause all native species to decline on the site and the site will become dominated by Kentucky bluegrass, dandelion and clover (Willoughby 1992).

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
SMOOTH WILLOW ( <i>Salix glauca</i> )	T	0-4	7
BRISTLY BLACK CURRANT ( <i>Ribes lacustre</i> )	1	0-7	7
<b>FORBS</b>			
DANDELION ( <i>Taraxacum officinale</i> )	10	1-29	100
YARROW ( <i>Achillea millefolium</i> )	5	2-15	100
FIREWEED ( <i>Epilobium angustifolium</i> )	3	0-19	43
BUNCHBERRY ( <i>Cornus canadensis</i> )	1	0-7	7
CLOVER ( <i>Trifolium spp.</i> )	4	0-15	46
<b>GRASSES</b>			
KENTUCKY BLUEGRASS ( <i>Poa pratensis</i> )	43	0-58	71
RUSH LIKE SEDGE ( <i>Carex scirpoidea</i> )	2	0-26	7
TUFTED HAIRGRASS ( <i>Deschampsia cespitosum</i> )	T	0-5	7
TIMOTHY ( <i>Phleum pratense</i> )	12	0-28	71

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:  
SUBHYGRIC

NUTRIENT REGIME:  
PERMESOTROPHIC

ELEVATION:  
1541(1340-1798)M

SOIL DRAINAGE:  
MODERATELY WELL

SLOPE:  
6(0-20)%

ASPECT:  
VARIABLE

### FORAGE PRODUCTION KG/HA

GRASS 1673(248-3700)  
FORB 593(420-889)  
SHRUB 75(0-198)  
TOTAL 2341(746-4589)

SUGGESTED GRAZING CAPACITY  
0.4 HA/AUM



## SASMA6. Yellow mountain avens

(*Dryas drummondii*)

n=1 This is similar to the community type described in the Central and Northern Rocky Mountain eodistricts. Corns and Achuff (1982) described this community type on recent fluvial and glacialfluvial landforms with gentle slopes, where the soils are rapidly drained. Willoughby et al. (2003), described a yellow mountain avens community on dry, gravelly river flats with nutrient poor soils in the Montane subregion. They found this community type to be successionaly immature and succession would be to a Balsam poplar dominated community type.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
SHRUBBY CINQUEFOIL ( <i>Potentilla fruticosa</i> )	4	-	100
<b>FORBS</b>			
YELLOW DRYAD ( <i>Dryas drummondii</i> )	13	-	100
LATE YELLOW LOCOWEED ( <i>Oxytropis monticola</i> )	5	-	100
SILVERY CINQUEFOIL ( <i>Potentilla argentea</i> )	3	-	100
LOW GOLDENROD ( <i>Solidago missouriensis</i> )	3	-	100
YELLOW HEDYSARUM ( <i>Hedysarum sulphurascens</i> )	3	-	100
<b>GRASSES</b>			
HAIRY WILDRYE ( <i>Elymus innovatus</i> )	6	-	100
JUNEGRASS ( <i>Koeleria macrantha</i> )	T	-	100
SLENDER WHEATGRASS ( <i>Agropyron trachycaulum</i> )	1	-	100
BLUNT SEDGE ( <i>Carex obtusata</i> )	67	-	100
FRINGED BROME ( <i>Bromus ciliatus</i> )	5	-	100

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:

MESIC

NUTRIENT REGIME:

SUBMESOTROPHIC

ELEVATION:

2165M

SOIL DRAINAGE:

WELL TO RAPIDLY

### FORAGE PRODUCTION KG/HA

GRASS 572

FORB 602

TOTAL 1372

\*ESTIMATE

SUGGESTED GRAZING CAPACITY

NON-USE

## SASMA7. Tufted hairgrass-Sedge

(*Deschampsia cespitosa*-*Carex spp.*)

n=3 This community type was described in the Savanna Creek range allotment and is located on moist sites that are better drained and slightly drier than the pure sedge meadows. Willoughby(2001), found that tufted hairgrass is a common plant species on lowland sites in the valley bottoms of the Upper Foothills subregion . Willoughby (1992), found when this community type is protected from grazing for 25-30 years, willow and bog birch expand and tufted hairgrass and sedge decline. The decline in graminoid cover also results in a decline in available forage production . Continuous heavy grazing causes hairgrass to decline and the site will be invaded by Kentucky bluegrass and dandelion.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
BARCLAY'S WILLOW ( <i>Salix barclayi</i> )	3	0-7	33
SHRUBBY CINQUEFOIL ( <i>Potentilla fruticosa</i> )	1	0-2	33
<b>FORBS</b>			
LINDLEY'S ASTER ( <i>Aster ciliolatus</i> )	1	0-2	33
YELLOW HEDYSARUM ( <i>Hedysarum sulphurcens</i> )1		0-3	33
YARROW ( <i>Achillea millefolium</i> )	6	3-10	100
GRACEFUL CINQUEFOIL ( <i>Potentilla gracilis</i> )	9	6-14	100
OLD MAN'S WHISKERS ( <i>Geum triflorum</i> )	3	0-9	68
AMERICAN VETCH ( <i>Vicia americana</i> )	T	0-1	67
<b>GRASSES</b>			
TWO-SEEDED SEDGE ( <i>Carex disperma</i> )	14	0-42	33
TUFTED HAIRGRASS ( <i>Deschampsia cespitosa</i> )	41	18-62	100
SLENDER WHEATGRASS ( <i>Agropyron trachycaulum</i> )2		1-4	100
ROUGH FESCUE ( <i>Festuca scabrella</i> )	2	0-6	33
HAIRY WILD RYE ( <i>Elymus innovatus</i> )	6	0-18	33

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:  
SUBHYGRIC

NUTRIENT REGIME:  
PERMESOTROPHIC

ELEVATION:  
1445(1445-1450)M

SOIL DRAINAGE: MOD. WELL

SLPOE 4%

ASPECT:NORTHERLY

### FORAGE PRODUCTION KG/HA

GRASS	1164
FORBS	162
SHRUBS	0
TOTAL	1326

SUGGESTED GRAZING CAPACITY  
0.7 HA/AUM OR 2.5 AC/AUM

**SASMA7a. Marsh reedgrass/Cow parsnip**  
(*Calamagrostis canadensis/Heracleum lanatum.*)

**n=1** This community type represents moist seepage areas in the foothills west of Turner valley. Often these sites are willow dominated to form the Willow/Marsh reedgrass dominated community type. Marsh reedgrass is characteristic of the lower elevation Lower Foothills subregion. Moving up in elevation into the Subalpine and Upper Foothills subregions, there is often a shift in dominance away from marsh reedgrass to tufted hairgrass on these moist sites. This community type is very similar to the Cow parsnip/Veiny meadow rue community described by Lane et al. (2000) in the Lower Foothills subregion. In the Lower Foothills this community type is very productive and is often heavily utilized by livestock. The Marsh reedgrass-Timothy/Cow parsnip (SASMC8) represents a grazing disclimax of this community type.

**PLANT COMPOSITION CANOPY COVER(%)**

MEAN RANGE CONST.

**SHRUBS**

PRICKLY ROSE ( <i>Rosa acicularis</i> )	4	-	100
RASPBERRY ( <i>Rubus idaeus</i> )	1	-	100

**FORBS**

COW PARSNIP ( <i>Heracleum lanatum</i> )	3	-	100
FIREWEED ( <i>Epilobium angustifolium</i> )	25	-	100
WESTERN MEADOW RUE ( <i>Thalictrum occidentale</i> )	10	-	100
HORSETAIL ( <i>Equisetum arvense</i> )	4	-	100
STICKY PURPLE GERANIUM ( <i>Geranium viscosissimum</i> )	4	-	100
AMERICAN VETCH ( <i>Vicia americana</i> )	2	-	100

**GRASSES**

SEDGE ( <i>Carex spp</i> )	15	-	100
MARSH REEDGRASS ( <i>Calamagrostis canadensis</i> )	24	-	100
SLENDER WHEATGRASS ( <i>Agropyron trachycaulum</i> )	1	-	100
IDAHO FESCUE ( <i>Festuca idahoensis</i> )	5	-	100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:

SUBHYGRIC

NUTRIENT REGIME:

PERMESOTROPHIC

ELEVATION:

1768M

SOIL DRAINAGE: MOD. WELL

SLOPE 3%

ASPECT: SOUTHERLY

**FORAGE PRODUCTION KG/HA**

TOTAL 2000\* ESTIMATE

<p>SUGGESTED GRAZING CAPACITY 0.5 HA/AUM</p>
--

## SASMA8. Rough fescue-Idaho fescue-Parry oatgrass

(*Festuca scabrella*-*Festuca idahoensis*-*Danthonia parryii*)

n=4 This community was described on lower slope positions in the Castle area. It is very similar to the Rough fescue-Idaho fescue-Parry oatgrass community described on Black Chernozemic soils in the Montane subregion from an elevation of 1300m up to 1900m (Willoughby et al. 2003) Willoughby (1992), described one Rough fescue-dominated site where the species composition had not changed in over 30 years, indicating this maybe the climax community type on river terraces and south facing slopes in the Montane subregion. Indeed Moss and Campbell (1947), found that rough fescue grows almost to the exclusion of other plants in the absence of disturbance. On rocky and gravelly slopes with shallow soils, rough fescue is replaced by Parry oatgrass and Idaho fescue. They also found Parry oatgrass and Idaho fescue increased and rough fescue declined with increased grazing pressure. Willoughby (1992), also described rough fescue and Idaho fescue dominated community types with little Parry oatgrass in the Castle area south of Blairmore. He also found that rose and shrubby cinquefoil tended to increase in cover at higher elevations in these grasslands.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
SHRUBBY CINQUEFOIL ( <i>Potentilla fruticosa</i> )	4	1-10	100
PRICKLY ROSE ( <i>Rosa acicularis</i> )	3	0-8	75
<b>FORBS</b>			
SHOWY ASTER ( <i>Aster conspicuus</i> )	8	0-29	75
COMMON FIREWEED ( <i>Epilobium angustifolium</i> )	4	0-6	75
CUT-LEAVED ANEMONE ( <i>Anemone multifida</i> )	2	1-2	100
GRACEFUL CINQUEFOIL ( <i>Potentilla gracilis</i> )	1	0-3	75
LONG-FRUITED WILD PARSLEY ( <i>Lomatium macrocarpum</i> )	1	0-4	25
LOW GOLDENROD ( <i>Solidago missouriensis</i> )	2	0-5	50
<b>GRASSES</b>			
ROUGH FESCUE ( <i>Festuca scabrella</i> )	21	12-27	100
IDAHO FESCUE ( <i>Festuca idahoensis</i> )	7	0-13	75
PARRY OATGRASS ( <i>Danthonia parryi</i> )	5	0-12	50
PINE REED GRASS ( <i>Calamagrostis rubescens</i> )	4	0-15	50
HAIRY WILD RYE ( <i>Elymus innovatus</i> )	1	0-1	50

### ENVIRONMENTAL VARIABLES

#### MOISTURE REGIME:

SUBXERIC TO SUBMESIC

#### NUTRIENT REGIME:

SUBMESOTROPHIC TO MESOTROPHIC

#### ELEVATION:

1838(1680-1974) M

#### SOIL DRAINAGE:

VERY RAPIDLY TO WELL

SLOPE: 13(6-22)%

ASPECT: VARIABLE

### FORAGE PRODUCTION KG/HA

GRASS 1146(934-1358)

FORBS 614(488-740)

SHRUBS 55(0-109)

TOTAL 1815(1674-1955)

### SUGGESTED GRAZING CAPACITY

0.5 HA/AUM

**SASMA9. Rough fescue-Sedge/Bearberry**  
*(Festuca scabrella-Carex obtusaia./Arctostaphylos uva-ursi)*

n=7 This community is characteristic of dry, south facing slopes, below the crest of the hill. It appears that snow accumulates in these areas favoring the growth of rough fescue. This community is similar to the Rough fescue-Sedge/Bearberry community type described by Willoughby et al. (2003) on hilltops in the Montane subregion. The shallow poorly developed soils appear to favour rough fescue, slender wheatgrass, and sedge over Parry oatgrass. This community is drier than the Rough fescue-Idaho fescue-Parry oatgrass grasslands characteristic of lower slope positions. This community type is similar to the Sedge/Bearberry community type but lacks the high cover of rough fescue. It appears that the Sedge/Bearberry community may represent a grazing disclimax of this community type.

**PLANT COMPOSITION CANOPY COVER(%)**

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
SHRUBBY CINQUEFOIL <i>(Potentilla fruticosa)</i>	5	1-11	100
PRICKLY ROSE <i>(Rosa acicularis)</i>	3	0-6	86
<b>FORBS</b>			
BEARBERRY <i>(Arctostaphylos uva-ursi)</i> 40		0-67	86
SILKY PERENNIAL LUPIN <i>(Lupinus sericeus)</i>	4	0-10	71
CUT-LEAVED ANEMONE <i>(Anemone multifida)</i>	3	0-2	71
WILD STRAWBERRY <i>(Fragaria virginiana)</i>	2	0-9	43
SMOOTH ASTER <i>(Aster laevis)</i>	1	0-8	43
LOW GOLDENROD <i>(Solidago missouriensis)</i>	1	0-6	71
<b>GRASSES</b>			
ROUGH FESCUE <i>(Festuca scabrella)</i>	22	6-43	100
SEDGES <i>(Carex spp.)</i>	2	0-5	71
PARRYS OATGRASS <i>(Danthonia Perryi)</i>	4	0-18	43
PINE REED GRASS <i>(Calamagrostis rubescens)</i> 4		0-15	50
HAIRY WILDRYE <i>(Elymus innovatus)</i>	3	0-9	57

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
XERIC TO MESIC

NUTRIENT REGIME:  
OLIGOTROPHIC TO MESOTROPHIC

ELEVATION:  
1740(1400-1939)M

SOIL DRAINAGE:  
VERY RAPIDLY-WELL

SLOPE:  
30(18-40)%

ASPECT:  
VARIABLE

**FORAGE PRODUCTION KG/HA**

GRASS 725(102-1612)  
 FORB 385(111-676)  
 SHRUB 1475(28-4400)  
 TOTAL 2585(970-5384)

SUGGESTED GRAZING CAPACITY  
0.3 HA/AUM

## SASMA10. Parry oatgrass-Rough fescue-Sedge

(*Danthonia parryi*-*Festuca scabrella*-*Carex spp*)

n=12 This community is situated upslope of the rough fescue dominated community types. This community type tends to be drier and better drained than the rough fescue dominated types found in the lower slope positions. The drier site conditions favour the growth of Parry oatgrass and sedge over rough fescue.

These sites are moderately productive and are easily accessible to livestock. They should be considered primary range. Continuous heavy grazing pressure will lead to a community type that is dominated by sedge and fringed sage.

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

#### SHRUBS

SHRUBBY CINQUEFOIL ( <i>Potentilla fruticosa</i> )	4	0-10	92
PRICKLY ROSE ( <i>Rosa acicularis</i> )	3	0-5	75
SASKATOON ( <i>Amelanchier alnifolia</i> )	2	0-16	50

#### FORBS

SILKY PERENNIAL LUPINE ( <i>Lupinus sericeus</i> )	5	0-18	75
CUT-LEAVED ANEMONE ( <i>Anemone multifida</i> )	2	0-2	83
OLD MAN'S WHISKERS ( <i>Geum triflorum</i> )	2	0-21	42
NORTHERN BEDSTRAW ( <i>Galium boreale</i> )	4	1-8	100
EARLY YELLOW LOCOWEED ( <i>Oxytropis sericea</i> )	2	0-6	67
GOLDENBEAN ( <i>Thermopsis rhombifolia</i> )	4	0-8	42

#### GRASSES

ROUGH FESCUE ( <i>Festuca scabrella</i> )	6	0-15	92
PARRY OATGRASS ( <i>Danthonia parryi</i> )	23	10-31	100
SEDGES ( <i>Carex spp.</i> )	3	0-8	94
JUNE GRASS ( <i>Koeleria macrantha</i> )	3	0-6	92
COLUMBIA NEEDLE GRASS ( <i>Stipa columbiana</i> )	5	0-13	75
SLENDER WHEATGRASS ( <i>Agropyron trachycaulum</i> )	1	0-5	75

### ENVIRONMENTAL VARIABLES

#### MOISTURE REGIME:

XERIC TO SUBXERIC

#### NUTRIENT REGIME :

OLIGOTROPHIC TO PERMESOTROPHIC

#### ELEVATION:

1473(1400-1962) M

#### SOIL DRAINAGE:

RAPIDLY TO WELL

#### SLOPE:

23(12-33) %

#### ASPECT:

VARIABLE

### FORAGE PRODUCTION KG/HA

GRASS 732(356-1284)

FORB 702(0-1550)

SHRUB 96(0-388)

TOTAL 1452(660-2866)

### SUGGESTED GRAZING CAPACITY

0.6 HA/AUM

## SASMA11. Sedge/Bearberry

(*Carex spp./Arctostaphylos uva-ursi*)

**n=11** This community was described on south facing slopes. This community is very similar to the Rough fescue-Sedge/Bearberry community type, but lacks the high cover of rough fescue. A number of the sites described in this community had extensive grazing pressure by bighorn sheep and domestic livestock. It is possible that this community type represents a grazing disclimax of the Rough fescue-Sedge/Bearberry community type.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>TREES</b>			
SUBALPINE FIR			
( <i>Abies lasiocarpa</i> )	T	0-2	20
WHITE BARK PINE			
( <i>Pinus albicaulis</i> )	T	0-3	20
<b>SHRUBS</b>			
BEARBERRY			
( <i>Arctostaphylos uva-ursi</i> )	27	6-70	100
SHRUBBY CINQUEFOIL			
( <i>Potentilla fruticosa</i> )	2	0-8	70
SASKATOON			
( <i>Amelanchier alnifolia</i> )	4	0-10	50
<b>FORBS</b>			
EARLY YELLOW LOCOWEED			
( <i>Oxytropis sericea</i> )	2	0-5	90
BROWN-BRACTED MOUNTAIN EVERLASTING			
( <i>Antennaria umbrinella</i> )	1	0-5	50
YELLOW VALSE DANDELION			
( <i>Agoseris glauca</i> )	1	0-3	70
ARCTIC SANDWART			
( <i>Minuartia obtusiloba</i> )	1	0-3	40
YELLOW HEDYSARUM			
( <i>Hedysarum sulphurescens</i> )	2	0-6	60
COMMON FIREWEED			
( <i>Epilobium angustifolium</i> )	1	0-3	30
<b>GRASSES</b>			
SEDGES			
( <i>Carex spp.</i> )	1	0-4	80
ROUGH FESCUE			
( <i>Festuca scabrella</i> )	2	0-7	70
IDAHO FESCUE			
( <i>Festuca idahoensis</i> )	1	0-2	80
BLUEBUNCH WHEATGRASS			
( <i>Agropyron spicatum</i> )	3	0-12	90

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:  
XERIC-SUBXERIC

NUTRIENT REGIME:  
OLIGOTROPHIC TO PERMESOTROPHIC

ELEVATION:  
1883(1706-2072)M

SOIL DRAINAGE:  
VERY RAPIDLY TO WELL

SLOPE: 31(16-50)%

ASPECT: SOUTH TO WEST

### FORAGE PRODUCTION KG/HA

GRASS 733(17-1238)  
FORB 391(34-583)  
SHRUB 1051(241-2382)  
TOTAL 2175(605-4121)

SUGGESTED GRAZING CAPACITY  
NON-USE

**SASMA12. Silverberry-Rose**  
(*Elaeagnus commutata*-*Rosa acicularis*)

n=1 This community type was described on the downwind side of the hill at lower elevations in the Castle area of the province (north of Waterton Lakes National Park). It appears this area accumulates snow so the moisture regime is favorable for the growth of silverberry and aspen. At 1800 m aspen is at its upper elevational limit. The aspen trees at this site are very stunted and lack the vigour of lower elevation sites. Silverberry is well adapted to growing on dry, gravelly, light soils in ravines, coulees and stream banks throughout Alberta (Wilkinson 1990). It is unusual to have silverberry and aspen growing at these higher elevations indicating that this site is somewhat protected and warmer so that the climate resembles the lower elevation sites.

**PLANT COMPOSITION CANOPY COVER(%)**

MEAN RANGE CONST.

**TREES**

ASPEN  
(*Populus tremuloides*) 4 - 100

**SHRUBS**

SILVERBERRY  
(*Elaeagnus commutata*) 15 - 100

PRICKLY ROSE  
(*Rosa acicularis*) 5 - 100

ASPEN  
(*Populus tremuloides*) 4 - 100

**FORBS**

MOUSE EARED CHICKWEED  
(*Cerastium arvense*) 6 - 100

FALSE DANDELION  
(*Agoseris glauca*) 2 - 100

SILVER PLANT  
(*Eriogonum ovalifolium*) T - 100

**GRASSES**

ROCKY MOUNTAIN FESCUE  
(*Festuca brachycaulum*) 1 - 100

BLUEBUNCH WHEATGRASS  
(*Agropyron spicatum*) 3 - 100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:

XERIC

NUTRIENT REGIME:

SUBMESOTROPHIC

ELEVATION: 1841 M

SOIL DRAINAGE: VERY RAPIDLY

SLOPE: 24%

ASPECT: SOUTHEASTERLY

**FORAGE PRODUCTION KG/HA**

TOTAL 500 \*ESTIMATE

SUGGESTED GRAZING CAPACITY  
1.8 HA/AUM



## SASMA13. Fescue-Junegrass/Early yellow locoweed

(*Festuca spp.-Koeleria macrantha/Oxytropis sericea*)

**n=17** This community type is characteristic of the dry, rocky, windswept ridges in the Castle area north of Waterton Lakes National Park. At higher elevations this community is often replaced by the White mountain avens community which occupies similar sites. This community type is somewhat variable false mountain dandelion and early yellow locoweed were found in the majority of the plots, but in areas where there is slightly more snow accumulation the cover of Idaho fescue increases, and there are larger patches of species like spotted saxifrage, brown bracted mountain everlasting and sandwort. Bareground on these moister areas is usually less than 40%. In contrast bareground on the drier parts of this community type exceed 60%. At lower elevations on Whistler Mtn. bluebunch wheatgrass becomes more prevalent.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
SHRUBBY CINQUEFOIL ( <i>Potentilla fruticosa</i> )	3	0-13	82
LIMBER PINE ( <i>Pinus flexilis</i> )	1	0-11	12
<b>FORBS</b>			
BEARBERRY ( <i>Arctostaphylos uva-ursi</i> )	1	0-8	24
KITTENTAILS ( <i>Besseyia wyomingensis</i> )	1	0-3	59
EARLY YELLOW LOCOWEED ( <i>Oxytropis sericea</i> )	1	0-5	88
SANDWORT ( <i>Minuartia obtusiloba</i> )	3	0-16	47
LITTLE CLUBMOSS ( <i>Selaginella densa</i> )	2	0-8	41
SPOTTED SAXIFRAGE ( <i>Saxifraga bronchialis</i> )	3	0-29	24
<b>GRASSES</b>			
ROUGH FESCUE ( <i>Festuca scabrella</i> )	1	0-6	24
JUNEGRASS ( <i>Koeleria macrantha</i> )	2	0-5	77
SEDGE ( <i>Carex spp.</i> )	3	0-9	82
IDAHO FESCUE ( <i>Festuca idahoensis</i> )	3	0-20	41
SMOOTH BROME ( <i>Bromus pumpellianus</i> )	1	0-5	53
BLUEBUNCH WHEATGRASS ( <i>Agropyron spicatum</i> )	1	0-3	41

### ENVIRONMENTAL VARIABLES

#### MOISTURE REGIME:

VERY XERIC TO SUBXERIC

#### NUTRIENT REGIME:

OLIGOTROPHIC TO MESOTROPHIC

#### ELEVATION:

1949(1536-2322) M

#### SOIL DRAINAGE:

VERY RAPIDLY TO WELL

#### SLOPE: 26(6-50)%

#### ASPECT: VARIABLE

### FORAGE PRODUCTION KG/HA

GRASS 512(18-2018)

FORB 291(44-680)

SHRUB 192(0-589)

TOTAL 994(519-2126)

SUGGESTED GRAZING CAPACITY

NON-USE

**SASMA14. White mountain avens**  
(*Dryas octopetala*)

**n=12** This community occurs on wind-exposed, snow free ridges and resembles the White mountain avens community described in the Alpine subregion. The soils are shallow, stoney, colluvial Regosols (Corns and Achuff 1982). Ogilvie (1969), found this community type to have an abundance of cushion and mat plants and a large number of lichens. This community is generally found at higher elevations than the Fescue-Junegrass/E. yellow locoweed community type. This community type appears to have no snow accumulation throughout the year, whereas, the fescue, Junegrass dominated community appears to have some snow accumulation. This may account for the differences in dominant plant species for each community type.

**PLANT COMPOSITION CANOPY COVER(%)**  
MEAN RANGE CONST.

**SHRUBS**

PRICKLY ROSE ( <i>Rosa acicularis</i> )	T	0	25
SHRUBBY CINQUEFOIL ( <i>Potentilla fruticosa</i> )	2	0-6	92
BUFFALOBERRY ( <i>Shepherdia canadensis</i> )	T	0-3	8
SILVERBERRY ( <i>Elaeagnus commutata</i> )	T	0-2	8

**FORBS**

WHITE MOUNTAIN AVENS ( <i>Dryas octopetala</i> )	31	14-48	100
EARLY YELLOW LOCOWEED ( <i>Oxytropis sericea</i> )	3	0-7	100
SPOTTED SAXIFRAGE ( <i>Saxifraga bronchialis</i> )	2	0-7	42
KITTENTAILS ( <i>Besseyia wyomingensis</i> )	1	0-2	83
YELLOW HEDYSARUM ( <i>Hedysarum sulphureum</i> )	2	0-4	67
FALSE MTN. DANDELION ( <i>Agoseris glauca</i> )	1	0-2	83

**GRASSES**

IDAHO FESCUE ( <i>Festuca idahoensis</i> )	1	0-4	33
SEDGE SPP. ( <i>Carex spp.</i> )	3	0-9	83
SMOOTH BROME ( <i>Bromus pumellianus</i> )	1	0-2	67
JUNEGRASS ( <i>Koeleria macrantha</i> )	1	0-5	67

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
VERY XERIC-SUBXERIC

NUTRIENT REGIME:  
OLIGOTROPHIC TO SUBMESOTROPHIC

ELEVATION:  
2136(2001-2423) M

SOIL DRAINAGE:  
VERY RAPIDLY TO RAPIDLY

SLOPE : 17(7-38)%

ASPECT: VARIABLE

**FORAGE PRODUCTION KG/HA**

GRASS 36(25-47)  
FORB 198(27-369)  
SHRUB 392(195-589)  
TOTAL 626(269-983)

SUGGESTED GRAZING CAPACITY  
NON-USE

**SASMA15. Pinegrass-Hairy wildrye/Strawberry**  
(*Calamagrostis rubescens*-*Elymus innovatus*/*Fragaria virginiana*)

n=4 This community type is similar to the pinegrass dominated community type described on west and north facing slopes in the Montane subregion (Willoughby et al 2003). In the Montane this community represents the transition from grassland to forest on moist sites with northerly aspects. There is usually high forb cover on these sites with strawberry, showy aster, american vetch and silky perennial lupine being common. Pinegrass and Hairy wildrye are common grass species in the understory of conifer and deciduous stands and their dominance in this community type may indicate a transition to a forested community. The higher moisture conditions on these sites allows for production of over 2000 kg/ha.

**PLANT COMPOSITION CANOPY COVER(%)**

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
PRICKLY ROSE ( <i>Rosa acicularis</i> )	9	1-20	100
SHRUBBY CINQUEFOIL ( <i>Potentilla fruticosa</i> )	1	1-2	100
WHITE MEADOWSWEET ( <i>Spiraea betulifolia</i> )	1	0-4	25
ASPEN ( <i>Populus tremuloides</i> )	1	0-3	50
<b>FORBS</b>			
SILKY PERENNIAL LUPINE ( <i>Lupinus sericeus</i> )	4	1-7	100
SHOWY ASTER ( <i>Aster conspicuus</i> )	3	1-6	100
COMMON DANDELION ( <i>Taraxacum officinale</i> )	3	1-8	100
CREAM-COLORED VETCHILING ( <i>Lathyrus ochroleucus</i> )	2	1-3	100
GRACEFUL CINQUEFOIL ( <i>Potentilla gracilis</i> )	1	1-2	100
FALSE MTN. DANDELION ( <i>Agoseris glauca</i> )	1	0	100
<b>GRASSES</b>			
PINEGRASS ( <i>Calamagrostis rubescens</i> )	11	1-22	100
JUNEGRASS ( <i>Koeleria macrantha</i> )	2	1-2	100
HAIRY WILDRYE ( <i>Elymus innovatus</i> )	4	0-11	75
KENTUCKY BLUEGRASS ( <i>Poa pratensis</i> )	3	0-13	25

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
SUBXERIC TO MESIC

NUTRIENT REGIME:  
SUBMESOTROPHIC TO MESOTROPHIC

ELEVATION:  
1701(1684-1710)M

SOIL DRAINAGE:  
WELL TO IMPERFECTLY

SLOPE (RANGE):  
31(23-40)%

ASPECT:  
SOUTHERLY

**FORAGE PRODUCTION KG/HA**

GRASS 758(392-1204)  
FORB 1170(506-1884)  
SHRUB 110(0-260)  
TOTAL 2037(1346-2739)

SUGGESTED GRAZING CAPACITY  
0.4 HA/AUM

**SASMA16. Forb meadow**  
(*Epilobium angustifolium*)

n=3 This site is similar to the Fireweed-Meadow rue/Sedge dominated community type described in the northern foothills area of the Subalpine. This community is characterized by a dominance of forb species (fireweed, graceful cinquefoil, yellow hedysarum) and only a small cover of grass species. It would appear that the moisture and nutrient regime are higher on this site compared to the other grassland community types which favors the growth of forb species. Succession in the absence of disturbance will likely be to aspen and then white spruce.

The forage production of this community type is fairly high because of the higher moisture and nutrient content of the soil, but the areas are so small and isolated they contribute little to the overall carrying capacity of a disposition.

**PLANT COMPOSITION CANOPY COVER(%)**  
MEAN RANGE CONST.

**SHRUBS**

PRICKLY ROSE ( <i>Rosa acicularis</i> )	2	0-2	100
SHRUBBY CINQUEFOIL ( <i>Potentilla fruticosa</i> )	3	0-5	100
CREEPING JUNIPER ( <i>Juniperus horizontalis</i> )	1	0-2	67

**FORBS**

GRACEFUL CINQUEFOIL ( <i>Potentilla gracilis</i> )	3	1-6	100
COMMON FIREWEED ( <i>Epilobium angustifolium</i> )	6	2-10	100
YELLOWHEDYSARUM ( <i>Hedysarum sulphurescens</i> )	4	0-9	100
WILD VETCH ( <i>Vicia americana</i> )	2	0-3	100
WILD STRAWBERRY ( <i>Fragaria virginiana</i> )	1	0-1	100
CUT-LEAVED ANEMONE ( <i>Anemone multifida</i> )	1	0-2	100

**GRASSES**

HAIRY WILDRYE ( <i>Elymus innovatus</i> )	3	0-5	100
PARRYS OATGRASS ( <i>Danthonia parryi</i> )	2	0-3	100
TIMOTHY ( <i>Phleum pratense</i> )	1	0-1	67
ROUGH FESCUE ( <i>Festuca scabrella</i> )	2	0-4	33

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
MESIC

NUTRIENT REGIME:  
PERMESOTROPHIC TO MESOTROPHIC

ELEVATION: 1701(1684-1710)

SOIL DRAINAGE:  
WELL TO IMPERFECTLY

SLOPE: 3(0-7)%

ASPECT: VARIABLE

**FORAGE PRODUCTION KG/HA**

GRASS 554(175-1126)  
FORB 734(567-1009)  
SHRUB 125(0-208)  
TOTAL 1413(968-2135)

SUGGESTED GRAZING CAPACITY  
0.6 HA/AUM

## SASMB1. Willow/Sedge

(*Salix spp./Carex spp.*)

n=2 Willow encroachment into moist grassland meadows eventually results in this community type. Historically fire has played an important role in the maintenance of the grassland community type in this subregion. Continued fire suppression will eventually allow willow and bog birch to invade many of the grassy meadows.

This community type is slightly drier than the Willow/Water sedge dominated community type.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
WILLOW			
( <i>Salix spp.</i> )	30	27-32	100
BOG BIRCH			
( <i>Betula glandulosa</i> )	4	2-6	100
SHRUBBY CINQUEFOIL			
( <i>Potentilla fruticosa</i> )	3	0-5	100
<b>FORBS</b>			
YARROW			
( <i>Achillea millefolium</i> )	1	0-1	50
WILD STRAWBERRY			
( <i>Fragaria virginiana</i> )	7	0-13	50
SMOOTH ASTER			
( <i>Aster laevis</i> )	3	0-6	50
SILVERY CINQUEFOIL			
( <i>Potentilla argentea</i> )	3	0-6	50
LOW GOLDENROD			
( <i>Solidago missouriensis</i> )	8	0-15	50
<b>GRASSES</b>			
TUFTED HAIRGRASS			
( <i>Deschampsia cespitosa</i> )	3	3	100
BALTIC RUSH			
( <i>Juncus balticus</i> )	7	0-13	100
SEDGE			
( <i>Carex spp.</i> )	42	35-49	100
HAIRY WILD RYE			
( <i>Elymus innovatus</i> )	3	0-6	50

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:

MESIC -HYGRIC

NUTRIENT REGIME :

OLIGOTROPIC TO PERMESOTROPIC

ELEVATION:

1599(1300-1631) M

SOIL DRAINAGE :

WELL TO VERY POORLY

### FORAGE PRODUCTION KG/HA

GRASS 1695(1622-1768)

FORB 373(0-1120)

SHRUB 149(0-446)

TOTAL 2478(2068-2888)

SUGGESTED GRAZING CAPACITY

NON-USE

## SASMB2. Willow/Richardson needlegrass

(*Salix spp./Stipa richardsonii*)

n=2 The ecology of this community type is unclear. Stringer (1973) described a Richardson needlegrass shrub savanna on small isolated areas, south facing slopes amongst subalpine fir, spruce, douglas fir forests in Banff and Jasper National Parks. He felt these grassland types were not closely related to any other grassland types.

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

#### SHRUBS

WILLOW SPP.

(*Salix myrtillifolia*) 30 20-40 100

BOG BIRCH

(*Betula glandulosa*) 5 0-10 50

SHRUBBY CINQUEFOIL

(*Potentilla fruticosa*) 5 0-10 50

#### FORBS

SHOWY LOCOWEED

(*Oxytropis splendens*) 1 1 100

YARROW

(*Achillea millefolium*) 1 1 100

FIREWEED

(*Epilobium angustifolium*) 1 1 100

STRAWBERRY

(*Fragaria virginiana*) 2 1-2 100

#### GRASSES

RICHARDSON NEEDLEGRASS

(*Stipa richardsonii*) 45 40-50 100

ROCKY MTN. FESCUE

(*Festuca saximontana*) 8 5-10 100

SEDGE

(*Carex spp*) 3 1-5 100

QUACK GRASS

(*Agropyron repens*) 1 0-2 50

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:

SUBMESIC

NUTRIENT REGIME:

MESOTROPHIC

ELEVATION:

1375(1300-1450) M

SOIL DRAINAGE:

WELL

### FORAGE CAPACITY KG/HA

SUGGESTED GRAZING CAPACITY

NON-USE

## SASMB3. Whitebark pine

(*Pinus albicaulis*)

n=1 This is a timberline community type found on steep south facing slopes with subxeric moisture regimes. The trees tend to be very small and shrub like. On moister sites the tree islands would be made up of the subalpine fir community type previously described.

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

#### TREES

WHITE BARK PINE ( <i>Pinus albicaulis</i> )	25	-	100
--	----	---	-----

#### SHRUBS

SMOOTH WILLOW ( <i>Salix glauca</i> )	8	-	100
BUFFALOBERRY ( <i>Shepherdia canadensis</i> )	8	-	100
GROUND JUNIPER ( <i>Juniperus communis</i> )	3	-	100

#### FORBS

BEARBERRY ( <i>Arctostaphylos uva-ursi</i> )	2	-	100
MOUNTAIN GOLDENROD ( <i>Solidago spathulata</i> )	2	-	100
STRAWBERRY ( <i>Fragaria virginiana</i> )	2	-	100
YARROW ( <i>Achillea millefolium</i> )	1	-	100

#### GRASSES

SMOOTH BROME ( <i>Bromus inermis</i> )	2	-	100
SEDGE ( <i>Carex spp.</i> )	1	-	100
HAIRY WILDRYE ( <i>Elymus innovatus</i> )	1	-	100

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:  
SUBXERIC

NUTRIENT REGIME :  
MESOTROPHIC

ELEVATION:  
2030 M

SOIL DRAINAGE:  
RAPIDLY

SLOPE:  
60%

ASPECT:  
SOUTHERLY

### FORAGE CAPACITY KG/HA

SUGGESTED GRAZING CAPACITY NON-USE
---------------------------------------

## SASMB4. Willow-Bog birch/Rough fescue-Kentucky bluegrass

(*Salix spp.-Betula glandulosa\Festuca scabrella-Poa pratensis*)

n=1 This community type represents the inside transect of a rangeland reference area which has been protected from grazing for over 30 years. This community type represents moist meadows and grasslands in the Sheep area west of Turner valley. Continuous heavy grazing pressure will cause the cover of shrubs and rough fescue to decline and allow Kentucky bluegrass, timothy and dandelion to invade onto the site.

This area of the province is classified as Lower Foothills (Alberta Environmental Protection 1994), but the plant species of this area are more characteristic of the Parkland and Montane subregions and the higher Subalpine subregion than the Lower Foothills. As a result this community type was placed within the Subalpine subregion guide.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
PUSSY WILLOW ( <i>Salix discolor</i> )	14	-	100
BOG BIRCH ( <i>Betula glandulosa</i> )	11	-	100
SHRUBBY CINQUEFOIL ( <i>Potentilla fruticosa</i> )	5	-	100
<b>FORBS</b>			
LINDLEY'S ASTER ( <i>Aster ciliolatus</i> )	7	-	100
WOOLLY CINQUEFOIL ( <i>Potentilla hippiana</i> )	6	-	100
STRAWBERRY ( <i>Fragaria virginiana</i> )	2	-	100
YARROW ( <i>Achillea millefolium</i> )	3	-	100
FIREWEED ( <i>Epilobium angustifolium</i> )	3	-	100
VEINY MEADOW RUE ( <i>Thalictrum venulosum</i> )	3	-	100
<b>GRASSES</b>			
SMOOTH BROME ( <i>Bromus inermis</i> )	2	-	100
SEDGE ( <i>Carex spp.</i> )	1	-	100
HAIRY WILDRYE ( <i>Elymus innovatus</i> )	1	-	100

MOISTURE REGIME:

SUBHYGRIC

NUTRIENT REGIME :

PERMESOTROPHIC

ELEVATION:

1450M

SOIL DRAINAGE:

MOD. WELL

SLOPE: 2%

ASPECT: SOUTHERLY

### FORAGE PRODUCTION KG/HA

GRASS	600
FORB	200
SHRUB	150
TOTAL	950* ESTIMATE

SUGGESTED GRAZING CAPACITY  
1.1 HA/AUM OR 2.5 AC/AUM

### ENVIRONMENTAL VARIABLES



## SASMB5. Willow/Marsh reedgrass

(*Salix spp./Calamagrostis canadensis*)

n=2 This community type was described in the Pekisko and Deep Creek allotments which are southwest of Longview. This community represents a small pocket of willow in depressional and seepage areas. It is unusual having a community type dominated by marsh reedgrass in the Subalpine subregion. Marsh reedgrass is more characteristic of wetland sites in the Boreal forest and Lower Foothills subregions of North and Central Alberta. Perhaps this community represents a transition between the Subalpine and lower elevation Montane subregion.

These sites can be highly productive because of the increased moisture and nutrients at the site, but livestock will rarely use these communities and they should be considered non-use.

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

#### SHRUBS

WILLOW			
( <i>Salix spp.</i> )	19	13-25	100
BOG BIRCH			
( <i>Betula glandulosa</i> )	2	0-3	50
ENGELMANN SPRUCE			
( <i>Picea engelmannii</i> )	8	0-15	50

#### FORBS

ARROW-LEAVED COLTSFOOT			
( <i>Petasites sagittatus</i> )	4	0-8	50
COMMON HORSETAIL			
( <i>Equisetum arvense</i> )	7	4-10	100
COW PARSNIP			
( <i>Heraclium lanatum</i> )	14	3-25	100
FIREWEED			
( <i>Epilobium angustifolium</i> )	8	2-13	100
TALL LARKSPUR			
( <i>Delphinium glaucum</i> )	4	1-7	100

#### GRASSES

MARSH REEDGRASS			
( <i>Calamagrostis canadensis</i> )	20	15-25	100
TIMOTHY			
( <i>Phleum pratense</i> )	2	1-3	100
SWEETGRASS			
( <i>Hierochloe odorata</i> )	1	0-1	50

### ENVIRONMENTAL VARIABLES

#### MOISTURE REGIME :

SUBHYGRIC-HYGRIC

#### NUTRIENT REGIME:

PERMESOTROPHIC-EUTROPHIC

#### ELEVATION:

1704(1671-1737)M

#### SOIL DRAINAGE:

IMPERFECTLY

#### SLOPE:

0%

#### ASPECT:

NORTH

### FORAGE PRODUCTION KG/HA

TOTAL 3500 \*ESTIMATE

SUGGESTED GRAZING CAPACITY

NON-USE

## SASMC1. Parry oatgrass-Rough fescue-Kentucky bluegrass

(*Danthonia parryi*-*Festuca scabrella*-*Poa pratensis*)

**n=7** This community type represents the grazing disclimax community of the Rough fescue-Idaho fescue-Parry oatgrass community type. Increased grazing pressure favours the growth of Parry oatgrass and sedge over rough fescue. Continued heavy grazing pressure eventually leads to a decline in all native species and the site is often dominated by only Kentucky bluegrass, timothy and dandelion. Recovery of this community type back to a rough fescue dominated site is possible with a reduction in grazing pressure, but once Kentucky bluegrass establishes in the stand it will likely remain as a co-dominant.

This community type is very productive and should be considered primary range.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
PRICKLY ROSE ( <i>Rosa acicularis</i> )	1	0-1	57
SASKATOON ( <i>Amelanchier alnifolia</i> )	1	0-1	14
SHRUBBY CINQUEFOIL ( <i>Potentilla fruticosa</i> )	4	0-9	86
<b>FORBS</b>			
YARROW ( <i>Achillea millefolium</i> )	3	1-10	100
WILD STRAWBERRY ( <i>Fragaria virginiana</i> )	1	0-6	72
THREE FLOWERED AVENS ( <i>Geum triflorum</i> )	10	0-21	86
GRACEFUL CINQUEFOIL ( <i>Potentilla gracilis</i> )	3	1-11	100
DANDELION ( <i>Taraxacum officinale</i> )	4	1-9	100
AMERICAN VETCH ( <i>Vicia americana</i> )	4	0-13	86
<b>GRASSES</b>			
PARRY OATGRASS ( <i>Danthonia parryi</i> )	17	6-25	100
ROUGH FESCUE ( <i>Festuca scabrella</i> )	7	1-13	100
KENTUCKY BLUEGRASS ( <i>Poa pratensis</i> )	6	0-13	86
IDAHO FESCUE ( <i>Festuca idahoensis</i> )	4	0-1	72
TIMOTHY ( <i>Phleum pratense</i> )	4	0-10	72

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME :	XERIC TO SUBMESIC
NUTRIENT REGIME :	SUBMESOTROPHIC TO MESOTROPHIC
ELEVATION:	1488(1397-1510)M
SOIL DRAINAGE :	RAPIDLY TO WELL
SLOPE:	16(0-50)%
ASPECT:	SOUTH EASTERLY

### FORAGE PRODUCTION KG/HA

GRASS	1160(654-1412)
FORB	712(382-1140)
SHRUB	39(0-118)
TOTAL	1574(660-2448)

SUGGESTED GRAZING CAPACITY  
0.6 HA/AUM

## SASMC2. Parry oatgrass-Kentucky bluegrass-Sedge (*Danthonia parryi*-*Poa pratensis*-*Carex* spp.)

n=4 This community type represents a heavily grazed Parry oatgrass-Rough fescue-Sedge community type. Heavy grazing pressure favours the growth of grazing resistant species of Kentucky bluegrass and sedge and causes rough fescue to decline. Continued heavy grazing pressure will eventually lead to a community type that is dominated by Kentucky bluegrass, timothy and sedge. If the grazing pressure is reduced on this community type there is a good possibility of recovery.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
SHRUBBY CINQUEFOIL ( <i>Potentilla fruticosa</i> )	2	1-3	100
SASKATOON ( <i>Amelanchier alnifolia</i> )	1	0-2	50
<b>FORBS</b>			
THREE FLOWERED AVENS ( <i>Geum triflorum</i> )	4	0-9	100
EARLY YELLOW LOCOWEED ( <i>Oxytropis sericea</i> )	6	0-16	75
SILKY PERENNIAL LUPINE ( <i>Lupinus sericeus</i> )	4	0-14	75
CUT-LEAVED ANEMONE ( <i>Anemone multifida</i> )	3	0-5	100
<b>GRASSES</b>			
PARRY OATGRASS ( <i>Danthonia parryi</i> )	13	6-17	100
KENTUCKY BLUEGRASS ( <i>Poa pratensis</i> )	8	2-16	100
SEDGE ( <i>Carex</i> spp)	3	0-3	75
SLENDER WHEATGRASS ( <i>Agropyron trachycaulum</i> )	1	0-1	25
TIMOTHY ( <i>Phleum pratense</i> )	4	0-15	75

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:  
XERIC TO SUBMESIC

NUTRIENT REGIME:  
SUBMESOTROPHIC TO MESOTROPHIC

ELEVATION:  
1419(1400-1441)M

SOIL DRAINAGE:  
RAPIDLY

SLOPE:  
25(13-30)%

ASPECT:  
SOUTHWESTERLY

### FORAGE PRODUCTION KG/HA

GRASS 919(0-2018)  
FORB 930(108-2330)  
SHRUB 60(0-238)  
TOTAL 1908(1408-2330)

SUGGESTED GRAZING CAPACITY  
0.5 HA/AUM

### SASMC3. Meadow foxtail-Kentucky bluegrass

(*Alopecurus pratensis*-*Poa pratensis*)

n=3 This community type was described in the South Sheep allotment. It represents small aspen stands that were cleared and seeded to a mixture of meadow brome, creeping red fescue, clover and meadow foxtail. These sites have continued to be heavily grazed which has favoured the growth of meadow foxtail, Kentucky bluegrass and clover. Meadow foxtail is not particularly palatable to livestock and therefore gains a competitive advantage over meadow brome and creeping red fescue, in these seeded areas.

#### PLANT COMPOSITION CANOPY COVER (%)

MEAN RANGE CONST.

##### SHRUBS

SHRUBBY CINQUEFOIL ( <i>Potentilla fruticosa</i> )	T	0-1	33
PRICKLY ROSE ( <i>Rosa acicularis</i> )	T	0-1	33

##### FORBS

COMMON DANDELION ( <i>Taraxacum officinale</i> )	8	4-12	100
STICKY PURPLE GERANIUM ( <i>Geranium viscosissimum</i> )	1	0-2	100
CLOVER ( <i>Trifolium spp.</i> )	11	0-32	67
WESTERN MEADOW RUE ( <i>Thalictrum occidentale</i> )	3	0-4	67

##### GRASSES

MEADOW FOXTAIL ( <i>Alopecurus pratensis</i> )	23	14-33	100
SEDGE ( <i>Carex spp.</i> )	4	1-6	100
KENTUCKY BLUEGRASS ( <i>Poa pratensis</i> )	24	14-36	100
TIMOTHY ( <i>Phleum pratense</i> )	1	1-2	100
RED FESCUE ( <i>Festuca rubra</i> )	6	0-15	67

#### ENVIRONMENTAL VARIABLES

##### MOISTURE REGIME:

MESIC TO SUBHYGRIC

##### NUTRIENT REGIME:

MESOTROPHIC TO PERMESOTROPHIC

##### ELEVATION:

1375(1300-1424)m

##### SOIL DRAINAGE:

WELL TO MODERATELY WELL

##### SLOPE:

7(2-10)%

##### ASPECT:

SOUTHERLY

#### FORAGE PRODUCTION KG/HA

GRASS 2775(2552-3132)

FORBS 507(306-608)

TOTAL 3282(2946-3160)

SUGGESTED GRAZING CAPACITY

0.3 HA/AUM

## SASMC4. Fringed sage/Kentucky bluegrass-Sedge

(*Artemisia frigida*/*Poa pratensis*-*Carex spp.*)

n=1 This community type was described on a south facing slope and ridge top in the South Sheep allotment. It appears to represent long-term heavy grazing pressure on a Parry oatgrass-Rough fescue-Sedge dominated community type. The increased grazing pressure on these south facing slopes favours the growth of fringed sage, sedge, Kentucky bluegrass and dandelion. It is unusual having such a high cover of Kentucky bluegrass on these slopes. Kentucky bluegrass usually prefers moister lower slope positions. Perhaps the higher precipitation received in the Subalpine subregion compared to the Montane makes the south facing slopes more favorable to Kentucky bluegrass invasion.

There is still a strong component of native species in this community type and recovery is likely if the grazing pressure is reduced.

### PLANT COMPOSITION CANOPY COVER (%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
PASTURE SAGE ( <i>Artemisia frigida</i> )	16	-	100
SHRUBBY CINQUEFOIL ( <i>Potentilla fruticosa</i> )	4	-	100
<b>FORBS</b>			
GOLDEN BEAN ( <i>Thermopsis rhombifolia</i> )	5	-	100
EARLY YELLOW LOCOWEED ( <i>Oxytropis sericea</i> )	2	-	100
COMMON DANDELION ( <i>Taraxacum officinale</i> )	2	-	100
NATIVE VETCH ( <i>Vicia americana</i> )	2	-	100
COMMON GOATS BEARD ( <i>Tragopogon dubius</i> )	1	-	100
<b>GRASSES</b>			
KENTUCKY BLUEGRASS ( <i>Poa pratensis</i> )	12	-	100
PARRY OATGRASS ( <i>Danthonia parryi</i> )	9	-	100
ROUGH FESCUE ( <i>Festuca scabrella.</i> )	8	-	100
COLUMBIA NEEDLE GRASS ( <i>Stipa columbiana</i> )	8	-	100
JUNEGRASS ( <i>Koeleria macrantha</i> )	6	-	100
SEDGE SPP. ( <i>Carex spp.</i> )	3	-	100

MOISTURE REGIME:

XERIC

NUTRIENT REGIME:

SUBMESOTROPHIC

ELEVATION:

1450M

SOIL DRAINAGE:

RAPIDLY

SLOPE:

33%

ASPECT:

SOUTHERLY

### FORAGE PRODUCTION KG/HA

GRASS	704
FORB	490
SHRUB	236
TOTAL	1430

SUGGESTED GRAZING CAPACITY

0.6 HA/AUM

### ENVIRONMENTAL VARIABLES

## SASMC5. Rough fescue-Kentucky bluegrass

*(Festuca scabrella-Poa pratensis)*

n=1 This community type was described in the South Sheep allotment west of Turner valley and represents a rough fescue grassland that has been heavily grazed to the point of Kentucky bluegrass invasion and is now recovering. Long-term heavy grazing pressure leads to a decline in rough fescue and an increase in Parry oatgrass and sedge species. Continued grazing pressure reduces the competitive advantage of rough fescue and the other native grass species and allows Kentucky bluegrass to establish on site. Protection or a reduction in stocking level at the point where Kentucky bluegrass becomes a significant of the community allows rough fescue to recover, but it seems Kentucky bluegrass also remains as co-dominant.

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

#### SHRUBS

LIMBER PINE  
*(Pinus flexilis)* 1 - 100

SHRUBBY CINQUEFOIL  
*(Potentilla fruticosa)* 10 - 100

#### FORBS

GOLDEN BEAN  
*(Thermopsis rhombifolia)* 1 - 100

SILKY PERENNIAL LUPINE  
*(Lupinus sericea)* 4 - 100

COMMON DANDELION  
*(Taraxacum officinale)* 1 - 100

NATIVE VETCH  
*(Vicia americana)* 1 - 100

MOUNTAIN GOLDENROD  
*(Solidago spathulata)* 2 - 100

#### GRASSES

KENTUCKY BLUEGRASS  
*(Poa pratensis)* 20 - 100

CALIFORNIA OATGRASS  
*(Danthonia californica)* 15 - 100

ROUGH FESCUE  
*(Festuca scabrella.)* 35 - 100

HAIRY WILDRYE  
*(Elymus innovatus)* 4 - 100

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:  
MESIC

NUTRIENT REGIME:  
PERMESOTROPHIC

ELEVATION:  
1500M

SOIL DRAINAGE:  
WELL

SLOPE:  
0%

### FORAGE PRODUCTION KG/HA

TOTAL 2000 \*ESTIMATE

SUGGESTED GRAZING CAPACITY 0.5HA/AUM
---

## SASMC6. Kentucky bluegrass-Rough fescue

(*Poa pratensis-Festuca scabrella*)

**n=13** Long-term heavy grazing pressure leads to a decline in rough fescue and an increase in Parry oatgrass and sedge species. Continued grazing pressure reduces the competitive advantage of rough fescue and the other native grass species and allows Kentucky bluegrass to establish on site. Continued heavy grazing pressure eventually leads to a decline in all native species and the plant community will resemble a Timothy-Kentucky bluegrass/dandelion type.

The forage productivity of this community type (2300 kg/ha) is equivalent to or better than a lightly grazed Rough fescue dominated community (1900 kg/ha). However, rough fescue is a more desirable forage species because it maintains its nutrient content into the dormant season. In contrast, Kentucky bluegrass loses its palatability and nutrient content if it is allowed to flower and set seed.

### PLANT COMPOSITION CANOPY COVER(%)

MEAN    RANGE    CONST.

#### SHRUBS

ASPEN  
(*Populus tremuloides*)    1    0-5    23

SHRUBBY CINQUEFOIL  
(*Potentilla fruticosa*)    2    1-8    92

#### FORBS

THREE FLOWERED AVENS  
(*Geum triflorum*)    3    0-16    69

WILD STRAWBERRY  
(*Fragaria virginiana*)    2    0-8    85

GRACEFUL CINQUEFOIL  
(*Potentilla gracilis*)    3    0-13    69

LATE YELLOW LOCOWEED  
(*Oxytropis monticola*)    1    0-5    54

YELLOW FALSE DANDELION  
(*Agoseris glauca*)    1    0-10    62

#### GRASSES

KENTUCKY BLUEGRASS  
(*Poa pratensis P.compressa*)<sup>7</sup>    2-21    100

TIMOTHY  
(*Phleum pratense*)    4    0-21    77

ROUGH FESCUE  
(*Festuca scabrella.*)    2    0-5    69

IDAHO FESCUE  
(*Festuca idahoensis*)    2    0-12    69

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:  
SUBXERIC TO SUBHYGRIC

NUTRIENT REGIME:  
SUBMESOTROPHIC TO PERMESOTROPHIC  
(POOR TO RICH)

ELEVATION:  
1630(1502-1798)M

SOIL DRAINAGE:  
VERY RAPIDLY TO MODERATELY WELL

SLOPE: 20(2-45)%

ASPECT: VARIABLE

### FORAGE PRODUCTION KG/HA

GRASS 1382(85-3584)  
FORB 887(126-2312)  
SHRUB 14(0-45)  
TOTAL 2258(421-4733)

SUGGESTED GRAZING CAPACITY  
0.4 ha/AUM

## SASMC7. Timothy-Kentucky bluegrass/Dandelion

(*Phleum pratense*-*Agropyron trachycaulum*/*Epilobium angustifolium*)

n=8 This community type was described in moist lower slope positions. Heavy grazing pressure has caused the native grass species to decline and allowed Canada bluegrass and timothy to invade onto the site. The presence of fireweed indicates that the site will likely succeed to an aspen dominated community type.

This community type is highly productive and it is easily accessible to domestic livestock. It should be rated as primary range.

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

#### SHRUBS

PRICKLY ROSE

(*Rosa acicularis*) 2 0-10 50

SHRUBBY CINQUEFOIL

(*Potentilla fruticosa*) 1 0-2 38

#### FORBS

GRACEFUL CINQUEFOIL

(*Potentilla gracilis*) 6 1-13 100

FIREWEED

(*Epilobium angustifolium*)2 0-8 86

COMMON DANDELION

(*Taraxacum officinale*) 11 1-26 100

NATIVE VETCH

(*Vicia americana*) 3 0-7 86

STICKY PURPLE GERANIUM

(*Geranium viscosissimum*)5 0-15 86

#### GRASSES

TIMOTHY

(*Phleum pratense*) 50 19-64 100

SLENDER WHEATGRASS

(*Agropyron trachycaulum*)2 0-10 42

ROUGH FESCUE

(*Festuca scabrella.*) 2 0-4 71

CALIFORNIA OATGRASS

(*Danthonia californica*) 2 0-15 14

KENTUCKY BLUEGRASS

(*Poa pratensis*) 4 0-11 86

SEDGE

(*Carex spp.*) 6 0-10 88

MOISTURE REGIME:

MESIC

NUTRIENT REGIME:

MESOTROPHIC

ELEVATION:

1511(1524-1655)M

SOIL DRAINAGE:

WELL

SLOPE:

5(1-5)%

ASPECT:

VARIABLE

### FORAGE PRODUCTION KG/HA

GRASS 4030(1368-7740)

FORB 863(550-1060)

SHRUB 33(0-144)

TOTAL 4926(2482-8494)

SUGGESTED GRAZING CAPACITY  
0.2 HA/AUM

### ENVIRONMENTAL VARIABLES



## SASMC8. Marsh reedgrass-Timothy/Cow parnsip

*(Calamagrostis canadensis-Phleum pratense/Heracleum lanatum)*

n=1 This community type represents seepage areas in the foothills west of Turner valley. Often these areas are invaded by willow to form the Willow/Marsh reedgrass dominated community type. This community type is very similar to the Marsh reedgrass/Cow parnsip community previously described, but this community has a high cover of timothy. Timothy, Kentucky bluegrass and dandelion will often invade these sites when exposed to heavy grazing pressure. This community type is highly productive and should be rated as primary range.

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

#### SHRUBS

RASPBERRY

*(Rubus idaeus)*

6 - 100

#### FORBS

COW PARSNIP

*(Heracleum lanatum)*

28 - 100

FIREWEED

*(Epilobium angustifolium)*3

- 100

COMMON DANDELION

*(Taraxacum officinale)*

3 - 100

CANADA VIOLET

*(Viola canadensis)*

20 - 100

WHITE GERANIUM

*(Geranium richardsonii)*

10 - 100

WESTERN MEADOW RUE

*(Thalictrum occidentale)*

15 - 100

#### GRASSES

TIMOTHY

*(Phleum pratense)*

29 - 100

SLENDER WHEATGRASS

*(Agropyron trachycaulum)*3

- 100

MOUNTAIN BROME

*(Bromus carinatus.)*

2 - 100

MARSH REEDGRASS

*(Calamagrostis canadensis)*37

- 100

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:

SUBHYGRIC

NUTRIENT REGIME:

PERMESOTROPHIC

ELEVATION:

1570M

SOIL DRAINAGE:

MOD. WELL

SLOPE:

10%

ASPECT:

VARIABLE

### FORAGE PRODUCTION KG/HA

GRASS 1369

FORB 1245

TOTAL 2613

SUGGESTED GRAZING CAPACITY

0.4 HA/AUM

## SASMC9. Idaho fescue-Rough fescue/Bearberry

*(Festuca idahoensis-Festuca scabrella/Arctostaphylos uva-ursi)*

**n=2** This community type represents a Rough fescue/Bearberry plant community that has been heavily to moderately grazed for a number of years. Increased grazing pressure causes rough fescue to decline and allows Idaho fescue, timothy and sedge species to increase. If grazing pressure is reduced on this site it will likely succeed back to a rough fescue dominated community.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
SHRUBBY CINQUEFOIL <i>(Potentilla fruticosa)</i>	1	1-2	100
COMMON JUNIPER <i>(Juniperus communis)</i>	7	0-15	50
<b>FORBS</b>			
BEARBERRY <i>(Arctostaphylos uva-ursi)</i>	22	13-30	100
OLD MANS WHISKERS <i>(Geum triflorum)</i>	16	4-27	100
STRAWBERRY <i>(Fragaria virginiana)</i>	11	4-16	100
YARROW <i>(Achillea millefolium)</i>	7	6-8	100
NORTHERN BEDSTRAW <i>(Galium boreale)</i>	6	5-7	100
AMERICAN VETCH <i>(Vicia americana)</i>	6	2-9	100
<b>GRASSES</b>			
IDAHO FESCUE <i>(Festuca idahoensis)</i>	26	20-31	100
ROUGH FESCUE <i>(Festuca scabrella)</i>	9	4-13	100
NORTHERN WHEATGRASS <i>(Agropyron dasystachyum)</i>	5	4-6	100
JUNEGRASS <i>(Koeleria macrantha)</i>	4	1-6	100

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:

SUBMESIC

NUTRIENT REGIME:

MESOTROPHIC

ELEVATION:

1500-1510(1505M)

SOIL DRAINAGE:

WELL

SLOPE:

2(1-3%)

ASPECT:

VARIABLE

### FORAGE PRODUCTION KG/HA

GRASS	1408(1012-1804)
FORB	862(434-1290)
SHRUB	43(0-86)
TOTAL	2313(2302-2324)

SUGGESTED GRAZING CAPACITY  
0.4 HA/AUM

**SUBALPINE SUBREGION**  
**SOUTHERN ECODISTRICTS**  
**FORESTED COMMUNITY TYPES**



**Figure 7.** Deciduous communities are not common in the Subalpine, but where they do occur they can be very productive for domestic livestock. This picture represents an Aspen/Rose/Pinegrass dominated community, which can be common on warmer sites in the Subalpine subregion of southern Alberta.

## **SUBALPINE SUBREGION**

### **FOREST ECOLOGY**

The forested plant communities in the subalpine can be split into three zones. These include the Pine, Spruce-Fir and Upper Subalpine zones (Strong and Leggat 1992). The Pine zone is the lowest elevation zone and is dominated by lodgepole pine and shows little evidence of succession to Engelmann spruce. It is within this zone that aspen and balsam poplar dominated plant communities can occur on warmer sites. The Spruce-Fir zone is located altitudinally above the Pine zone (Strong and Leggat 1992). This zone displays evidence of lodgepole pine succession to Engelmann spruce and subalpine fir. The Upper subalpine zone is characterized by open canopied vegetation that occurs between the Spruce-Fir zone and timberline. This zone is dominated by dwarfed Engelmann spruce, alpine fir, whitebark pine and alpine larch. Plant growth in this upper zone is limited by low temperatures, wind, low moisture and a short growing season (Strong and Leggat 1992). There are only a handful of forested community types described in this guide. Archibald et al. (1996), provide a good descriptions of the major forested community types in the Subalpine subregion that are not represented in this guide. Generally, these forested plant communities only provide limited forage for domestic livestock, but where deciduous communities are extensive they can be heavily utilized by livestock and should be considered secondary range.

**Table 6.** Deciduous and conifer communities of the Southern Rocky Mountains of the Subalpine subregion

Community number	Community type	Productivity (kg/ha)			Moisture	Drainage	Carrying capacity (ha/AUM)	
		Grass	Forb	Shrub				Total
<b>D. DECIDUOUS</b>								
SASMD1	Pb/Silverberry	-	-	-	444*	Submesic	Rapidly	Non-use
SASMD2	Aw/Rose/Pinegrass	603	621	112	1336	Mesic	Well	1.4
SASMD3	Aw/Fireweed/Meadow foxtail	1612	679	92	2383	Subxeric	Well	0.8
SASMD4	Aw/Rose/Canada bluegrass	1036	348	78	1462	Submesic	Well	1.2
SASMD5	Aw-Pb/Cow parsnip	328	1200	92	1620	Subhygric	Well	1.1
SASMD6	Aw-Pb/Cow parsnip/Timothy	328	1200	92	1620	Subhygric	Well	1.1
<b>E. CONIFER</b>								
SASME1	Pl/Juniper	-	-	-	-	Submesic	Rapidly	Non-use
SASME2	Pl/Pinegrass	210	180	154	472	Mesic	Well	Non-use
SASME3	Pl-Se/Moss	187	707	88	981	Mesic	Well	Non-use
SASME4	Sw-Aw/Alder/Hairy wildrye	-	-	-	155*	Mesic	Well	Non-use

\*Estimate

**SASMD1. Pb/Silverberry**  
(*Populus balsamifera/Elaeagnus commutata*)

**n=1** This community type was described along the banks of Pekisko creek in the Pekisko Creek allotment. It is characteristic of dry gravelly, river flats, which are periodically flooded in the spring. This community type is very similar to the yellow mountain avens community which have been described in both the Montane and Subalpine subregions.

Generally, there is little forage for domestic livestock in this community type and it should be rated as non-use.

**PLANT COMPOSITION**      **CANOPY COVER(%)**

	MEAN	RANGE	CONST.
<b>TREES</b>			
BALSAM POPLAR ( <i>Populus balsamifera</i> )	15	-	100
<b>SHRUBS</b>			
PRICKLY ROSE ( <i>Rosa acicularis</i> )	1	-	100
SILVERBERRY ( <i>Elaeagnus commutata</i> )	5	-	100
SHRUBBY CINQUEFOIL ( <i>Potentilla fruticosa</i> )	1	-	100
FALSE MOUNTAIN WILLOW ( <i>Salix pseudomonticola</i> )	2	-	100
<b>FORBS</b>			
ALPINE HEDYSARUM ( <i>Hedysarum alpinum</i> )	3	-	100
COMMON HORSETAIL ( <i>Equisetum arvense</i> )	1	-	100
CUT-LEAVED ANEMONE ( <i>Anemone multifida</i> )	1	-	100
GRACEFUL CINQUEFOIL ( <i>Potentilla gracilis</i> )	1	-	100
<b>GRASSES</b>			
BLUE BUNCH WHEATGRASS ( <i>Agropyron spicatum</i> )	1	-	100
JUNE GRASS ( <i>Koeleria macrantha</i> )	1	-	100
CANADA BLUEGRASS ( <i>Poa compressa</i> )	1	-	100
HAIRY WILD RYE ( <i>Elymus innovatus</i> )	1	-	100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME :  
SUBMESIC

NUTRIENT REGIME :  
MESOTROPHIC

ELEVATION:  
1828M

SOIL DRAINAGE :  
RAPIDLY

SLOPE:  
18%

ASPECT:  
SOUTHEAST

**FORAGE PRODUCTION KG/HA**

TOTAL 444 \*ESTIMATE

SUGGESTED GRAZING CAPACITY  
NON-USE

**SASMD2. Aw/Rose/Pinegrass**  
(*Populus tremuloides/Rosa/Calamagrostis rubescens*)

**n=20** This community type is similar to the Aw/Rose/Pinegrass community which is described on mesic sites with medium nutrient regimes in the Montane subregion. It may represent the transition from the Montane to the Subalpine. These sites were described at lower elevations in the Subalpine and generally had southerly aspects, making the conditions more suitable for growing aspen. The forage productivity on this community type is moderate and it should be rated as secondary range.

**PLANT COMPOSITION CANOPY COVER(%)**

MEAN RANGE CONST.

**TREES**

ASPEN

(*Populus tremuloides*) 34 20-50 100

WHITE SPRUCE

(*Picea glauca*) 2 0-5 25

BALSAM POPULAR

(*Populus balsamifera*) 2 0-15 20

**SHRUBS**

PRICKLY ROSE

(*Rosa acicularis*) 6 1-28 90

SNOWBERRY

(*Symphoricarpose albus*) 1 0-5 50

WHITE MEADOWSWEET

(*Spiraea betulifolia*) 4 0-31 50

WILD RED RASPBERRY

(*Rubus idaeus*) 1 0-8 30

**FORBS**

WILD VETCH

(*Vicia americana*) 5 1-23 100

CREAM COLORED VETCHLING

(*Lathyrus ochroleucus*) 7 0-17 95

WILD STRAWBERRY

(*Fragaria virginiana*) 5 0-11 95

SHOWY ASTER

(*Aster conspicuus*) 4 0-24 80

COMMON FIREWEED

(*Epilobium angustifolium*)5 0-24 80

COMMON DANDYLION

(*Taraxacum officinale*) 1 0-5 70

WESTERN CANADIAN VIOLET

(*Viola canadensis*) 3 0-10 30

**GRASSES**

PINEGRASS

(*Calamagrostis rubescens*)14 3-32 100

HAIRY WILDRYE

(*Elymus innovatus*) 5 0-9 85

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME :

SUBMESIC TO SUBHYGRIC

NUTRIENT REGIME :

MESOTROPHIC

ELEVATION:

1508(1400-1768)M

SOIL DRAINAGE :

WELL TO IMPERFECTLY

SLOPE:

13(6-20)%

ASPECT:

VARIABLE

**FORAGE PRODUCTION KG/HA**

GRASS 603(160-1512)

FORB 621(44-1151)

SHRUB 112(0-726)

TOTAL 1336(416-2034)

SUGGESTED GRAZING CAPACITY

1.4HA/AUM

### SASMD3. Aw/Fireweed/Meadow foxtail

(*Populus tremuloides/Epilobium angustifolium/Alopecurus pratensis*)

**n=1** This community type was described in the South Sheep allotment on an old range improvement area. Many aspen stands in the South Sheep were cleared and seeded to mixture of brome, meadow foxtail, creeping red fescue and clover. This site represents invasion of aspen back into these range improvement areas. Meadow foxtail is generally unpalatable to livestock and has persisted on these sites.

This community is very productive and efforts should be made to control the aspen invasion.

#### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>TREES</b>			
ASPEN			
( <i>Populus tremuloides</i> )	36	-	100
<b>SHRUBS</b>			
PRICKLY ROSE			
( <i>Rosa acicularis</i> )	8	-	100
PIN CHERRY			
( <i>Prunus pensylvanica</i> )	1	-	100
<b>FORBS</b>			
COMMON FIREWEED			
( <i>Epilobium angustifolium</i> )	12	-	100
STICKY PURPLE GERANIUM			
( <i>Geranium viscosissimum</i> )	8	-	100
LINDLEY'S ASTER			
( <i>Aster ciliolatus</i> )	4	-	100
COMMON VETCH			
( <i>Vicia americana</i> )	3	-	100
CREAM COLORED VETCHLING			
( <i>Lathyrus ochroleucus</i> )	2	-	100
<b>GRASSES</b>			
MEADOW FOXTAIL			
( <i>Alopecurus pratensis</i> )	32	-	100
ORCHARDGRASS			
( <i>Dactylis glomerata</i> )	17	-	100
KENTUCKY BLUEGRASS			
( <i>Poa pratensis</i> )	3	-	100
TIMOTHY			
( <i>Phleum pratense</i> )	7	-	100

#### ENVIRONMENTAL VARIABLES

MOISTURE REGIME :  
MESIC TO SUBHYGRIC

NUTRIENT REGIME :  
MESOTROPHIC

ELEVATION:  
1400M

SOIL DRAINAGE :  
WELL

SLOPE:  
10%

ASPECT:  
SOUTHWEST

#### FORAGE PRODUCTION KG/HA

GRASS 1612  
FORB 679  
SHRUB 92  
TOTAL 2383

SUGGESTED GRAZING CAPACITY  
0.8HA/AUM



**SASMD4. Aw/Rose/Canada bluegrass**  
(*Populus tremuloides*/*Rosa spp./Poa compressa*)

**n=1** This community type represents a Aw/Rose/Pinegrass community that has been heavily grazed and invaded by Canada bluegrass. Canada bluegrass is an introduced grass that increases with increased grazing pressure. As grazing pressure increases in these aspen dominated community types there is a shift away from native species (rose, pinegrass, asters, fireweed) to a community that is dominated by bluegrass, timothy, dandelion and clover species. The invasion of non-native invaders onto the site makes this community very productive for domestic livestock, but the presence of overgrazed communities indicates some type of distribution problem and the management of the disposition should be discussed.

**PLANT COMPOSITION CANOPY COVER(%)**

	MEAN	RANGE	CONST.
<b>TREES</b>			
ASPEN ( <i>Populus tremuloides</i> )	35	-	100
WHITE SPRUCE ( <i>Picea glauca</i> )	5	-	100
<b>SHRUBS</b>			
PRICKLY ROSE ( <i>Rosa acicularis</i> )	4	-	100
WILD RED RASPBERRY ( <i>Rubus idaeus</i> )	10	-	100
NORTHERN GOOSEBERRY ( <i>Ribes oxycanthoides</i> )	1	-	100
<b>FORBS</b>			
LINDLEY'S ASTER ( <i>Aster ciliolatus</i> )	22	-	100
WILD STRAWBERRY ( <i>Fragaria virginiana</i> )	6	-	100
COMMON DANDYLION ( <i>Taraxacum officinal</i> )	5	-	100
COMMON FIREWEED ( <i>Epilobium angustifolium</i> )	4	-	100
CREAM COLORED VETCHLING ( <i>Lathyrus ochroleucus</i> )	4	-	100
<b>GRASSES</b>			
CANADA BLUEGRASS ( <i>Poa compressa</i> )	30	-	100
PINE REEDGRASS ( <i>Calamagrostis rubescens</i> )	9	-	100
HAIRY WILDRYE ( <i>Elymus innovatus</i> )	9	-	100
AWNLESS BROME ( <i>Bromus inermis</i> )	2	-	100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME :  
SUBMESIC

NUTRIENT REGIME :  
MESOTROPHIC

ELEVATION:  
1420M

SOIL DRAINAGE :  
WELL

SLOPE:  
10%

ASPECT:  
SOUTHWEST

**FORAGE PRODUCTION KG/HA**

GRASS 1036  
FORB 348  
SHRUB 78  
TOTAL 1462

SUGGESTED GRAZING CAPACITY  
1.2HA/AUM

## SASMD5. Aw-Pb/Cow parsnip

(*Populus tremuloides*-*Populus balsamifera*/*Heracleum lanatum*)

n=3 Nutrient seepage occurs at some point in the growing season favouring the growth of cow parsnip. This community type is very similar to the Aw/Thimbleberry community described in the Montane subregion. Forage productivity on these sites is generally quite high because of the favourable moisture and nutrient conditions. Cow parsnip is palatable to livestock and maybe extensively utilized. This community type should be rated as secondary range.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>TREES</b>			
ASPEN			
( <i>Populus tremuloides</i> )	17	0-30	67
BALSAM POPLAR			
( <i>Populus balsamifera</i> )	18	10-30	100
<b>SHRUBS</b>			
WILD RED RASPBERRY			
( <i>Rubus idaeus</i> )	1	0-1	67
NORTHERN GOOSEBERRY			
( <i>Ribes oxycanthoides</i> )	2	0-5	33
<b>FORBS</b>			
CREAM COLORED VETCHLING			
( <i>Lathyrus ochroleucus</i> )	5	1-13	100
COW PARSNIP			
( <i>Heracleum lanatum</i> )	31	8-48	100
LINDLEY'S ASTER			
( <i>Aster ciliolatus</i> )	22	20-24	100
WESTERN CANADA VIOLET			
( <i>Viola canadensis</i> )	6	1-15	100
TALL LUNGWORT			
( <i>Mertensia paniculata</i> )	5	1-12	100
TALL LARKSPUR			
( <i>Delphinium glaucum</i> )	5	1-11	100
COMMON FIREWEED			
( <i>Epilobium angustifolium</i> )	4	3-5	100
COMMON DANDYLION			
( <i>Taraxacum officinale</i> )	2	1-2	100
<b>GRASSES</b>			
MARSH REEDGRASS			
( <i>Calamagrostis canadensis</i> )	4	0-7	67
HAIRY WILDRYE			
( <i>Elymus innovatus</i> )	2	1-2	100
AWNLESS BROME			
( <i>Bromus inermis</i> )	2	1-5	100

### TIMOTHY

(*Phleum pratense*) 2 1-3 100

### SEDGE

(*Carex spp.*) 1 1-2 100

### ENVIRONMENTAL VARIABLES

#### MOISTURE REGIME :

SUBHYGRIC

#### NUTRIENT REGIME :

MESOTROPHIC-PERMESOTROPHIC

#### ELEVATION:

1400M

#### SOIL DRAINAGE :

WELL

#### SLOPE:

8%

#### ASPECT:

SOUTHWEST

### FORAGE PRODUCTION KG/HA

GRASS 328  
 FORB 1200  
 SHRUB 92  
 TOTAL 1620

### SUGGESTED GRAZING CAPACITY

1.1HA/AUM

## SASMD6. Aw-Pb/Cow parsnip/Timothy

(*Populus tremuloides*-*Populus balsamifera*/*Heracleum lanatum*/*Phleum pratense*)

n=1 Nutrient seepage occurs at some point in the growing season favouring the growth of cow parsnip. This community type is very similar to the previously described Aw-Pb/Cow parsnip community, but this community has a high cover of timothy. Increased grazing pressure will often allow timothy, Kentucky bluegrass and dandelion to invade onto these sites. Forage productivity on these sites is generally quite high because of the favourable moisture and nutrient conditions. Cow parsnip is palatable to livestock and maybe extensively utilized. This community type should be rated as secondary range.

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>TREES</b>			
<b>ASPEN</b>			
( <i>Populus tremuloides</i> )	26	-	100
<b>BALSAM POPLAR</b>			
( <i>Populus balsamifera</i> )	16	-	100
<b>SHRUBS</b>			
<b>WILD RED RASPBERRY</b>			
( <i>Rubus idaeus</i> )	4	-	100
<b>FORBS</b>			
<b>CREAM COLORED VETCHLING</b>			
( <i>Lathyrus ochroleucus</i> )	2	-	100
<b>COW PARSNIP</b>			
( <i>Heracleum lanatum</i> )	2	-	100
<b>LINDLEY'S ASTER</b>			
( <i>Aster ciliolatus</i> )	18	-	100
<b>WESTERN CANADA VIOLET</b>			
( <i>Viola canadensis</i> )	2	-	100
<b>TALL LUNGWORT</b>			
( <i>Mertensia paniculata</i> )	1	-	100
<b>WESTERN MEADOW RUE</b>			
( <i>Thalictrum occidentale</i> )	4	-	100
<b>COMMON FIREWEED</b>			
( <i>Epilobium angustifolium</i> )	1	-	100
<b>COMMON DANDELION</b>			
( <i>Taraxacum officinale</i> )	6	-	100
<b>GRASSES</b>			
<b>MARSH REEDGRASS</b>			
( <i>Calamagrostis rubescens</i> )	4	-	100
<b>SMOOTH WILD RYE</b>			
( <i>Elymus glaucus</i> )	7	-	100
<b>AWNLESS BROME</b>			
( <i>Bromus inermis</i> )	2	-	100
<b>TIMOTHY</b>			
( <i>Phleum pratense</i> )	18	-	100

SEDGE  
(*Carex spp.*) 3 - 100

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME :  
SUBHYGERIC

NUTRIENT REGIME :  
PERMESOTROPHIC

ELEVATION:  
1510M

SOIL DRAINAGE :  
MOD. WELL

SLOPE:  
10%

ASPECT:  
VARIABLE

### FORAGE PRODUCTION KG/HA

GRASS 328  
FORB 1200  
SHRUB 92  
TOTAL 1620

SUGGESTED GRAZING CAPACITY  
1.1HA/AUM

**SASME1. PI/Juniper**  
(*Pinus contorta*/*Juniperus spp.*)

n=1 Dry site conditions from south exposures or coarse textured soils are characteristic of this community type (Archibald et al. 1996). The dry site conditions limit the amount of forage this site can produce and the steep slope limits access to livestock. As a result, this community would be considered non-use.

**PLANT COMPOSITION CANOPY COVER(%)**

	MEAN	RANGE	CONST.
<b>TREES</b>			
LODGEPOLE PINE ( <i>Pinus contorta</i> )	15	-	100
<b>SHRUBS</b>			
GROUND JUNIPER ( <i>Juniperus communis</i> )	15	-	100
CREEPING JUNIPER ( <i>Juniperus horizontalis</i> )	2	-	100
CANADA BUFFALOBERRY ( <i>Shepherdia canadensis</i> )	6	-	100
PRICKLY ROSE ( <i>Rosa acicularis</i> )	3	-	100
CHOKO CHERRY ( <i>Prunus virginiana</i> )	2	-	100
<b>FORBS</b>			
YELLOW HEDYSARUM ( <i>Hedysarum sulphurescens</i> )	3	-	100
BEARBERRY ( <i>Arctostaphylos uva-ursi</i> )	8	-	100
SPREADING DOGBANE ( <i>Apocynum androsaemifolium</i> )	15	-	100
LINDLEY'S ASTER ( <i>Aster ciliolatus</i> )	1	-	100
<b>GRASSES</b>			
HAIRY WILD RYE ( <i>Elymus innovatus</i> )	1	-	100
SEDGE ( <i>Carex spp.</i> )	T	-	100

MOISTURE REGIME :  
SUBMESIC

NUTRIENT REGIME :  
OLIGOTROPHIC

ELEVATION:  
1659M

SOIL DRAINAGE :  
RAPIDLY

SLOPE:  
22%

ASPECT:  
SOUTH

**FORAGE PRODUCTION KG/HA**

TOTAL 350 \*ESTIMATE

SUGGESTED GRAZING CAPACITY  
NON-USE

**ENVIRONMENTAL VARIABLES**

**SASME2. Pl/Pinegrass**  
(*Pinus contorta/Calamagrostis rubescens*)

n=1 This community type is very similar to the Pl/Pinegrass dominated community described in the Montane subregion (Willoughby et al. 2003). Succession will be to white spruce, but the extensive fire history in the area has resulted in a predominance of lodgepole pine (Archibald et al 1996). Pinegrass is generally unpalatable to livestock, but if grazed early in the spring they will utilize it as a forage source. The forage productivity of this community type is quite low. As a result this community type should be rated as non-use.

**PLANT COMPOSITION CANOPY COVER(%)**

	MEAN	RANGE	CONST.
<b>TREES</b>			
LODGEPOLE PINE ( <i>Pinus contorta</i> )	70	-	100
<b>SHRUBS</b>			
PRICKLY ROSE ( <i>Rosa acicularis</i> )	3	-	100
WHITE MEADOWSWEET ( <i>Spiraea betulifolia</i> )	11	-	100
DWARF BILBERRY ( <i>Vaccinium caespitosum</i> )7		-	100
GROUND JUNIPER ( <i>Juniperus communis</i> )	5	-	100
<b>FORBS</b>			
TWINFLOWER ( <i>Linnaea borealis</i> )	17	-	100
SHOWY ASTER ( <i>Aster conspicuus</i> )	4	-	100
CREAM COLORED VETCHLING ( <i>Lathyrus ochroleucus</i> )	2	-	100
WILD STRAWBERRY ( <i>Fragaria virginiana</i> )	1	-	100
HEART-LEAVED ARNICA ( <i>Arnica cordifolia</i> )	3	-	100
WESTERN LOUSEWORT ( <i>Pedicularis bracteosa</i> )	1	-	100
<b>GRASSES</b>			
PINEGRASS ( <i>Calamagrostis rubescens</i> )27		-	100
HAIRY WILDRYE ( <i>Elymus innovatus</i> )	4	-	100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME :  
MESIC

NUTRIENT REGIME :  
MESOTROPHIC

ELEVATION:  
1676M

SOIL DRAINAGE :  
WELL

SLOPE:  
10%

ASPECT:  
EAST

**FORAGE PRODUCTION KG/HA**

GRASS 210  
FORB 180  
SHRUB 154  
TOTAL 472

SUGGESTED GRAZING CAPACITY  
NON-USE

**SASME3. PI-Se/Moss**  
(*Pinus contorta-Picea engelmannii/Moss spp.*)

n=5 This community type represents the modal conditions for the Subalpine subregion at mid to lower elevations. Lodgepole pine, Engelmann spruce and subalpine fir can all occur as the dominant tree species on this ecological site. In general succession is from lodgepole pine to Engelmann spruce and subalpine fir. However, lodgepole pine is the most common tree species because of the frequency of fire.

There is little forage for domestic livestock in this community type. As a result, this community should be rated as non-use.

**PLANT COMPOSITION**

	MEAN	RANGE	CONST.
<b>TREES</b>			
ENGELMANN SPRUCE ( <i>Picea engelmannii</i> )	3	0-10	40
WHITE SPRUCE ( <i>Picea glauca</i> )	2	0-10	20
LODGEPOLE PINE ( <i>Pinus contorta</i> )	30	15-60	100
<b>SHRUBS</b>			
PRICKLY ROSE ( <i>Rosa acicularis</i> )	1	0-1	80
DWARF BILBERRY ( <i>Vaccinium caespitosum</i> ) <sup>2</sup>		1-3	100
WHITE MEADOWSWEET ( <i>Spiraea betulifolia</i> )	2	0-5	80
GREEN ALDER ( <i>Alnus crispa</i> )	3	0-8	40
<b>FORBS</b>			
BUNCHBERRY ( <i>Cornus canadensis</i> )	8	1-14	100
WILD STRAWBERRY ( <i>Fragaria virginiana</i> )	2	1-4	100
SHOWY ASTER ( <i>Aster conspicuus</i> )	2	0-3	80
TWINFLOWER ( <i>Linnaea borealis</i> )	2	0-6	80
BROAD-LEAVED ARNICA ( <i>Arnica latifolia</i> )	1	0-4	40
<b>GRASSES</b>			
HAIRY WILDRYE ( <i>Elymus innovatus</i> )	1	0-1	80
PINEGRASS ( <i>Calamagrostis rubescens</i> ) <sup>3</sup>		0-5	60
RICHARDSON NEEDLEGRASS ( <i>Stipa richardsonii</i> )	T	0-1	20

**Moss**

(*Moss spp*)                      10            1-38            100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME :

XERIC TO MESIC

NUTRIENT REGIME :

SUBMESOTROPHIC TO MESOTROPHIC

ELEVATION:

1647(1536-1770)M

SOIL DRAINAGE : RAPIDLY TO WELL

SLOPE: 9(0-22)%

ASPECT: VARIABLE

**FORAGE PRODUCTION KG/HA**

GRASS 187

FORB 707

SHRUB 88

TOTAL 981

SUGGESTED GRAZING CAPACITY  
NON-USE

**SASME4. Sw-Aw/Alder/Hairy wildrye**  
*(Picea glauca-Populus tremuloides/Alnus crispa/Elymus innovatus)*

**n=1** This community type represents the succession of spruce onto an aspen dominated community type. As succession occurs from aspen to spruce there is a corresponding drop in forage production. This community type has little forage available for domestic livestock and should be rated as non-use.

**PLANT COMPOSITION CANOPY COVER(%)**

	MEAN	RANGE	CONST.
<b>TREES</b>			
WHITE SPRUCE <i>(Picea glauca)</i>	31	-	100
ASPEN <i>(Populus tremuloides)</i>	16	-	100
LODGEPOLE PINE <i>(Pinus contorta)</i>	3	-	100
<b>SHRUBS</b>			
PRICKLY ROSE <i>(Rosa acicularis)</i>	3	-	100
DWARF BILBERRY <i>(Vaccinium caespitosum)</i>	3	-	100
WHITE MEADOWSWEET <i>(Spiraea betulifolia)</i>	5	-	100
GREEN ALDER <i>(Alnus crispa)</i>	10	-	100
<b>FORBS</b>			
ONE SIDED WINTERGREEN <i>(Orthilia secunda)</i>	2	-	100
SHOWY ASTER <i>(Aster conspicuus)</i>	2	-	100
TWINFLOWER <i>(Linnaea borealis)</i>	1	-	100
<b>GRASSES</b>			
HAIRY WILDRYE <i>(Elymus innovatus)</i>	3	-	100
PINEGRASS <i>(Calamagrostis rubescens)</i>	3	-	100
<b>MOSS</b>			
<i>(Moss spp)</i>	10	-	100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME :

MESIC

NUTRIENT REGIME :

MESOTROPHIC

ELEVATION:

1557M

SOIL DRAINAGE : WELL

SLOPE: 22%

ASPECT: NORTHERLY

**FORAGE PRODUCTION KG/HA**

TOTAL 155

SUGGESTED GRAZING CAPACITY

NON-USE

## ALPINE SUBREGION

### NATIVE GRASSLANDS AND SHRUBLANDS



**Figure 8.** Low growing plant communities of white mountain avens and bog sedge on windswept ridges, with arctic willow, heather and blackening sedge in snow accumulation areas are typical of the Alpine subregion



## **Alpine communities**

The alpine environment generally occurs above timberline. Ogilvie (1969), found timberline to be controlled by low temperature, wind dessication, avalanching and snow depth. The alpine plants and communities show adaptations to these extreme environmental conditions. The alpine plant communities tend to be low growing where they are protected from the wind and benefit from the warmer temperatures close to the ground (Ogilvie 1969). Figure 4 outlines the alpine communities in the landscape of the Rocky Mountains. On south facing, wind swept ridges the bog sedge and white dryad communities are found. At slightly lower elevations where snow accumulates the low growing willow communities predominate (arctic willow, snow willow, rock willow). On the north facing slopes where snow accumulates the blackening sedge and heather community types are found. In the valley bottoms below timberline the willow, bog birch, and grassy meadow community types are typical. The sequence of the valley bottom community types in the landscape is described in Figure 2. The plant community types found in the Alpine subregion are listed in Table 7.

Many of these alpine grass and shrublands are very fragile because of exposure and cold climate. The forage productivity is very low compared to the valley bottoms. Consequently, recovery from overgrazing will likely take some time. As a result grazing by domestic livestock in the Alpine subregion should be discouraged.

**Table 7.** Native grass and shrublands of the Alpine subregion

Community number	Community type	Productivity (kg/ha)		Moisture	Drainage	Carrying capacity (ac/AUM)
		Grass	Forb Shrub Total			
<b>A. GRASSLANDS</b>						
ALPA1.	Bog sedge		N/A	Subxeric	Rapidly	Non-use
ALPA2.	White Mountain Avens			N/A	Subxeric	Rapidly Non-use
ALPA3.	Mountain heather		N/A	Submesic	Well	Non-use
ALPA4.	Blackening sedge		N/A	Subhygric	Mod. Well	Non-use
ALPA5.	Simple bog sedge		N/A	Subhygric	Mod. Well	Non-use
ALPA6.	Arctic willow		N/A	Subhygric	Mod. Well	Non-use

## ALPA1. Bog sedge

(*Kobresia myosuroides*)

n=5 This community type occurs at higher elevations on snow-free, wind-exposed south facing slopes and ridge crests. The soils are shallow, stoney colluvial Regosols (Corns and Achuff 1982). Ogilvie (1969), found that there was rich herb layer, and an abundant lichen and bryophyte layer in this community type.

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

#### SHRUBS

SMOOTH WILLOW

(*Salix glauca*) 3 0-8 60

BOG BIRCH

(*Betula glandulosa*) 3 0-7 60

WHITE MOUNTAIN AVENS

(*Dryas integrifolia*,  
*D. octopetala*) 4 0-7 40

#### FORBS

ALPINE HEDYSARUM

(*Hedysarum alpinum*) 5 0-8 80

LITTLE CLUBMOSS

(*Selaginella densa*) 1 0-3 40

ALPINE BISTORT

(*Polygonum viviparum*) 2 0-4 60

SMOOTH LEAVED CINQUEFOIL

(*Potentilla diversifolia*) 1 0-2 60

#### GRASSES

BOG SEDGE

(*Kobresia myosuroides*) 44 35-65 100

HAIRY WILDRYE

(*Elymus innovatus*) 4 0-10 80

ROCKY MTN. FESCUE

(*Festuca saximontana*) 1 0-5 40

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:

SUBXERIC

NUTRIENT REGIME:

OLIGOTROPHIC

ELEVATION:

2042( 1900-2260) M

SOIL DRAINAGE:

RAPIDLY

SLOPE:

0-48(19)%

ASPECT:

SOUTH -SOUTHWEST

### FORAGE PRODUCTION KG/HA

SUGGESTED GRAZING CAPACITY

NON-USE

## ALPA2. White mountain avens

(*Dryas octopetala*, *D. integrifolia*)

n=43 This community type occurs on wind-exposed, snow-free ridges. The soils are shallow, stoney, colluvial Regosols (Corns and Achuff 1982). Ogilvie (1969), found this community to have an abundance of cushion and mat-plants and a large number of lichens.

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

#### SHRUBS

WHITE MOUNTAIN AVENS

(*Dryas octopetala*,  
*D. integrifolia*) 33 1-70 95

SNOW WILLOW

(*Salix reticulata*) 2 0-15 65

WHITE MOUNTAIN HEATHER

(*Cassiope tetragona*) 3 0-35 23

#### FORBS

ELEPHANT'S HEAD

(*Pedicularis groenlandicum*)T 0-2 9

WOOLY EVERLASTING

(*Antennaria lanata*) 1 0-25 19

MOSS CAMPION

(*Silene acaulis*) 1 0-5 72

ALPINE BISTORT

(*Polygonum viviparum*) 1 0-5 65

LITTLE CLUBMOSS

(*Selaginella densa*) 1 0-15 33

#### GRASSES

BOG SEDGE

(*Kobresia myosuroides*) 3 0-35 35

SEDE SPP.

(*Carex spp.*) 2 0-65 77

SPIKED TRisetum

(*Trisetum spicatum*) 1 0-7 42

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:

SUBXERIC

NUTRIENT REGIME:

SUBMESOTROPHIC

ELEVATION:

2192( 1410-2490) M

SOIL DRAINAGE:

RAPIDLY

SLOPE:

0-60(25)%

ASPECT:

EAST TO WEST

### FORAGE PRODUCTION

SUGGESTED GRAZING CAPACITY

NON-USE

## ALPA3. Mountain heather

(*Cassiope spp.*-*Phyllodoce spp.*)

**n=58** This community type occurs on north facing slopes, with deep snow cover. The soils are Brunisolic, Podzolic and Regosolic and they have seepage and solifluction (Corns and Achuff 1982). Ogilvie (1969), found this community type to have an abundant low shrub layer, a rich byrophyte and lichen layer and a moderately developed herb layer.

This community type includes both the *Cassiope tetragona*-*Dryas octopetala*-*Salix nivalis* and *Phyllodoce glanduliflora*-*Cassiope mertensiana*-*Antennaria lanata* community types described by Corns and Achuff (1982).

### PLANT COMPOSITION CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
WILLOW SPP. ( <i>Salix spp.</i> )	13	0-30	67
WESTERN MOUNTAIN HEATHER ( <i>Cassiope mertensiana</i> )	27	0-75	86
WHITE MOUNTAIN HEATHER ( <i>Cassiope tetragona</i> )	1	0-15	16
YELLOW HEATHER ( <i>Phyllodoce glanduliflora</i> )	20	0-60	20
RED HEATHER ( <i>Phyllodoce empetriformis</i> )	5	0-25	40
<b>FORBS</b>			
LANCED -LEAVED PAINT BRUSH ( <i>Castilleja occidentalis</i> )	T	0-1	16
WOOLLY EVERLASTING ( <i>Antennaria lanata</i> )	3	0-15	85
WESTERN ANEMONE ( <i>Anemone occidentalis</i> )	1	0-30	26
MOUNTAIN SAGE ( <i>Artemisia norvegica</i> )	1	0-15	53
<b>GRASSES</b>			
REDDISH WOOD RUSH ( <i>Luzula piperi</i> )	T	0-5	33
SEDGE ( <i>Carex spp.</i> )	1	0-12	66

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:  
SUBMESIC

NUTRIENT REGIME:  
MESOTROPHIC

ELEVATION:  
2154(194-2410) M

SOIL DRAINAGE:  
WELL

SLOPE:  
24(2-75)%

ASPECT:  
NORTHERLY

### FORAGE PRODUCTION KG/HA

NOT AVAILABLE

<p>SUGGESTED GRAZING CAPACITY NON-USE</p>
---

## ALPA4. Blackening sedge

(*Carex nigricans*)

n=7 This is a snowpatch community occurring in small channels and depressions where there is very deep snow accumulation (Ogilvie 1969). The soils are predominantly Brunisols, Podzols and Regosols and are moist and free of snow for only a brief period of time (Corns and Achuff 1982). Ogilvie (1969) found these community types to have an abundant herb layer of sedges, grasses and forbs, with only a minor occurrence of dwarf shrubs and mosses.

### PLANT COMPOSITION CANOPY COVER(%)

MEAN RANGE CONST.

#### SHRUBS

WILLOW

(*Salix spp*)

T 0-2 14

#### FORBS

WOOLLY EVERLASTING

(*Antennaria lanata*)

6 0-34 57

MOUNTAIN MARIGOLD

(*Caltha leptosepala*)

7 0-40 43

MOUNTAIN BUTTERCUP

(*Ranunculus eschscholtzii*)

1 0-5 29

MARE'S TAIL

(*Hippus vulgaris*)

2 0-12 14

#### GRASSES

BLACKENING SEDGE

(*Carex nigricans*)

49 0-90 100

TUFTED HAIRGRASS

(*Deschampsia caespitosa*)

4 0-30 14

REDDISH WOOD RUSH

(*Luzula piperi*)

2 0-8 57

MOUNTAIN HAIRGRASS

(*Vahlodea atropurpurea*)

4 0-15 57

WHITE RUSH

(*Juncus albenscens*)

2 0-15 14

### ENVIRONMENTAL VARIABLES

MOISTURE REGIME:

MESIC TO SUBHYGRIC

NUTRIENT REGIME:

MESOTROPHIC

ELEVATION:

2164(1990-2240) M

SOIL DRAINAGE:

MODERATELY WELL

SLOPE:

25( 1-64)%

ASPECT:

NORTHERLY

### FORAGE PRODUCTION KG/HA

SUGGESTED GRAZING CAPACITY

NON-USE

**ALPA5. Simple bog sedge**  
(*Kobresia simpliciuscula*)

**n=1** This community type was described at only one site. Simple bog sedge is typical of boggy areas at higher elevations. This community type was described on a poorly drained, level site at higher elevations. It is likely found in association with the willow and sedge dominated community types found in the valley bottoms of the lower alpine and upper subalpine.

**PLANT COMPOSITION** CANOPY COVER(%)

	MEAN	RANGE	CONST.
<b>SHRUBS</b>			
WILLOW SPP. ( <i>Salix spp.</i> )	3	-	100
BOG BIRCH ( <i>Betula glandulosa</i> )	2	-	100
<b>FORBS</b>			
BOG ASPHODEL ( <i>Tofieldia pusilla</i> )	8	-	100
YELLOW MOUNTAIN SAXIFRAGE ( <i>Saxifraga aizoides</i> )	5	-	100
SMALL WOOD ANEMONE ( <i>Anemone parviflora</i> )	2	-	100
BROAD LEAVED FIREWEED ( <i>Epilobium latifolium</i> )	2	-	100
ALPINE BISTORT ( <i>Polygonum viviparum</i> )	1	-	100
<b>GRASSES</b>			
SIMPLE BOG SEDGE ( <i>Kobresia simpliciuscula</i> )	25	-	100
BALTIC RUSH ( <i>Juncus balticus</i> )	1	-	100

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:  
SUBHYGRIC

NUTRIENT REGIME:  
PERMESOTROPHIC

ELEVATION:  
1900 M

SOIL DRAINAGE:  
MODERATELY WELL

**FORAGE PRODUCTION KG/HA**

SUGGESTED GRAZING CAPACITY  
NON-USE

**ALPA6. Arctic willow**  
(*Salix arctica*)

**n=20** This dwarf shrub community occurs in areas of deep snow accumulation, on soils which receive seepage for much of the growing season (Corns and Achuff 1982). Ogilvie (1969), found that there was a rich herb layer of sedges, grasses and forbs and an abundant dwarf shrub layer. This community is similar to the blackening sedge community previously described, but it appears melt out occurs earlier in this community type (Corns and Achuff 1982)

**PLANT COMPOSITION CANOPY COVER(%)**

MEAN RANGE CONST.

**SHRUBS**

ARCTIC WILLOW . ( <i>Salix arctica</i> )	17	0-50	80
SNOW WILLOW ( <i>Salix reticulata</i> )	4	0-30	40
ROCK WILLOW ( <i>Salix vestita</i> )	6	0-65	15

**FORBS**

WOOLY EVERLASTING ( <i>Antennaria lanata</i> )	7	0-35	50
WILD STRAWBERRY ( <i>Fragaria virginiana</i> )	T	0-5	15
SMOOTH LEAVED CINQUEFOIL ( <i>Potentilla diversifolia</i> )	1	0-3	35
MOUNTAIN SAGE ( <i>Artemisia norvegica</i> )	2	0-5	50
ALPINE BISTORT ( <i>Polygonum viviparum</i> )	1	0-5	55

**GRASSES**

SEDGE ( <i>Carex spp</i> )	3	0-20	35
HAIRY WILDRYE ( <i>Elymus innovatus</i> )	1	0-15	5
MOUNTAIN TIMOTHY ( <i>Phleum commutatum</i> )	1	0-10	25

**ENVIRONMENTAL VARIABLES**

MOISTURE REGIME:

MESIC TO SUBHYGRIC

NUTRIENT REGIME:

MESOTROPHIC

ELEVATION:

2141(1830-2330)m

SOIL DRAINAGE:

MODERATELY WELL TO IMPERFECT

SLOPE:

25(11-58)%

ASPECT:

VARIABLE

**FORAGE PRODUCTION KG/HA**

SUGGESTED GRAZING CAPACITY  
NON-USE



### Literature cited

Archibald, J.H., G. Klappstein and I.G.W. Corns. 1996. Field guide to ecosites of Southwestern Alberta. Nat. Resour. Can., Can. For. Serv. Northwest Reg., North. For. Cent., Edmonton, AB. Spec rep. no. 8.

Bailey, A.W., M.G. Willoughby, R. Johansen and S. Smith. (1992) Management of Yukon Rangelands. Renewable Resources, Yukon Territorial Government, Whitehorse, Yukon. 55pp. ISBN-1-55018-138-6.

Bork, E. 1990. Clearwater allotment: Prescribed burn vegetation survey, evaluation, and discussion. Range Management Section, Alberta Forest Service, Edmonton, Alta. 29pp.

Bork, E. 1994. Ecological classification and management of native ranges in Willmore Wilderness Park. Range management section, Alberta Environmental Protection. Edmonton, Alta. Pub. no. T/282 65pp.

Beckingham, J., I.G.W. Corns and J.H. Archibald. 1996. Field guide to Ecosites of West-Central Alberta. Nat. Resour. Can., Can. For. Serv. Northwest Reg., North. For. Cent., Edmonton, AB. Spec rep. no. 9.

Beckingham, J. and J.H. Archibald. 1996. Field guide to ecosites of Northern Alberta. Nat. Resour. Can., Can. For. Serv. Northwest Reg., North. For. Cent., Edmonton, AB. Spec rep. no. 5.

Corns, I.G.W. and P. Achuff. 1982. In Ecological (Biophysical) Land Classification of Banff and Jasper National Parks. Vol II: Soil and Vegetation Resources. Edited by Holland, W.D. and G.M. Coen. Environment Canada and Alberta Institute of Pedology, University of Alberta. Pub. no. SS-82-44. 540pp.

Corns, I.G.W. and R.M. Annas. 1986. Field guide to forest ecosystems of West-Central Alberta. Northern Forestry Center, Canadian Forestry Service, Edmonton, Alta. 251pp.

Daubenmire, R. 1952. Forest vegetation of Northern Idaho and adjacent Washington and its bearing on concepts of vegetation classification. Ecol. Mongr. 22: 301-330.

Department of Environmental Protection. 1994. Natural Regions of Alberta. Alberta Environmental Protection. Edmonton, Alta. Pub. no.: I/531. 18pp.

Gauch, H.G. 1982. Multivariate analysis in community ecology. Cambridge University Press, Cambridge, 298pp.

Holland, W.D. and G.M. Coen. 1982. Ecological (Biophysical) Land Classification of Banff and Jasper National Parks. Vol II: Soil and Vegetation Resources. Environment Canada and Alberta Institute of Pedology, University of Alberta. Pub. no. SS-82-44. 540pp.

Jaques, D.R. 1976. Winter alpine-subalpine wildlife habitat in the southern Rocky Mountains of Alberta. Kananaskis Center for Environmental Research, University of Calgary. Calgary, Alta. 113pp.

Lane, C.T., M.G. Willoughby and M.J. Alexander. 2000. Range plant community types and Carrying capacity for the Lower Foothills Subregion. 2<sup>nd</sup> Approximation. Alberta Environment. Land and Forest Service. Edmonton. Pub.no. T/532. 232 pp.

Mackinnon, A., J. Pojar and R. Coupe. 1992. Plants of Northern British Columbia. Lone Pine Publishing, Edmonton, Alta. 345pp.

Mueggler, W.F. 1988. Aspen community types of the Intermountain Region. U.S.D.A. Intermountain Research Station. INT-250. 133pp.

Morgantini, L. E. and W.B. Russell. 1983. An assesement of three selected elk winter ranges in the Rocky Mountains Region. Alberta Fish and Wildlife Division. Edmonton. AB. 264pp.

Moss, E.H. and J.W. Campbell. 1947. The fescue grassland of Alberta. Can. J. Res. 25:209-227.

Ogilvie, R.T. 1969. The Mountain Forest and Alpine Zones of Alberta. In, Vegetation, Soils and Wildlife. edited by J.G. Nelson and M.J. Chambers. Methuen Publications, Toronto. pg 24-44.

Range Survey Manual. 1992. Range Management Section, Alberta Forest Service. Edmonton, Alta. 39pp.

Scoggan, H.J. 1978. The Flora of Canada. National Museum of Natural Sciences, Publications in Botany, Ottawa, Canada. No. 7(2).

Stringer, P.W. 1973. An ecological study of grasslands in Banff, Jasper and Waterton National Parks. Can. J. Bot. Vol. 51. 383-411.

Strong, W.L. 1992. Ecoregions and ecodistricts of Alberta. Vol 1. Alberta Forestry Lands and Wildlife, Land Information Services Division, Resource Information Branch, Edmonton, Alta. T/244. 77pp.

Strong, W.L. and J.M. Thompson. 1995. Ecodistricts of Alberta: Summary of Biophysical

Attributes. Alberta Environmental Protection, Resource Data Division. Edmonton, Alta. Pub. no. T/319. 91pp.

Strong, W.L. and K.R. Leggat. 1992. Ecoregions of Alberta. Alberta Forestry, Lands and Wildlife, Resource Information Branch, Edmonton, Alta. T/245. 77pp.

Wilkinson, K. 1990. Trees and shrubs of Alberta. Lone Pine Publishing. Edmonton. Alta. 191pp.

Willoughby, M.G. 1992. Rangeland Reference Areas, Plant communities, ecology and response to grazing in Division 3. Forestry, Lands and Wildlife, Alberta Forest Service, Edmonton, Alta. Pub. no. T/269. 36pp.

Willoughby, M.G. 1992. Rangeland Reference Areas, Plant communities, ecology and response to grazing in Division 2. Forestry, Lands and Wildlife, Alberta Forest Service, Edmonton, Alta. Pub. no. T/268. 42pp.

Willoughby, M.G. 1998. Rangeland Reference Areas, Seven mile creek range condition and trend from 1953-1994. Environmental Protection, Lands and Forest Service. Edmonton, Alta. Pub. No. T/403. 21 pp.

Willoughby, M.G. 2001. Range plant community types and carrying capacity for the Upper Foothills subregion . 3<sup>rd</sup> Approximation. Sustainable Resource Development. Public Lands Division, Alta. Pub. no. T/003 120pp.

Willoughby, M.G., M.J. Alexander and B.W. Adams. 2003. Range plant communities and carrying capacity for the Montane subregion. 5<sup>th</sup> Approximation. Sustainable Resource Development. Public Lands Division. Pub. No. T/033. 223pp.











3 3286 52548021 2